

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 498 491 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

19.01.2005 Bulletin 2005/03

(21) Application number: **03723098.4**

(22) Date of filing: **09.04.2003**

(51) Int Cl.7: **C12P 21/08**, C12N 15/09,
C12N 5/10, C07K 16/18,
G01N 33/50, C07K 16/46
// (A61K39/395, A61P9:00,
29:00, 31:04, 31:12, 31:14,
35:00, 37:02, 37:04, 37:08,
43:00, G01N33:15,
C12P21:08), C12R1:91,
C12N5:10, C12R1:91

(86) International application number:

PCT/JP2003/004504

(87) International publication number:

WO 2003/085119 (16.10.2003 Gazette 2003/42)

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**

Designated Extension States:

AL LT LV MK

(30) Priority: **09.04.2002 JP 2002106950**

(71) Applicant: **KYOWA HAKKO KOGYO CO., LTD.**
Tokyo 100-8185 (JP)

(72) Inventors:

- **NAKAMURA, Kazuyasu,**
Kyowa Hakko Kogyo Co., Ltd.
3-chome, Machida-shi, Tokyo 194-8533 (US)
- **SHITARA, Kenya, Kyowa Hakko Kogyo Co., Ltd.**
3-chome, Machida-shi, Tokyo 194-8533 (JP)

(74) Representative: **VOSSIUS & PARTNER**

Siebertstrasse 4
81675 München (DE)

(54) **METHOD OF ENHANCING ACTIVITY OF ANTIBODY COMPOSITION OF BINDING TO Fc
GAMMA RECEPTOR IIIa**

(57) A method for enhancing a binding activity of an antibody composition to Fcγ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule; a method for enhancing an antibody-dependent cell-mediated cytotoxic activity of an antibody composition; a process for producing an antibody composition having an enhanced binding activity to Fcγ receptor IIIa; a method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the

reducing end in the sugar chain among total complex *N*-glycoside-linked sugar chains bound to the Fc region in an antibody composition; an Fc fusion protein composition produced by using a cell resistant to a lectin which recognizes a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond in a complex *N*-glycoside-linked sugar chain; and a process for producing the same.

EP 1 498 491 A1

Description

TECHNICAL FIELD

[0001] The present invention relates to a method for enhancing a binding activity of an antibody composition to Fc γ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule; a method for enhancing an antibody-dependent cell-mediated cytotoxic activity of an antibody composition; a process for producing an antibody composition having an enhanced binding activity to Fc γ receptor IIIa; a method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among total complex *N*-glycoside-linked sugar chains bound to the Fc region in an antibody composition; an Fc fusion protein composition produced by using a cell resistant to a lectin which recognizes a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain; and a process for producing the same.

BACKGROUND ART

[0002] Since antibodies have high binding activity, binding specificity and high stability in blood, their applications to diagnosis, prevention and treatment of various human diseases have been attempted [*Monoclonal Antibodies: Principles and Applications*, Wiley-Liss, Inc., Chapter 2.1 (1995)]. Also, production of a humanized antibody such as a human chimeric antibody or a human complementarity determining region (hereinafter referred to as "CDR")-grafted antibody from a non-human animal antibody have been attempted by using genetic recombination techniques. The human chimeric antibody is an antibody in which its antibody variable region (hereinafter referred to as "V region") is derived from a non-human animal antibody and its constant region (hereinafter referred to as "C region") is derived from a human antibody. The human CDR-grafted antibody is an antibody in which the CDR of a human antibody is replaced by CDR derived from a non-human animal antibody.

[0003] It has been revealed that five classes, namely IgM, IgD, IgG, IgA and IgE, are present in mammal antibodies. Antibodies of human IgG class are mainly used for the diagnosis, prevention and treatment of various human diseases because they have functional characteristics such as long half-life in blood and various effector functions [*Monoclonal Antibodies: Principles and Applications*, Wiley-Liss, Inc., Chapter 1 (1995)]. The human IgG class antibody is further classified into the following 4 subclasses: IgG1, IgG2, IgG3 and IgG4. A large number of studies have so far been conducted for antibody-dependent cell-mediated cytotoxic activity (hereinafter referred to as "ADCC activity") and complement-dependent cytotoxic activity (hereinafter referred to as "CDC activity") as effector functions of the IgG class antibody, and it has been reported that among antibodies of the human IgG class, the IgG1 subclass has the highest ADCC activity and CDC activity [*Chemical Immunology*, 65, 88 (1997)]. In view of the above, most of the anti-tumor humanized antibodies, including commercially available Rituxan and Herceptin, which require high effector functions for the expression of their effects, are antibodies of the human IgG1 subclass.

[0004] Expression of ADCC activity and CDC activity of the human IgG1 subclass antibodies requires binding of the Fc region of the antibody to a receptor for an antibody (hereinafter referred to as "Fc γ R") existing on the surface of effector cells such as killer cells, natural killer cells or activated macrophages and various complement components are bound. Regarding the binding, it has been suggested that several amino acid residues in the hinge region and the second domain of C region (hereinafter referred to as "C γ 2 domain") of the antibody are important [*Eur. J. Immunol.*, 23, 1098 (1993), *Immunology*, 86, 319 (1995), *Chemical Immunology*, 65, 88 (1997)] and that a sugar chain in the C γ 2 domain [*Chemical Immunology*, 65, 88 (1997)] is also important.

[0005] Regarding the sugar chain, Boyd *et al.* have examined effects of a sugar chain on the ADCC activity and CDC activity by treating a human CDR-grafted antibody CAMPATH-1H (human IgG1 subclass) produced by a Chinese hamster ovary cell (hereinafter referred to as "CHO cell") or a mouse myeloma NSO cell (hereinafter referred to as "NSO cell") with various glycosidases, and reported that elimination of sialic acid in the non-reducing end did not have influence upon both activities, but the CDC activity alone was affected by further removal of galactose residue and about 50% of the activity was decreased, and that complete removal of the sugar chain caused disappearance of both activities [*Molecular Immunol.*, 32, 1311 (1995)]. Also, Lively *et al.* have analyzed the sugar chain bound to a human CDR-grafted antibody CAMPATH-1H (human IgG1 subclass) which was produced by CHO cell, NSO cell or rat myeloma Y0 cell (hereinafter referred to as "Y0 cell"), measured its ADCC activity, and reported that the CAMPATH-1H derived from Y0 cell showed the highest ADCC activity, suggesting that *N*-acetylglucosamine (hereinafter sometimes referred to as "GlcNAc") at the bisecting position is important for the activity [*Glycobiology*, 5, 813 (1995); WO99/54342]. These reports indicate that the structure of the sugar chain plays an important role in the effector functions of human antibodies of IgG1 subclass and that it is possible to prepare an antibody having much higher effector function by modifying the structure of the sugar chain. However, structures of sugar chains are various and complex actually, and it cannot be said that an important structure for the effector function was completely identified.

[0006] Thus, the sugar chain bound to the CH2 domain of an IgG class antibody has great influence on the induction of effector functions of an antibody. As described above, some of effector functions of an antibody are exerted via interaction with FcγR present on the effector cell surface [*Annu. Rev. Immunol.*, 18, 709 (2000), *Annu. Rev. Immunol.*, 19, 275 (2001)].

[0007] It has been found that 3 different types are present in FcγR, and they are respectively called FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16). In human, FcγRII and FcγRIII are further classified into FcγRIIa and FcγRIIb, and FcγRIIIa and FcγRIIIb, respectively. FcγR is a membrane protein belonging to the immunoglobulin super family, FcγRII and FcγRIII have an α chain having an extracellular region containing two immunoglobulin-like domains, FcγRI has an α chain having an extracellular region containing three immunoglobulin-like domains, as a constituting component, and the α chain is involved in the IgG binding activity. In addition, FcγRI and FcγRIII have a γ chain or ζ chain as a constituting component which has a signal transduction function in association with the α chain [*Annu. Rev. Immunol.*, 18, 709 (2000), *Annu. Rev. Immunol.*, 19, 275 (2001)].

[0008] FcγRI is a high affinity receptor having a binding constant (hereinafter referred to as "K_A") value of 10⁸ to 10⁹ M⁻¹ and also has high binding activity for monomeric IgG [*Ann. Hematol.*, 76, 231 (1998)]. On the other hand, FcγRII and FcγRIII are low affinity receptors which have a low K_A value of 10⁵ to 10⁷ M⁻¹ for monomeric IgG and efficiently bind to an IgG immune complex polymerized by binding to an antigen or the like [*Ann. Hematol.*, 76, 231 (1998)]. Based on its functions, FcγR is classified into an activating receptor and an inhibitory receptor [*Annu. Rev. Immunol.*, 19, 275 (2001)].

[0009] In the activating receptor, there is a sequence consisting of 19 amino acid residues, called immunoreceptor tyrosine-based activation motif (hereinafter referred to as "ITAM") in the intracellular region of the α chain or the associating γ chain or ζ chain. According to the binding of an IgG immune complex, a tyrosine kinase such as Src or Syk which interacts with ITAM is activated to induce various activation reactions.

[0010] In the inhibitory receptor, there is a sequence consisting of 13 amino acid residues, called immunoreceptor tyrosine-based inhibitory motif (hereinafter referred to as "ITIM") in the intracellular region of the α chain. When ITIM is phosphorylated via its association with the activating receptor, various reactions including activation of a phosphatase called SHIP are induced to suppress activation signal from the activation receptor.

[0011] In human, the high affinity FcγRI and the low affinity FcγRIIa and FcγRIIIa function as activating receptors. In FcγRI, an ITAM sequence is present in the intracellular region of the associated γ chain. FcγRI is expressed on macrophages, monocytes, dendritic cells, neutrophils, eosinophils and the like. FcγRIIa comprises a single α chain, and an ITAM-like sequence is present in the intracellular region. FcγRIIa is expressed on macrophages, mast cells, monocytes, dendritic cells, Langerhans cells, neutrophils, eosinophils, platelets and a part of B cells. FcγRIIIa has an ITAM sequence present in the intracellular region of the associated γ chain or ζ chain and is expressed on NK cells, macrophages, monocytes, mast cells, dendritic cells, Langerhans cells, eosinophils and the like, but is not expressed on neutrophils, B cells and T cells.

[0012] On the other hand, the low affinity receptor FcγRIIb comprises a single α chain, and the amino acid sequence in the extracellular region has homology of about 90% with FcγRIIa. However, since an ITIM sequence is present in the intracellular region, it functions as a suppressing receptor. FcγRIIb is expressed on B cells, macrophages, mast cells, monocytes, dendritic cells, Langerhans cells, basophils, neutrophils and eosinophils, but is not expressed on NK cells and T cells. FcγRIIIb comprises a single α chain, and the amino acid sequence in the extracellular region has a homology of about 95% with FcγRIIIa. However, it is specifically expressed on neutrophils as a glycosylphosphatidylinositol (hereinafter referred to as "GPI")-anchored membrane protein. FcγRIIIb binds to an IgG immune complex but cannot activate cells by itself, and it is considered to function via its association with a receptor having an ITAM sequence such as FcγRIIa. Thus, *in vivo* effector functions of IgG class antibodies are obtained as the result of complex interaction with activating and suppressing FcγRs expressed on various effector cells.

[0013] It is considered that ADCC activity as one of the effector functions of IgG class antibodies is generated as a result of activation of effector cells such as NK cells, neutrophils, monocytes and macrophages, and among these, NK cells play an important role [*Blood*, 76, 2421 (1990), *Trends in Immunol.*, 22, 633 (2001), *Int. Rev. Immunol.*, 20, 503 (2001)].

[0014] FcγR expressed on NK cells is FcγRIIIa. Accordingly, it is considered that the ADCC activity can be enhanced by enhancing the activation signal from FcγRIIIa expressed on the NK cells.

[0015] As Fc fusion protein, Etanercept (trade name: Enbrel, manufactured by Immunex) (USP 5,605,690) and Alefacept (trade name: Amevive, manufactured by Biogen) (USP 5,914,111) and the like are known. Also, it is known that it has no ADCC activity when CH2 domain of an antibody is absent.

DISCLOSURE OF THE INVENTION

[0016] The present invention relates to the following (I) to (48).

(1) A method for enhancing a binding activity of an antibody composition to Fc γ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule.

(2) The method according to (1), wherein the modification of a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule is to bind a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain to the Fc region in the antibody molecule.

(3) The method according to (1) or (2), wherein the sugar chain is synthesized by a cell in which the activity of a protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is decreased or deleted.

(4) The method according to (3), wherein the protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is selected from the group consisting of the following (a), (b) and (c):

(a) a protein relating to synthesis of an intracellular sugar nucleotide, GDP-fucose;

(b) a protein relating to modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain;

(c) a protein relating to transport of an intracellular sugar nucleotide, GDP-fucose to the Golgi body.

(5) The method according to (3) or (4), wherein the cell is resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain.

(6) The method according to any one of (3) to (5), wherein the cell is resistant to at least one lectin selected from the group consisting of the following (a) to (d):

(a) a *Lens culinaris* lectin;

(b) a *Pisum sativum* lectin;

(c) a *Vicia faba* lectin;

(d) an *Aleuria aurantia* lectin.

(7) The method according to any one of (3) to (6), wherein the cell is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.

(8) The method according to any one of (3) to (7), wherein the cell is selected from the group consisting of the following (a) to (i):

(a) a CHO cell derived from a Chinese hamster ovary tissue;

(b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 cell;

(c) a BHK cell derived from a Syrian hamster kidney tissue;

(d) a mouse myeloma cell line NS0 cell;

(e) a mouse myeloma cell line SP2/0-Ag14 cell;

(f) a hybridoma cell;

(g) a human leukemic cell line Namalwa cell;

(h) an embryonic stem cell;

(i) a fertilized egg cell.

(9) The method according to any one of (1) to (8), wherein the antibody molecule is selected from the group consisting of the following (a) to (d):

(a) a human antibody;

(b) a humanized antibody;

(c) an antibody fragment comprising the Fc region of (a) or (b);

(d) a fusion protein comprising the Fc region of (a) or (b).

(10) The method according to any one of (1) to (9), wherein the antibody molecule belongs to an IgG class.

(11) The method according to any one of (1) to (10), wherein, in the complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule, the ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is 20% or more of total complex *N*-glycoside-linked sugar chains.

(12) A method for enhancing an antibody-dependent cell-mediated cytotoxic activity of an antibody composition,

which comprises using the method according to any one of (1) to (11).

(13) A process for producing an antibody composition having an enhanced binding activity to Fc γ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule.

(14) The process according to (13), wherein the modification of a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule is to bin a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain to the Fc region in the antibody molecule.

(15) The process according to (13) or (14), wherein the sugar chain is synthesized by a cell in which the activity of a protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is decreased or deleted.

(16) The process according to (15), wherein the protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is selected from the group consisting of the following (a), (b) and (c):

- (a) a protein relating to synthesis of an intracellular sugar nucleotide, GDP-fucose;
- (b) a protein relating to modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain;
- (c) a protein relating to transport of an intracellular sugar nucleotide, GDP-fucose to the Golgi body.

(17) The process according to (15) or (16), wherein the cell is resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain.

(18) The process according to any one of (15) to (17), wherein the cell is resistant to at least one lectin selected from the group consisting of the following (a) to (d):

- (a) a *Lens culinaris* lectin;
- (b) a *Pisum sativum* lectin;
- (c) a *Vicia faba* lectin;
- (d) an *Aleuria aurantia* lectin.

(19) The process according to any one of (15) to (18), wherein the cell is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.

(20) The process according to any one of (15) to (19), wherein the cell is selected from the group consisting of the following (a) to (i):

- (a) a CHO cell derived from a Chinese hamster ovary tissue;
- (b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 cell;
- (c) a BHK cell derived from a Syrian hamster kidney tissue;
- (d) a mouse myeloma cell line NSO cell;
- (e) a mouse myeloma cell line SP2/0-Ag14 cell;
- (f) a hybridoma cell;
- (g) a human leukemic cell line Namalwa cell;
- (h) an embryonic stem cell;
- (i) a fertilized egg cell.

(21) The process according to any one of (13) to (20), wherein the antibody molecule is selected from the group consisting of the following (a) to (d):

- (a) a human antibody;
- (b) a humanized antibody;
- (c) an antibody fragment comprising the Fc region of (a) or (b);
- (d) a fusion protein comprising the Fc region of (a) or (b).

(22) The process according to any one of (13) to (21), wherein the antibody molecule belongs to an IgG class.

(23) The method according to any one of (13) to (22), wherein, in the complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule, the ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is 20% or more of total

complex *N*-glycoside-linked sugar chains.

(24) A process for producing an antibody composition having an increased antibody-dependent cell-mediated cytotoxic activity, which comprises using the process according to (12).

(25) An antibody composition produced by the process according to any one of (13) to (24).

(26) A method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among total complex *N*-glycoside-linked sugar chains bound to the Fc region in an antibody composition, which comprises: reacting an antigen with a tested antibody composition to form a complex of the antigen and the antibody composition; contacting the complex with an Fc γ receptor IIIa to measure the binding activity to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

(27) A method for detecting the antibody-dependent cell-mediated cytotoxic activity in an antibody composition, which comprises: reacting an antigen with a tested antibody composition to form a complex of the antigen and the antibody composition; contacting the complex with an Fc γ receptor IIIa to measure the binding activity to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

(28) A method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among total complex *N*-glycoside-linked sugar chains bound to the Fc region in an antibody composition, which comprises: contacting a tested antibody composition with a Fc γ receptor IIIa to measure the binding activity of the antibody composition to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

(29) A method for detecting the antibody-dependent cell-mediated cytotoxic activity in an antibody composition, which comprises: contacting a tested antibody composition with a Fc γ receptor IIIa to measure the binding activity of the antibody composition to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

(30) An Fc fusion protein composition produced by using a cell resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain.

(31) The Fc fusion protein composition according to (30), wherein the cell is selected from the group consisting of the following (a), (b) and (c):

- (a) an enzyme protein relating to synthesis of an intracellular sugar nucleotide, GDP-fucose;
- (b) an enzyme protein relating to modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain;
- (c) a protein relating to transport of an intracellular sugar nucleotide, GDP-fucose to the Golgi body,

wherein the activity of the protein is decreased or deleted.

(32) The Fc fusion protein composition according to (30) or (31), wherein the cell is resistant to at least one lectin selected from the group consisting of the following (a) to (d):

- (a) a *Lens culinaris* lectin;
- (b) a *Pisum sativum* lectin;
- (c) a *Vicia faba* lectin;
- (d) an *Aleuria aurantia* lectin.

(33) The Fc fusion protein composition according to any one of (30) to (32), wherein the cell is a cell into which a gene encoding an Fc fusion protein is introduced.

(34) The Fc fusion protein composition according to (33), wherein the Fc is derived from an IgG class of an antibody molecule.

(35) The Fc fusion protein composition according to any one of (30) to (34), wherein the cell is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.

(36) The Fc fusion protein composition according to any one of (30) to (35), wherein the cell is a mouse myeloma cell.

(37) The Fc fusion protein composition according to (36), wherein the mouse myeloma cell is NS0 cell or SP2/0-Ag14 cell.

(38) The Fc fusion protein composition according to any one of (30) to (37), wherein the cell is selected from the group consisting of the following (a) to (g):

- (a) a CHO cell derived from a Chinese hamster ovary tissue;

- (b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 line;
- (c) a BHK cell derived from a Syrian hamster kidney tissue;
- (d) an antibody-producing hybridoma cell;
- (e) a human leukemic cell line Namalwa cell;
- (f) an embryonic stem cell;
- (g) a fertilized egg cell.

(39) An Fc fusion protein composition comprising an Fc fusion protein having an complex *N*-glycoside-linked sugar chain at the Fc region of an antibody molecule, wherein the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain is 20% or more of total complex *N*-glycoside-linked sugar chains which are bound to the Fc region in the composition.

(40) The Fc fusion protein composition according to (39), wherein the sugar chain in which fucose is not bound is a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain through α -bond.

(41) The Fc fusion protein composition according to (39) or (40), wherein the antibody molecule belongs to an IgG class.

(42) The Fc fusion protein composition according to any one of (30) to (41), wherein the Fc fusion protein composition is Fc-fused fibroblast growth factor-8.

(43) A cell which produces the Fc fusion protein composition according to any one of (30) to (42).

(44) The cell according to (43), which is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.

(45) The cell according to (43) or (44), which is a mouse myeloma cell.

(46) The cell according to (45), wherein the mouse myeloma cell is NSO cell or SP2/0-Ag14 cell.

(47) The cell according to any one of (43) to (46), which is selected from the group consisting of the following (a) to (g):

- (a) a CHO cell derived from a Chinese hamster ovary tissue;
- (b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 line;
- (c) a BHK cell derived from a Syrian hamster kidney tissue;
- (d) an antibody-producing hybridoma cell;
- (e) a human leukemic cell line Namalwa cell;
- (f) an embryonic stem cell;
- (g) a fertilized egg cell.

(48) A process for producing an Fc fusion protein composition, which comprises culturing the cell according to any one of (43) to (47) in a medium to form and accumulate an Fc fusion protein composition in the culture, and recovering the Fc fusion protein composition from the culture.

DETAILED DESCRIPTION OF THE INVENTION

[0017] The present invention relates to a method for enhancing a binding activity of an antibody composition to Fc γ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule.

[0018] In the present invention, the antibody molecule may be any antibody molecule, so long as it is a molecule comprising the Fc region of an antibody. Examples include an antibody, an antibody fragment, a fusion protein comprising an Fc region, and the like.

[0019] An antibody is a protein, which is *produced in vivo* by immunization as the result of extra-antigen stimulation. The antibody has a specific binding activity to an antigen. The antibody includes an antibody secreted by a hybridoma cell prepared from a spleen cell of an animal immunized with an antigen; an antibody prepared by genetic engineering technique, i.e., an antibody obtained by introducing an antibody gene-inserted antibody expression vector into a host cell; and the like. Examples include an antibody produced by a hybridoma, a humanized antibody, a human antibody and the like.

[0020] A hybridoma is a cell which is obtained by cell fusion between a B cell obtained by immunizing a non-human mammal with an antigen and a myeloma cell derived from mouse or the like and can produce a monoclonal antibody having the desired antigen specificity.

[0021] The humanized antibody includes a human chimeric antibody, a human CDR-grafted antibody and the like.

[0022] A human chimeric antibody is an antibody which comprises an antibody heavy chain V region (hereinafter referred to heavy chain as "H chain", and referred to as "HV" or "VH") and an antibody light chain V region (hereinafter

referred to light chain as "L chain", and referred to as "LV" or "VL"), both of a non-human animal, a human antibody H chain C region (hereinafter also referred to as "CH") and a human antibody L chain C region (hereinafter also referred to as "CL"). The non-human animal may be any animal such as mouse, rat, hamster or rabbit, so long as a hybridoma can be prepared therefrom.

[0023] The human chimeric antibody can be produced by obtaining cDNAs encoding VH and VL from a monoclonal antibody-producing hybridoma, inserting them into an expression vector for host cell having genes encoding human antibody CH and human antibody CL to thereby construct a vector for expression of human chimeric antibody, and then introducing the vector into a host cell to express the antibody.

[0024] The CH of human chimeric antibody may be any CH, so long as it belongs to human immunoglobulin (hereinafter referred to as "hlg") can be used. Those belonging to the hlgG class are preferred and any one of the subclasses belonging to the hlgG class such as hlgG1, hlgG2, hlgG3 and hlgG4 can be used. Also, as the CL of human chimeric antibody, any CL can be used, so long as it belongs to the hlg class, and those belonging to the κ class or λ class can also be used.

[0025] A human CDR-grafted antibody is an antibody in which amino acid sequences of CDRs of VH and VL of a non-human animal antibody are grafted into appropriate positions of VH and VL of a human antibody.

[0026] The human CDR-grafted antibody can be produced by constructing cDNAs encoding V regions in which CDR sequences of VH and VL of a non-human animal antibody are grafted into CDR sequences of VH and VL of a desired human antibody, inserting them into an expression vector for host cell having genes encoding human antibody CH and human antibody CL to thereby construct a vector for expression of human CDR-grafted antibody, and then introducing the expression vector into a host cell to express the human CDR-grafted antibody.

[0027] The CH of human CDR-grafted antibody may be any CH, so long as it belongs to the hlg. Those of the hlgG class are preferred and any one of the subclasses belonging to the hlgG class, such as hlgG1, hlgG2, hlgG3 and hlgG4, can be used. Also, as the CL of human CDR-grafted antibody, any CL can be used, so long as it belongs to the hlg class, and those belonging to the κ class or λ class can also be used.

[0028] A human antibody is originally an antibody naturally existing in the human body, but it also includes antibodies obtained from a human antibody phage library, a human antibody-producing transgenic non-human animal and a human antibody-producing transgenic plant, which are prepared based on the recent advance in genetic engineering, cell engineering and developmental engineering techniques.

[0029] Regarding the antibody existing in the human body, a lymphocyte capable of producing the antibody can be cultured by isolating a human peripheral blood lymphocyte, immortalizing it by its infection with EB virus or the like and then cloning it, and the antibody can be purified from the culture.

[0030] The human antibody phage library is a library in which antibody fragments such as Fab and single chain antibody are expressed on the phage surface by inserting a gene encoding an antibody prepared from a human B cell into a phage gene: A phage expressing an antibody fragment having the desired antigen binding activity can be recovered from the library based on the activity to bind to an antigen-immobilized substrate. The antibody fragment can be converted further into a human antibody molecule comprising two full H chains and two full L chains by genetic engineering techniques.

[0031] A human antibody-producing transgenic non-human animal is an animal in which a human antibody gene is introduced into cells. Specifically, a human antibody-producing transgenic mouse can be prepared by introducing a human antibody gene into ES cell of a mouse, transplanting the ES cell into an early stage embryo of other mouse and then developing it. By introducing a human antibody-gene into a fertilized egg of an animal and developing it, the transgenic non-human animal can be also prepared. Regarding the preparation method of a human antibody from the human antibody-producing transgenic non-human animal, the human antibody can be produced and accumulated in a culture by obtaining a human antibody-producing hybridoma by a hybridoma preparation method usually carried out in non-human mammals and then culturing it.

[0032] The transgenic non-human animal includes cattle, sheep, goat, pig, horse, mouse, rat, fowl, monkey, rabbit and the like.

[0033] Moreover, in the present invention, the antibody is preferably an antibody which recognizes a tumor-related antigen, an antibody which recognizes an allergy- or inflammation-related antigen, an antibody which recognizes cardiovascular disease-related antigen, an antibody which recognizes autoimmune disease-related antigen or an antibody which recognizes a viral or bacterial infection-related antigen. Also, the class of the antibody is preferably IgG.

[0034] An antibody fragment is a fragment which comprises at least part of the Fc region of the above antibody. The Fc region is a region at the C-terminal side of H chain of an antibody, such as CH2 region and CH3 region, and includes a natural type and a mutant type. The at least part of the Fc region is preferably a fragment comprising CH2 region, and more preferably a region comprising aspartic acid at position 1 existing in CH2 region. The Fc region of the IgG class is from Cys at position 226 to the C-terminal or from Pro at position 230 to the C-terminal according to the numbering of EU Index of Kabat *et al.* [*Sequences of Proteins of Immunological Interest*, 5th Ed., Public Health Service, National Institutes of Health, Bethesda, MD. (1991)]. The antibody fragment includes an H chain monomer, an H chain

dimer and the like.

[0035] A fusion protein comprising a part of Fc region is a substance in which an antibody comprising the part of Fc region of an antibody or the antibody fragment is fused with a protein such as an enzyme or a cytokine (hereinafter referred to as "Fc fusion protein").

[0036] In the present invention, *N*-glycoside-linked sugar chain bound to the Fc region of the antibody molecule includes a complex type in which the non-reducing end side of the core structure has one or more parallel branches of galactose-*N*-acetylglucosamine (hereinafter referred to as "Gal-GlcNAc") and further the non-reducing end side of Gal-GlcNAc has a structure of sialic acid, bisecting *N*-acetylglucosamine or the like.

[0037] Since the Fc region in the antibody molecule has positions to which *N*-glycoside-linked sugar chains are separately bound, two sugar chains are bound per one antibody molecule. Since many sugar chains having different structures are present for the two *N*-glycoside-linked sugar chains bound to the antibody, homology of antibody molecules can be judged in view of the sugar chain structure bound to the Fc region.

[0038] The antibody composition is a composition which comprises an antibody molecule having complex *N*-glycoside-linked sugar chains in the Fc region, and may comprise an antibody molecule having the same sugar chain structure or an antibody molecule having different sugar chain structures.

[0039] Modification of the *N*-glycoside-linked sugar chain bound to the Fc region of an antibody molecule is preferably carried out by binding a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain to the Fc region of an antibody molecule.

[0040] In the present invention, the sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is a complex *N*-glycoside-linked sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

[0041] The sugar chain can be synthesized by a cell in which the activity of an enzyme protein relating to the modification of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is decreased or deleted.

[0042] In the present invention, the enzyme protein relating to the modification of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain includes:

- (a) an enzyme protein relating to synthesis of an intracellular sugar nucleotide, GDP-fucose (hereinafter referred to as "GDP-fucose synthase");
- (b) an enzyme protein relating to modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain (hereinafter referred to as " α 1,6-fucose modifying enzyme"); and
- (c) a protein relating to transport of an intracellular sugar nucleotide, GDP-fucose, to the Golgi body (hereinafter referred to as "GDP-fucose transport protein").

[0043] In the present invention, the GDP-fucose synthase may be any enzyme, so long as it is an enzyme relating to the synthesis of the intracellular sugar nucleotide, GDP-fucose, as a supply source of fucose to a sugar chain, and includes an enzyme which has influence on the synthesis of the intracellular sugar nucleotide, GDP-fucose.

[0044] The intracellular sugar nucleotide, GDP-fucose, is supplied by a *de novo* synthesis pathway or a salvage synthesis pathway. Thus, all enzymes relating to the synthesis pathways are included in the GDP-fucose synthase.

[0045] The GDP-fucose synthase relating to the *de novo* synthesis pathway includes GDP-mannose 4-dehydratase (hereinafter referred to as "GMD"), GDP-keto-6-deoxymannose 3,5-epimerase, 4-reductase (hereinafter referred to as "Fx") and the like.

[0046] The GDP-fucose synthase relating to the salvage synthesis pathway includes GDP-beta-L-fucose pyrophosphorylase (hereinafter referred to as "GFPP"), fucokinase and the like.

[0047] As the enzyme which has influence on the synthesis of an intracellular sugar nucleotide, GDP-fucose, an enzyme which has influence on the activity of the enzyme relating to the synthesis pathway of the intracellular sugar nucleotide, GDP-fucose, and an enzyme which has influence on the structure of substances as the substrate of the enzyme are also included.

[0048] In the present invention, the GMD includes a protein encoded by a DNA selected from the group consisting of (a) and (b), a protein selected from the group consisting of (c), (d) and (e), and the like:

- (a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:65;
- (b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:65 under stringent conditions and encodes a protein having GMD activity,
- (c) a protein comprising the amino acid sequence represented by SEQ ID NO:71,

- (d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:71 and has GMD activity, and
 (e) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:71 and has GMD activity.

[0049] Fx includes a protein encoded by a DNA selected from the group consisting of (a) and (b), a protein selected from the group consisting of (c), (d) and (e), and the like:

- (a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:48;
 (b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:48 under stringent conditions and encodes a protein having Fx activity,
 (c) a protein comprising the amino acid sequence represented by SEQ ID NO:19,
 (d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:19 and has Fx activity, and
 (e) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:19 and has Fx activity.

[0050] GFPP includes a protein encoded by a DNA selected from the group consisting of (a) and (b), a protein selected from the group consisting of (c), (d) and (e), and the like:

- (a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:51;
 (b) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:51 under stringent conditions and encodes a protein having GFPP activity,
 (c) a protein comprising the amino acid sequence represented by SEQ ID NO:20,
 (d) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:20 and has GFPP activity, and
 (e) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:20 and has GFPP activity.

[0051] In the present invention, the α 1,6-fucose modifying enzyme includes any enzyme, so long as it is an enzyme relating to the reaction of binding of 1-position of fucose to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain. The enzyme relating to the reaction of binding of 1-position of fucose to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain includes an enzyme which has influence on the reaction of binding of 1-position of fucose to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain.

[0052] The α 1,6-fucose modifying enzyme includes α 1,6-fucosyltransferase, α -L-fucosidase and the like.

[0053] Also, an enzyme which has influence on the activity of the above enzyme relating to the reaction of binding of 1-position of fucose to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain and an enzyme which has influence on the structure of substances as the substrate of the enzyme are included.

[0054] The α 1,6-fucosyltransferase includes a protein encoded by a DNA selected from the group consisting of the following (a), (b), (c) and (d), a protein selected from the group consisting of the following (e), (f), (g), (h), (i) and (j), and the like:

- (a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:1;
 (b) a DNA comprising the nucleotide sequence represented by SEQ ID NO:2;
 (c) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:1 under stringent conditions and encodes a protein having α 1,6-fucosyltransferase activity;
 (d) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:2 under stringent conditions and encodes a protein having α 1,6-fucosyltransferase activity;
 (e) a protein comprising the amino acid sequence represented by SEQ ID NO:23,
 (f) a protein comprising the amino acid sequence represented by SEQ ID NO:24,
 (g) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:23 and has α 1,6-fucosyltransferase activity,
 (h) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:24 and has α 1,6-fucosyltransferase activity,

- (i) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:23 and has α 1,6-fucosyltransferase activity, and
- (j) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:24 and has α 1,6-fucosyltransferase activity.

[0055] The GDP-fucose transport protein may be any protein, so long as it is a protein relating to the transportation of the intracellular sugar nucleotide, GDP-fucose to the Golgi body or a protein which has an influence on the reaction to transport the intracellular sugar nucleotide, GDP-fucose to the Golgi body.

[0056] The GDP-fucose transport protein includes a GDP-fucose transporter and the like.

[0057] Furthermore, the protein which has an influence on the reaction to transport the intracellular sugar nucleotide, GDP-fucose to the Golgi body includes a protein which has an influence on the above GDP-fucose transport protein or has an influence on the expression thereof.

[0058] In the present invention, the GDP-fucose transporter includes a protein encoded by a DNA selected from the group consisting of the following (a) to (h), and the like:

- (a) a DNA comprising the nucleotide sequence represented by SEQ ID NO:91;
- (b) a DNA comprising the nucleotide sequence represented by SEQ ID NO:93;
- (c) a DNA comprising the nucleotide sequence represented by SEQ ID NO:95;
- (d) a DNA comprising the nucleotide sequence represented by SEQ ID NO:97;
- (e) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:91 under stringent conditions and encodes a protein having GDP-fucose transporter activity;
- (f) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:93 under stringent conditions and encodes a protein having GDP-fucose transporter activity;
- (g) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:95 under stringent conditions and encodes a protein having GDP-fucose transporter activity; and
- (h) a DNA which hybridizes with the DNA comprising the nucleotide sequence represented by SEQ ID NO:97 under stringent conditions and encodes a protein having GDP-fucose transporter activity.

Furthermore, the GDP-fucose transporter of the present invention includes a protein selected from the group consisting of the following (i) to (t), and the like:

- (i) a protein comprising the amino acid sequence represented by SEQ ID NO:92,
- (j) a protein comprising the amino acid sequence represented by SEQ ID NO:94,
- (k) a protein comprising the amino acid sequence represented by SEQ ID NO:96,
- (l) a protein comprising the amino acid sequence represented by SEQ ID NO:98,
- (m) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:92 and has GDP-fucose transporter activity,
- (n) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:94 and has GDP-fucose transporter activity,
- (o) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:96 and has GDP-fucose transporter activity,
- (p) a protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:98 and has GDP-fucose transporter activity,
- (q) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:92 and has GDP-fucose transporter activity,
- (r) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:94 and has GDP-fucose transporter activity,
- (s) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:96 and has GDP-fucose transporter activity, and
- (t) a protein which comprises an amino acid sequence having a homology of 80% or more with the amino acid sequence represented by SEQ ID NO:98 and has GDP-fucose transporter activity.

[0059] A DNA which hybridizes under stringent conditions is a DNA obtained, e.g., by a method such as colony hybridization, plaque hybridization or Southern blot hybridization using a DNA such as the DNA having the nucleotide sequence represented by SEQ ID NO:1, 2, 48, 51, 65, 91, 93, 95 or 97 or a partial fragment thereof as the probe, and specifically includes a DNA which can be identified by carrying out hybridization at 65°C in the presence of 0.7 to 1.0

mol/l sodium chloride using a filter to which colony- or plaque-derived DNAs are immobilized, and then washing the filter at 65°C using 0.1 to 2 × SSC solution (composition of the 1 × SSC solution comprising 150 mmol/l sodium chloride and 15 mmol/l sodium citrate). The hybridization can be carried out in accordance with the methods described, e.g., in *Molecular Cloning, A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press (1989) (hereinafter referred to as "*Molecular Cloning, Second Edition*"); *Current Protocols in Molecular Biology*, John Wiley & Sons, 1987-1997 (hereinafter referred to as "*Current Protocols in Molecular Biology*"); *DNA Cloning 1: Core Techniques, A Practical Approach*, Second Edition, Oxford University (1995); and the like. The hybridizable DNA includes a DNA having at least 60% or more, preferably 70% or more, more preferably 80% or more, still more preferably 90% or more, far more preferably 95% or more, and most preferably 98% or more, of homology with the nucleotide sequence represented by SEQ ID NO:1, 2, 48, 51, 65, 91, 93, 95 or 97.

[0060] The protein which comprises an amino acid sequence in which at least one amino acid is deleted, substituted, inserted and/or added in the amino acid sequence represented by SEQ ID NO:19, 20, 23, 24, 71, 92, 94, 96 or 98 and has α1,6-fucosyltransferase activity, GMD activity, Fx activity, GFPP activity or GFP-fucose transporter activity can be obtained, e.g., by introducing a site-directed mutation into a DNA encoding a protein having the amino acid sequence represented by SEQ ID NO:1, 2, 48, 51 or 65, respectively, using the site-directed mutagenesis described, e.g., in *Molecular Cloning, Second Edition*; *Current Protocols in Molecular Biology*; *Nucleic Acids Research*, **10**, 6487 (1982); *Proc. Natl. Acad. Sci. USA*, **79**, 6409 (1982); *Gene*, **34**, 315 (1985); *Nucleic Acids Research*, **13**, 4431 (1985); *Proc. Natl. Acad. Sci. USA*, **82**, 488 (1985); and the like. The number of amino acids to be deleted, substituted, inserted and/or added is one or more, and the number is not particularly limited, but is a number which can be deleted, substituted or added by a known technique such as the site-directed mutagenesis, e.g., it is 1 to several tens, preferably 1 to 20, more preferably 1 to 10, and most preferably 1 to 5.

[0061] Also, in order to maintain the α1,6-fucosyltransferase activity, GMD activity, Fx activity, GFPP activity or GDP-fucose transporter activity of the protein to be used in the present invention, it has at least 80% or more, preferably 85% or more, more preferably 90% or more, still more preferably 95% or more, far more preferably 97% or more, and most preferably 99% or more, of homology with the amino acid sequence represented by SEQ ID NO:19, 20, 23, 24, 71, 92, 94, 96 or 98, when calculated using an analyzing soft such as BLAST [*J. Mol. Biol.*, **215**, 403 (1990)] or FASTA [*Methods in Enzymology*, **183**, 63 (1990)].

[0062] As the method for obtaining the above cells, any technique can be used, so long as it can decrease or delete the enzyme activity of interest. The technique for decreasing or deleting the enzyme activity includes:

- (a) a gene disruption technique which comprises targeting a gene encoding the enzyme,
- (b) a technique for introducing a dominant negative mutant of a gene encoding the enzyme,
- (c) a technique for introducing mutation into the enzyme, and
- (d) a technique for surprising transcription and/or translation of a gene encoding the enzyme, and the like.

[0063] Also, the method includes a method for selecting a cell having resistance to lectin which recognizes the structure of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond in a complex *N*-glycoside-linked sugar chain.

[0064] The growth of lectin-resistant cell is not inhibited in the presence of a lectin at an effective concentration during cell culturing.

[0065] In the present invention, the effective concentration of a lectin which does not inhibit the growth can be decided depending on the cell line, and is generally 10 μg/ml to 10.0 mg/ml, preferably 0.5 to 2.0 mg/ml. The effective concentration of lectin in the case where mutation is introduced into a parent cell is a concentration in which the parent cell cannot normally grow or higher than the concentration, and is a concentration which is preferably similar to, more preferably 2 to 5 times, still more preferably 10 times, and most preferably 20 times or more, higher than the concentration in which the parent cell cannot normally grow.

[0066] The lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine through α-bond includes any lectin, so long as it is a lectin which is capable of recognizing the sugar chain structure. Examples include lentil agglutinin derived from *Lens culinaris* (*Lens culinaris* lectin LCA), pea lectin derived from *Pisum sativum* (pea lectin PSA), (agglutinin derived from *Vicia faba* (broad bean lectin VFA), lectin derived from *Aleuria aurantia* (*Aleuria aurantia* lectin AAL) and the like.

[0067] The parent cell is a cell before a certain treatment is applied, namely a cell before the step for selecting the lectin-resistant cell used in the present invention is carried out or a cell before genetic engineering techniques for decreasing or deleting the above enzyme activity is carried out.

[0068] Although the parent cell is not particularly limited, the following cells are exemplified.

[0069] The parent cell of NSO cell includes NSO cells described in literatures such as *BIO/TECHNOLOGY*, **10**, 169 (1992) and *Biotechnol. Bioeng.*, **73**, 261 (2001). Furthermore, it includes NSO cell line (RCB 0213) registered at RIKEN Cell Bank, The Institute of Physical and Chemical Research, sub-cell lines obtained by acclimating these cell lines to

media in which they can grow, and the like.

[0070] The parent cell of SP2/0-Ag14 cell includes SP2/0-Ag14 cells described in literatures such as *J. Immunol.*, 126, 317 (1981), *Nature*, 276, 269 (1978) and *Human Antibodies and Hybridomas*, 3, 129 (1992). Furthermore, it includes SP2/0-Ag14 cell (ATCC CRL-1581) registered at ATCC, sub-cell lines obtained by naturalizing these cell lines to media in which they can grow (ATCC CRL-1581.1), and the like.

[0071] The parent cell of CHO cell derived from Chinese hamster ovary tissue includes CHO cells described in literatures such as *Journal of Experimental Medicine*, 108, 945 (1958), *Proc. Natl. Acad. Sci. USA*, 60, 1275 (1968), *Genetics*, 55, 513 (1968), *Chromosoma*, 41, 129 (1973), *Methods in Cell Science*, 18, 115 (1996), *Radiation Research*, 148, 260 (1997), *Proc. Natl. Acad. Sci. USA*, 77, 4216 (1980), *Proc. Natl. Acad. Sci. USA*, 60, 1275 (1968), *Cell*, 6, 121 (1975) and *Molecular Cell Genetics*, Appendix I, II (p. 883-900). Furthermore, it includes cell line CHO-K1 (ATCC CCL-61), cell line DUXB11 (ATCC CRL-9060) and cell line Pro-5 (ATCC CRL-1781) registered at ATCC, commercially available cell line CHO-S (Cat # 11619 of Life Technologies), sub-cell lines obtained by acclimating these cell lines to media in which they can grow, and the like.

[0072] The parent cell of a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 cell includes cell lines established from Y3/Ag1.2.3 cell (ATCC CRL-1631) such as YB2/3HL.P2.G11.16Ag.20 cell described in literatures such as *J. Cell. Biol.*, 93, 576 (1982) and *Methods Enzymol.*, 73B, 1 (1981). Furthermore, it include YB2/3HL.P2.G11.16Ag.20 cell (ATCC CRL-1662) registered at ATCC, sub-lines obtained by acclimating these cell lines to media in which they can grow, and the like.

[0073] In the present invention, Fc γ R means an Fc receptor (hereinafter also referred to as "FcR") against an IgG class antibody. FcR means a receptor which binds to the Fc region of an antibody [*Annu. Rev. Immunol.*, 9, 457 (1991)]. Furthermore, Fc γ R includes Fc γ RI, Fc γ RII and Fc γ RIII subclasses and their allele mutants and isoforms formed by alternative splicing. In addition, Fc γ RII includes Fc γ RIIa and Fc γ RIIb, and Fc γ RIII includes Fc γ RIIIa and Fc γ RIIIb [*Annu. Rev. Immunol.*, 9, 457 (1991)].

[0074] The binding activity to Fc γ RIIIa can be increased by binding sugar chains to the Fc region of an antibody molecule so as to adjust the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among the total complex *N*-glycoside-linked sugar chains bound to the Fc region to preferably 20% or more, more preferably 30% or more, still more preferably 40% or more, particularly preferably 50% or more, and most preferably 100%.

[0075] The ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among the total complex *N*-glycoside-linked sugar chains bound to the Fc region contained in the antibody composition is a ratio of the number of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain to the total number of the complex *N*-glycoside-linked sugar chains bound to the Fc region contained in the composition. Also, the ratio of a sugar chain is preferably a ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the sugar chain.

[0076] The sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain. Preferably, it is a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the complex *N*-glycoside-linked sugar chain through α -bond.

[0077] The ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain contained in the composition which comprises an antibody molecule having complex *N*-glycoside-linked sugar chains in the Fc region can be determined by releasing the sugar chain from the antibody molecule using a known method such as hydrazinolysis, enzyme digestion or the like [*Biochemical Experimentation Methods 23 - Method for Studying Glycoprotein Sugar Chain* (Japan Scientific Societies Press), edited by Reiko Takahashi (1989)], carrying out fluorescence labeling or radioisotope labeling of the released sugar chain, and then separating the labeled sugar chain by chromatography. Also, the released sugar chain can be determined by analyzing it with the HPAED-PAD method [*J. Liq. Chromatogr.*, 6, 1577 (1983)]. The antibody composition in which binding activity to Fc γ RIIIa has been enhanced by the method of the present invention has high ADCC activity.

[0078] In the present invention, the ADCC activity is a cytotoxic activity in which an antibody bound to a cell surface antigen on a cell such as a tumor cell in the living body activates an effector cell mediated the antibody Fc region and an Fc receptor existing on effector cell surface and thereby injures the tumor cell and the like [*Monoclonal Antibodies: Principles and Applications*, Wiley-Liss, Inc., Chapter 2.1 (1995)]. The effector cell includes killer cells, natural killer cells, monocytes, macrophages, and the like.

[0079] A process for producing a host cell in which the activity of a protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is decreased or deleted used in the method of the present invention is explained below in detail.

1. Preparation of host cell used in the method of the invention

[0080] The host cell used in the method of the present invention can be prepared by the following techniques.

(1) Gene disruption technique which comprises targeting a gene encoding an enzyme

[0081] The host cell used in the method of the present invention can be prepared by targeting a gene encoding a GDP-fucose synthase, α 1,6-fucose modifying enzyme or a GDP-fucose transport protein by using a gene disruption technique. The GDP-fucose synthase includes GMD, Fx, GFPP, fucokinase and the like. The α 1,6-fucose modifying enzyme includes α -1,6-fucosyltransferase, α -L-fucosidase and the like. The GDP-fucose transport protein includes GDP-fucose transporter.

[0082] The gene as used herein includes DNA and RNA.

[0083] The gene disruption method may be any method, so long as it can disrupt the gene encoding the target enzyme. Examples include an antisense method, a ribozyme method, a homologous recombination method, an RNA-DNA oligonucleotide (RDO) method, an RNA interference (RNAi) method, a method using retrovirus, a method using transposon and the like. The methods are specifically described below.

(a) Preparation of host cell used in the present invention by the antisense method or the ribozyme method

[0084] The host cell used in the method of the present invention can be prepared by targeting the GDP-fucose synthase, α 1,6-fucose modifying enzyme or the GDP-fucose transport protein according to the antisense or ribozyme method described in *Cell Technology*, 12, 239 (1993); *BIO/TECHNOLOGY*, 1.7 1097 (1999); *Hum. Mol. Genet.*, 5, 1083 (1995); *Cell Technology*, 13, 255 (1994); *Proc. Natl. Acad. Sci. USA*, 96, 1886 (1999); or the like, e.g., in the following manner.

[0085] A cDNA or a genomic DNA encoding GDP-fucose synthase, α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is prepared.

[0086] The nucleotide sequence of the prepared cDNA or genomic DNA is determined.

[0087] Based on the determined DNA sequence, an antisense gene or ribozyme construct of an appropriate length comprising a DNA moiety which encodes the GDP-fucose synthase, α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, a part of its untranslated region or an intron is designed.

[0088] In order to express the antisense gene or ribozyme in a cell, a recombinant vector is prepared by inserting a fragment or total length of the prepared DNA into downstream of the promoter of an appropriate expression vector.

[0089] A transformant is obtained by introducing the recombinant vector into a host cell suitable for the expression vector.

[0090] The host cell used in the method of the present invention can be obtained by selecting a transformant based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein. The host cell of the present invention can also be obtained by selecting a transformant based on the sugar chain structure of a glycoprotein on the cell membrane or the sugar chain structure of the produced antibody molecule.

[0091] As the host cell used for preparing the host cell used in the method of the present invention, any cell such as yeast, an animal cell, an insect cell or a plant cell can be used, so long as it has a gene encoding the target GDP-fucose synthase, α 1,6-fucose modifying enzyme or GDP-fucose transport protein. Examples include host cells described in the following item 3.

[0092] As the expression vector, a vector which is autonomously replicable in the host cell or can be integrated into the chromosome and comprises a promoter at such a position that the designed antisense gene or ribozyme can be transferred is used. Examples include expression vectors described in the following item 3.

[0093] As the method for introducing a gene into various host cells, the methods for introducing recombinant vectors suitable for various host cells described in the following item 3, can be used.

[0094] The method for selecting a transformant based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes biochemical methods or genetic engineering techniques described in *New Biochemical Experimentation Series (Shin-Jikken Kagaku Koza) 3 - Saccharides (Toshitsu) I*, Glycoprotein (Totanpakushitu) (Tokyo Kagaku Dojin), edited by Japanese Biochemical Society (1988); *Cell Engineering (Saibo Kogaku)*, Supplement, Experimental Protocol Series, Glycobiology Experimental Protocol, Glycoprotein, Glycolipid and Proteoglycan (Shujun-sha), edited by Naoyuki Taniguchi, Akemi Suzuki, Kiyoshi Furukawa and Kazuyuki Sugawara (1996); *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*; and the like. The biochemical method includes a method in which the enzyme activity is evaluated using an enzyme-specific substrate and the like. The genetic engineering technique include the Northern analysis, RT-PCR and the like which measures the amount of mRNA of a gene encoding the enzyme.

[0095] The method for selecting a transformant based on the sugar chain structure of a glycoprotein on the cell

membrane includes the methods described in the following item 1(5). The method for selecting a transformant based on the sugar chain structure of a produced antibody molecule includes the methods described in the following items 6 and 7.

[0096] As the method for preparing cDNA encoding the GDP-fucose synthase, α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, the following method is exemplified.

Preparation of DNA:

[0097] A total RNA or mRNA is prepared from a human or non-human animal tissue or cell.

[0098] The mRNA of a human or non-human tissue or cell may be a commercially available product (e.g., manufactured by Clontech) or may be prepared from a human or non-human animal tissue or cell as follows. The method for preparing a total RNA from a human or non-human animal tissue or cell includes the guanidine thiocyanate-caesium trifluoroacetate method [*Methods in Enzymology*, 154, 3 (1987)], the acidic guanidine thiocyanate phenol chloroform (AGPC) method [*Analytical Biochemistry*, 162, 156 (1987); *Experimental Medicine*, 9, 1937 (1991)] and the like.

[0099] Also, the mRNA can be prepared as poly(A)⁺ RNA from a total RNA by the oligo(dT)-immobilized cellulose column method (*Molecular Cloning*, Second Edition) and the like.

[0100] In addition, mRNA can be prepared using a kit such as Fast Track mRNA Isolation Kit (manufactured by Invitrogen) or Quick Prep mRNA Purification Kit (manufactured by Pharmacia).

[0101] A cDNA library is prepared from the prepared mRNA of a human or non-human animal tissue or cell. The method for preparing cDNA library includes the methods described in *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*; and the like, or methods using a commercially available kit such as Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by Life Technologies) or ZAP-cDNA Synthesis Kit (manufactured by STRATAGENE).

[0102] As the cloning vector for the preparation of the cDNA library, any vector such as a phage vector or a plasmid vector can be used, so long as it is autonomously replicable in *Escherichia coli* K12. Examples include ZAP Express [manufactured by STRATAGENE, *Strategies*, 5, 58 (1992)], pBluescript II SK(+) [*Nucleic Acids Research*, 17, 9494 (1989)], Lambda ZAP II (manufactured by STRATAGENE), λ gt10 and λ gt11 [DNA Cloning, A Practical Approach, 1, 49 (1985)], λ TriplEx (manufactured by Clontech), λ ExCell (manufactured by Pharmacia), pT7T318U (manufactured by Pharmacia), pcD2 [*Mol. Cell. Biol.*, 3, 280 (1983)], pUC18 [*Gene*, 33, 103 (1985)] and the like.

[0103] Any microorganism can be used as the host microorganism for the preparation of the cDNA library, and *Escherichia coli* is preferably used. Examples include *Escherichia coli* XL1-Blue MRF' [manufactured by STRATAGENE, *Strategies*, 5, 81 (1992)], *Escherichia coli* C600 [*Genetics*, 39 440 (1954)], *Escherichia coli* Y1088 [*Science*, 222, 778 (1983)], *Escherichia coli* Y1090 [*Science*, 222, 778 (1983)], *Escherichia coli* NM522 [*J. Mol. Biol.*, 166, 1 (1983)], *Escherichia coli* K802 [*J. Mol. Biol.*, 16, 118 (1966)], *Escherichia coli* JM105 [*Gene*, 38, 275 (1985)] and the like.

[0104] The cDNA library can be used as such in the following analysis, and in order to obtain a full length cDNA as efficient as possible by decreasing the ratio of an infull length cDNA, a cDNA library prepared by using the oligo cap method developed by Sugano *et al.* [*Gene*, 138, 171 (1994); *Gene*, 200, 149 (1997); *Protein, Nucleic Acid and Protein*, 41, 603 (1996); *Experimental Medicine*, 11, 2491 (1993); *cDNA Cloning* (Yodo-sha) (1996); *Methods for Preparing Gene Libraries* (Yodo-sha) (1994)] can be used in the following analysis.

[0105] Based on the amino acid sequence of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, degenerative primers specific for the 5'-terminal and 3'-terminal nucleotide sequences of a nucleotide sequence presumed to encode the amino acid sequence are prepared, and DNA is amplified by PCR [*PCR Protocols*, Academic Press (1990)] using the prepared cDNA library as the template to obtain a gene fragment encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein.

[0106] It can be confirmed that the obtained gene fragment is a DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein by a method generally used for analyzing a nucleotide such as the dideoxy method of Sanger *et al.* [*Proc. Natl. Acad. Sci. USA*, 74, 5463 (1977)] or by using a nucleotide sequence analyzer such as ABI PRISM 377 DNA Sequencer (manufactured by PE Biosystems).

[0107] A DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein can be obtained by carrying out colony hybridization or plaque hybridization (*Molecular Cloning*, Second Edition) for the cDNA or cDNA library synthesized from the mRNA contained in the human or non-human animal tissue or cell, using the gene fragment as a DNA probe.

[0108] Also, using the primers used for obtaining the gene fragment encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, a DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein can also be obtained by carrying out screening by PCR using the cDNA or cDNA library synthesized from the mRNA contained in a human or non-human animal tissue or cell as the template.

[0109] The nucleotide sequence of the obtained DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying

enzyme or the GDP-fucose transport protein is analyzed from its terminus and determined by a method generally used for analyzing a nucleotide such as the dideoxy method of Sanger *et al.* [*Proc. Natl. Acad. Sci. USA*, 74, 5463 (1977)] or by using a nucleotide sequence analyzer such as ABIPRISM 377 DNA Sequencer (manufactured by PE Biosystems).

[0110] A gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein can also be determined from genes in data bases by searching nucleotide sequence data bases such as GenBank, EMBL and DDBJ using a homology retrieving program such as BLAST based on the determined cDNA nucleotide sequence.

[0111] The nucleotide sequence of the gene encoding the GDP-fucose synthase obtained by the above method includes the nucleotide sequence represented by SEQ ID NO:48, 51 or 65. The nucleotide sequence of the gene encoding the α 1,6-fucose modifying enzyme includes the nucleotide sequence represented by SEQ ID NO: or 2. The nucleotide sequence of the gene encoding the GDP-fucose transport protein includes the nucleotide sequence represented by SEQ ID NO:91, 93, 95 or 97.

[0112] The cDNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein can also be obtained by chemically synthesizing it with a DNA synthesizer such as DNA Synthesizer model 392 manufactured by Perkin Elmer using the phosphoramidite method, based on the determined DNA nucleotide sequence.

[0113] The method for preparing a genomic DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes known methods described in *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*; and the like. Furthermore, the genomic DNA can be prepared by using a kit such as Genome DNA Library Screening System (manufactured by Genome Systems) or Universal GenomeWalker™ Kits (manufactured by CLONTECH).

[0114] The nucleotide sequence of the genomic DNA encoding the GDP-fucose synthase obtained by the method includes the nucleotide sequence represented by SEQ ID NO:67 or 70. The nucleotide sequence of the genomic DNA encoding the α 1,6-fucose modifying enzyme includes the nucleotide sequence represented by SEQ ID NO:3. The nucleotide sequence of the genomic DNA encoding the GDP-fucose transport protein includes the nucleotide sequence represented by SEQ ID NO:99 or 100.

[0115] In addition, the host cell can also be obtained without using an expression vector, by directly introducing an antisense oligonucleotide or ribozyme into a host cell, which is designed based on the nucleotide sequence encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein.

[0116] The antisense oligonucleotide or ribozyme can be prepared in the usual method or by using a DNA synthesizer. Specifically, it can be prepared based on the sequence information of an oligonucleotide having a corresponding sequence of continued 5 to 150 bases, preferably 5 to 60 bases, and more preferably 5 to 40 bases, among nucleotide sequences of a cDNA and a genomic DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein by synthesizing an oligonucleotide which corresponds to a sequence complementary to the oligonucleotide (antisense oligonucleotide) or a ribozyme comprising the oligonucleotide sequence.

[0117] The oligonucleotide includes oligo RNA and derivatives of the oligonucleotide (hereinafter referred to as "oligonucleotide derivatives").

[0118] The oligonucleotide derivatives includes oligonucleotide derivatives in which a phosphodiester bond in the oligonucleotide is converted into a phosphorothioate bond, an oligonucleotide derivative in which a phosphodiester bond in the oligonucleotide is converted into an N3'-P5' phosphoramidate bond, an oligonucleotide derivative in which ribose and a phosphodiester bond in the oligonucleotide are converted into a peptide-nucleic acid bond, an oligonucleotide derivative in which uracil in the oligonucleotide is substituted with C-5 propynyluracil, an oligonucleotide derivative in which uracil in the oligonucleotide is substituted with C-5 thiazoleuracil, an oligonucleotide derivative in which cytosine in the oligonucleotide is substituted with C-5 propynylcytosine, an oligonucleotide derivative in which cytosine in the oligonucleotide is substituted with phenoxazine-modified cytosine, an oligonucleotide derivative in which ribose in the oligonucleotide is substituted with 2'-O-propylribose and an oligonucleotide derivative in which ribose in the oligonucleotide is substituted with 2'-methoxyethoxyribose [*Cell Technology (Saibo Kogaku)*, 16, 1463 (1997)].

(b) Preparation of host cell used in the method of the present invention by homologous recombination

[0119] The host cell used in the method of the present invention can be prepared by targeting a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein and modifying the target gene on chromosome through a homologous recombination technique.

[0120] The target gene on the chromosome can be modified by using a method described in *Manipulating the Mouse Embryo, A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press (1994) (hereinafter referred to as "*Manipulating the Mouse Embryo, A Laboratory Manual*"); *Gene Targeting, A Practical Approach*, IRL Press at Oxford University Press (1993); *Biomanual Series 8, Gene Targeting, Preparation of Mutant Mice using ES*, Yodo-sha (1995) (hereinafter referred to as "*Preparation of Mutant Mice using ES Cells*"); or the like, for example, as follows:

[0121] A genomic DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is prepared.

[0122] Based on the nucleotide sequence of the genomic DNA, a target vector is prepared for homologous recombination of a target gene to be modified (e.g., structural gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein or a promoter gene).

[0123] The host cell used in the method of the present invention can be produced by introducing the prepared target vector into a host cell and selecting a cell in which homologous recombination occurred between the target gene and target vector.

[0124] As the host cell, any cell such as yeast, an animal cell, an insect cell or a plant cell can be used, so long as it has a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein. Examples include the host cells described in the following item 3.

[0125] The method for preparing a genomic DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods described in "Preparation method of genomic DNA" in the item 1(1)(a) and the like.

[0126] The nucleotide sequence of the genomic DNA encoding the GDP-fucose synthase obtained by the above method includes the nucleotide sequence represented by SEQ ID NO:67 or 70. The nucleotide sequence of the genomic DNA of the α 1,6-fucose modifying enzyme includes the nucleotide sequence represented by SEQ ID NO:3. The nucleotide sequence of the genomic DNA of the GDP-fucose transport protein includes the nucleotide sequence represented by SEQ ID NO:99 or 100.

[0127] The target vector for the homologous recombination of the target gene can be prepared in accordance with a method described in *Gene Targeting, A Practical Approach*, IRL Press at Oxford University Press (1993); *Preparation of Mutant Mice using ES Cells*; or the like. The target vector can be used as either a replacement type or an insertion type.

[0128] For introducing the target vector into various host cells, the methods for introducing recombinant vectors suitable for various host cells described in the following item 3 can be used.

[0129] The method for efficiently selecting a homologous recombinant includes a method such as the positive selection, promoter selection, negative selection or polyA selection described in *Gene Targeting, A Practical Approach*, IRL Press at Oxford University Press (1993); *Preparation of Mutant Mice using ES Cells*; or the like. The method for selecting the homologous recombinant of interest from the selected cell lines includes the Southern hybridization method for genomic DNA (*Molecular Cloning*, Second Edition), PCR [*PCR Protocols*, Academic Press (1990)], and the like.

(c) Preparation of cell of the present invention by RDO method

[0130] The host cell used in the method of the present invention can be prepared by targeting a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein according to an RDO method, for example, as follows.

[0131] A cDNA or a genomic DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is prepared.

[0132] The nucleotide sequence of the prepared cDNA or genomic DNA is determined.

[0133] Based on the determined DNA sequence, an RDO construct of an appropriate length comprising a DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, a DNA encoding an untranslated region or a DNA encoding an intron, is designed and synthesized.

[0134] The host cell used in the method of the present invention can be obtained by introducing the synthesized RDO into a host cell and then selecting a transformant in which a mutation occurred in the target enzyme, i.e., the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein.

[0135] As the host cell, any cell such as yeast, an animal cell, an insect cell or a plant cell can be used, so long as it has a gene encoding the target GDP-fucose synthase, α 1,6-fucose modifying enzyme or GDP-fucose transport protein. Examples include the host cells described in the following item 3.

[0136] The method for introducing RDO into various host cells includes the methods for introducing recombinant vectors suitable for various host cells described in the following item 3.

[0137] The method for preparing cDNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods described in "Preparation of DNA" in the item 1(1)(a) and the like.

[0138] The method for preparing a genomic DNA encoding the GDP-fucose synthase, α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods described in "Preparation method of genomic DNA" in the item 1(1)(a) and the like.

[0139] The nucleotide sequence of the DNA can be determined by digesting it with appropriate restriction enzymes, cloning the fragments into a plasmid such as pBluescript SK(-) (manufactured by Stratagene), subjecting the clones

to the reaction generally used as a method for analyzing a nucleotide sequence such as the dideoxy method of Sanger *et al.* [*Proc. Natl. Acad. Sci. USA*, 74, 5463 (1977)] or the like, and then analyzing the clones using an automatic nucleotide sequence analyzer such as ABI PSISM 377DNA Sequencer (manufactured by PE Biosystems) or the like.

[0140] The RDO can be prepared in the usual method or by using a DNA synthesizer.

[0141] The method for selecting a cell in which a mutation occurred, by introducing the RDO into the host cell, in the gene encoding the targeting enzyme, the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods for directly detecting mutations in chromosomal genes described in *Molecular Cloning*, Second Edition, *Current Protocols in Molecular Biology* and the like.

[0142] Furthermore, the method described in the item 1(1)(a) for selecting a transformant based on the activity of the introduced GDP-fucose synthase, α 1,6-fucose modifying enzyme or GDP-fucose transport protein and the method for selecting a transformant based on the sugar chain structure of a glycoprotein on the cell membrane described later in the item 1(5), and the method for selecting a transformant based on the sugar structure of a produced antibody molecule described later in the item 6 or 7 can also be used.

[0143] The construct of the RDO can be designed in accordance with the methods described in *Science*, 273, 1386 (1996); *Nature Medicine*, 4, 285 (1998); *Hepatology*, 25, 1462 (1997); *Gene Therapy*, 5, 1960 (1999); *J. Mol. Med.*, 75, 829 (1997); *Proc. Natl. Acad. Sci. USA*, 96, 8774 (1999); *Proc. Natl. Acad. Sci. USA*, 96, 8768 (1999); *Nuc. Acids. Res.*, 27, 1323 (1999); *Invest. Dermatol.*, 111, 1172 (1998); *Nature Biotech.*, 16, 1343 (1998); *Nature Biotech.*, 18, 43 (2000); *Nature Biotech.*, 18, 555 (2000); and the like.

(d) Preparation of host cell used in the method of the present invention by RNAi method

[0144] The host cell used in the method of the present invention can be prepared by targeting a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein according to the RNAi method, for example, as follows.

[0145] A cDNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is prepared.

[0146] The nucleotide sequence of the prepared cDNA is determined.

[0147] Based on the determined DNA sequence, an RNAi gene construct of an appropriate length comprising a DNA encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein or a DNA encoding a untranslated region, is designed.

[0148] In order to express the RNAi gene in a cell, a recombinant vector is prepared by inserting a fragment or full length of the prepared DNA into downstream of the promoter of an appropriate expression vector.

[0149] A transformant is obtained by introducing the recombinant vector into a host cell suitable for the expression vector.

[0150] The host cell used in the method of the present invention can be obtained by selecting a transformant based on the activity of the introduced GDP-fucose synthase, α 1,6-fucose modifying enzyme or GDP-fucose transport protein, or the sugar chain structure of the produced antibody molecule or of a glycoprotein on the cell membrane.

[0151] As the host cell, any cell such as yeast, an animal cell, an insect cell or a plant cell can be used, so long as it has a gene encoding the target GDP-fucose synthase, α 1,6-fucose modifying enzyme or GDP-fucose transport protein. Examples include the host cells described in the following item 3.

[0152] As the expression vector, a vector which is autonomously replicable in the above host cell or can be integrated into the chromosome and comprises a promoter at such a position that the designed RNAi gene can be transferred is used. Examples include the expression vectors described in the following item 3.

[0153] As the method for introducing a gene into various host cells, the methods for introducing recombinant vectors suitable for various host cells described in the following item 3 can be used.

[0154] The method for selecting a transformant based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods described in the item 1(1)(a).

[0155] The method for selecting a transformant based on the sugar chain structure of a glycoprotein on the cell membrane includes the methods described in the following item 1(5). The method for selecting a transformant based on the sugar chain structure of a produced antibody molecule includes the methods described in the following item 6 or 7.

[0156] The method for preparing cDNA of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods described in "Preparation of DNA" in the item 1(1)(a) and the like.

[0157] In addition, the host cell used in the method of the present invention can also be obtained without using an expression vector, by directly introducing an RNAi gene designed based on the nucleotide sequence of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein.

[0158] The RNAi gene can be prepared in the usual method or by using a DNA synthesizer.

[0159] The RNAi gene construct can be designed in accordance with the methods described in *Nature*, 391, 806

(1998); *Proc. Natl. Acad. Sci. USA*, 95, 15502 (1998); *Nature*, 395, 854 (1998); *Proc. Natl. Acad. Sci. USA*, 96, 5049 (1999); *Cell*, 95, 1017 (1998); *Proc. Natl. Acad. Sci. USA*, 96, 1451 (1999); *Proc. Natl. Acad. Sci. USA*, 95, 13959 (1998); *Nature Cell Biol.*, 2, 70 (2000); and the like.

(e) Preparation of host cell used in the method of the present invention by method using transposon

[0160] The host cell used in the method of the present invention can be prepared by selecting a mutant based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein or the sugar chain structure of a produced antibody molecule or of a glycoprotein on the cell membrane by using a transposon system described in *Nature Genet.*, 25, 35 (2000) or the like.

[0161] The transposon system is a system in which a mutation is induced by randomly inserting an exogenous gene into chromosome, wherein an exogenous gene interposed between transposons is generally used as a vector for inducing a mutation, and a transposase expression vector for randomly inserting the gene into chromosome is introduced into the cell at the same time.

[0162] Any transposase can be used, so long as it is suitable for the sequence of the transposon to be used.

[0163] As the exogenous gene, any gene can be used, so long as it can induce a mutation in the DNA of a host cell.

[0164] As the host cell, any cell such as yeast, an animal cell, an insect cell or a plant cell can be used, so long as it has a gene encoding the targeting GDP-fucose synthase, α 1,6-fucose modifying enzyme or GDP-fucose transport protein. Examples include the host cells described in the following item 3. For introducing the gene into various host cells, the method for introducing recombinant vectors suitable for various host cells described in the following item 3, can be used.

[0165] The method for selecting a mutant based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods which will be described above in the item 1 (1)(a).

[0166] The method for selecting a mutant based on the sugar chain structure of a glycoprotein on the cell membrane includes the methods described in the following item 1(5). The method for selecting a transformant based on the sugar chain structure of a produced antibody molecule includes the methods described in the following item 6 or 7.

(2) Method for introducing dominant negative mutant of enzyme

[0167] The host cell used in the method of the present invention can be prepared by targeting a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein according to a technique for introducing a dominant negative mutant of the enzyme. The GDP-fucose synthase includes GMD, Fx, GFPP, fucokinase and the like. The α 1,6-fucose modifying enzyme includes α 1,6-fucosyltransferase, α -L-fucosidase and the like. The GDP-fucose transport protein includes GDP-fucose transporter and the like.

[0168] The enzymes catalyze specific reactions having substrate specificity, and dominant negative mutants of the enzymes can be prepared by disrupting the active center of the enzymes which have the catalytic activity having substrate specificity. The method for preparing a dominant negative mutant is specifically described as follows with reference to GMD among the target enzymes.

[0169] As a result of the analysis of the three-dimensional structure of *E. coli*-derived GMD, it has been found that 4 amino acids (threonine at position 133, glutamic acid at position 135, tyrosine at position 157 and lysine at position 161) have an important function on the enzyme activity [*Structure*, 8, 2 (2000)]. That is, when mutants were prepared by substituting the 4 amino acids with other different amino acids based on the three-dimensional structure information, the enzyme activity of all of the mutants was significantly decreased. On the other hand, changes in the ability of GMD to bind to GMD coenzyme NADP and its substrate GDP-mannose were hardly observed in the mutants. Accordingly, a dominant negative mutant can be prepared by substituting the 4 amino acids which control the enzyme activity of GMD. A dominant negative mutant can be prepared by comparing the homology and predicting the three-dimensional structure using the amino acid sequence information based on the results of the *E. coli*-derived GMD. For example, in GMD (SEQ ID NO:65) derived from CHO cell, a dominant negative mutant can be prepared by substituting threonine at position 155, glutamic acid at position 157, tyrosine at position 179 and lysine at position 183. Such a gene into which amino acid substitution is introduced can be prepared by the site-directed mutagenesis described in *Molecular Cloning*, Second Edition, *Current Protocols in Molecular Biology* or the like.

[0170] The host cell can be prepared by using the above prepared dominant negative mutant gene of the target enzyme according to the method described in *Molecular Cloning*, Second Edition, *Current Protocols in Molecular Biology*, *Manipulating the Mouse Embryo*, Second Edition or the like, for example, as follows.

[0171] A gene encoding the dominant negative mutant of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein (hereinafter referred to as "dominant negative mutant gene") is prepared.

[0172] Based on the full length DNA of the prepared dominant negative mutant gene, a DNA fragment of an appro-

prate length containing a DNA encoding the antibody molecule is prepared, if necessary.

[0173] A recombinant vector is prepared by inserting the DNA fragment or full length DNA into downstream of the promoter of an appropriate expression vector.

[0174] A transformant is obtained by introducing the recombinant vector into a host cell suitable for the expression vector.

[0175] The host cell used in the method of the present invention can be prepared by selecting a transformant based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, or the sugar chain structure of a produced antibody molecule or of a glycoprotein on the cell membrane.

[0176] As the host cell, any cell such as yeast, an animal cell, an insect cell or a plant cell can be used, so long as it has a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein. Examples include the host cells described in the following item 3.

[0177] As the expression vector, a vector which is autonomously replicable in the host cell or can be integrated into the chromosome and comprises a promoter at a position where transcription of the DNA encoding the dominant negative mutant of interest can be effected is used. Examples include the expression vectors described in the following item 3.

[0178] For introducing the gene into various host cells, the method for introducing recombinant vectors suitable for various host cells described in the following item 3, can be used.

[0179] The method for selecting a mutant based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods which will be described in the above item 1 (1)(a).

[0180] The method for selecting a mutant based on the sugar chain structure of a glycoprotein on the cell membrane includes the methods described in the following item 1(5). The method for selecting a transformant based on the sugar chain structure of a produced antibody molecule includes the methods described in the following item 6 or 7.

(3) Method for introducing mutation into enzyme

[0181] The host cell used in the method of the present invention can be prepared by introducing a mutation into a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, and then selecting a cell line of interest in which the mutation occurred in the enzyme.

[0182] The GDP-fucose synthase includes GMD, Fx, GFPP, fucokinase and the like. The α 1,6-fucose modifying enzyme includes α 1,6-fucosyltransferase, α -L-fucosidase and the like. The GDP-fucose transport protein includes GDP-fucose transporter and the like.

[0183] The method for introducing mutation into an enzyme includes 1) a method in which a desired clone is selected from mutants obtained by a mutation-inducing treatment of a parent cell line with a mutagen or spontaneously generated mutants, based on the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, 2) a method in which a desired clone is selected from mutants obtained by a mutation-inducing treatment of a parent cell line with a mutagen or spontaneously generated mutants, based on the sugar chain structure of a produced antibody molecule, 3) a method in which a desired clone is selected from mutants obtained by a mutation-inducing treatment of a parent cell line with a mutagen or spontaneously generated mutants, based on the sugar chain structure of a glycoprotein on the cell membrane, and the like.

[0184] As the mutation-inducing treatment, any treatment can be used, so long as it can induce a point mutation or a deletion or frame shift mutation in the DNA of cells of the parent cell line.

[0185] Examples include treatment with ethyl nitrosourea, nitrosoguanidine, benzopyrene or an acridine pigment and treatment with radiation. Also, various alkylating agents and carcinogens can be used as mutagens. The method for allowing a mutagen to act upon cells includes the methods described in *Tissue Culture Techniques*, 3rd edition (Asakura Shoten), edited by Japanese Tissue Culture Association (1996), *Nature Genet.*, 24, 314 (2000) and the like.

[0186] The spontaneously generated mutant includes mutants which are spontaneously formed by continuing sub-culture under general cell culture conditions without applying special mutation-inducing treatment.

[0187] The method for measuring the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein includes the methods described above in the item 1(1)(a). The method for distinguishing the sugar chain structure of a produced antibody molecule includes the methods described in the following item 6 or 7. The method for distinguishing the sugar chain structure of a glycoprotein on the cell membrane includes the methods described in the following item 1(5).

(4) Method for suppressing transcription and/or translation of enzyme

[0188] The host cell used in the method of the present invention can be prepared by targeting a gene encoding the GDP-fucose synthase or the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein and suppressing

transcription and/or translation of the target gene according to the antisense RNA/DNA technique [*Bioscience and Industry*, 50, 322 (1992); *Chemistry*, 46, 681 (1991); *Biotechnology*, 9, 358 (1992); *Trends in Biotechnology*, 10, 87 (1992); *Trends in Biotechnology*, 10, 152 (1992); *Cell Engineering*, 16, 1463 (1997)], the triple helix technique [*Trends in Biotechnology*, 10, 132 (1992)] or the like.

[0189] The GDP-fucose synthase includes GMD, Fx, GFPP, fucokinase and the like. The α 1,6-fucose modifying enzyme includes α 1,6-fucosyltransferase, α -L-fucosidase and the like. The GDP-fucose transport protein includes GDP-fucose transporter and the like.

(5) Method for selecting clone resistant to lectin which recognizes sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain

[0190] The host cell used in the method of the present invention can be prepared by using a method for selecting a clone resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain.

[0191] The method for selecting a clone resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain includes the methods using lectin described in *Somatic Cell Mol. Genet.*, 12, 51 (1986) and the like.

[0192] As the lectin, any lectin can be used, so long as it is a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain. Examples include a *Lens culinaris* lectin LCA (lentil agglutinin derived from *Lens culinaris*), a pea lectin PSA (pea lectin derived from *Pisum sativum*), a broad bean lectin VFA (agglutinin derived from *Vicia faba*), an *Aleuria aurantia* lectin AAL (lectin derived from *Aleuria aurantia*) and the like.

[0193] Specifically, the clone of the present invention resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the *N*-glycoside-linked sugar chain can be selected by culturing cells by using a medium comprising the lectin at a concentration of 1 μ g/ml to 1 mg/ml for 1 day to 2 weeks, preferably 1 day to 1 week, subculturing surviving cells or picking up a colony and transferring it into a culture vessel, and subsequently continuing the culturing using the lectin-containing medium.

[0194] The method for confirming that the cell is a lectin-resistant cell includes a method for confirming expression of the GDP-fucose synthase, α 1,6-fucose modifying enzyme or the GDP-fucose transport protein, a method for culturing the cell in a medium to which lectin is directly added. Specifically, when the expression amount of the mRNA of α 1,6-fucosyltransferase which is one of α 1,6-fucose modifying enzymes in the cell is measured, a lectin-resistant cell decreases in an amount of the mRNA expressed.

2. Preparation of transgenic non-human animal or plant or the progenies

[0195] The cell used in the method of the present invention can be prepared by using a transgenic non-human animal or plant or the progenies thereof in which a genomic gene is modified in such a manner that the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is decreased or deleted. The transgenic non-human animal or plant or the progenies thereof can be prepared by targeting a gene encoding the above protein according to the method similar to that in the item 1.

[0196] In a transgenic non-human animal, the embryonic stem cell used in the process of the present invention in which the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is decreased or deleted can be prepared applying the method similar to that in the item 1 to an embryonic stem cell of the intended non-human animal such as cattle, sheep, goat, pig, horse, mouse, rat, fowl, monkey or rabbit.

[0197] Specifically, a mutant clone is prepared in which a gene encoding the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is inactivated or substituted with any nucleotide sequence, by a known homologous recombination technique [e.g., *Nature*, 326, 6110, 295 (1987); *Cell*, 51, 3, 503 (1987); etc.]. Using the prepared mutant clone, a chimeric individual comprising an embryonic stem cell clone and a normal cell can be prepared by an injection chimera method into blastocyst of fertilized egg of an animal or by an aggregation chimera method. The chimeric individual is crossed with a normal individual, so that a transgenic non-human animal in which the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is decreased or deleted in the whole body cells can be obtained.

[0198] The target vector for the homologous recombination of the target gene can be prepared in accordance with a method described in *Gene Targeting, A Practical Approach*, IRL Press at Oxford University Press (1993); *Preparation of Mutant Mice using ES Cells*, or the like. The target vector can be used as any of a replacement type, an insertion type and a gene trap type. As the method for introducing the target vector into the embryonic stem cell, any method can be used, so long as it can introduce DNA into an animal cell. Examples include electroporation [*Cytotechnology*,

3, 133 (1990)], the calcium phosphate method (Japanese Published Unexamined Patent Application No. 227075/90), the lipofection method [*Proc. Natl. Acad. Sci. USA*, 84, 7413 (1987)], the injection method [*Manipulating the Mouse Embryo*, Second Edition], a method using particle gun (gene gun) (Japanese Patent No. 2606856, Japanese Patent No. 2517813), the DEAE-dextran method [*Biomanual Series 4-Gene Transfer and Expression Analysis* (Yodo-sha), edited by Takashi Yokota and Kenichi Arai (1994)], the virus vector method [*Manipulating Mouse Embryo*, Second Edition] and the like.

[0199] The method for efficiently selecting a homologous recombinant includes a method such as the positive selection, promoter selection, negative selection or poly A selection described in *Gene Targeting, A Practical Approach*, IRL Press at Oxford University Press (1993), *Preparation of Mutant Mice using ES Cells*, or the like. Specifically, in the case of the target vector containing hprt gene, it is introduced into the hprt gene-defected embryonic stem cell, the embryonic stem cell is cultured in a medium containing aminopterin, hypoxanthine and thymidine, and positive selection which selects the homologous recombinant of the hprt gene can be carried out by selecting an a homogenous recombinant containing an aminopterin-resistant clone. In the case of the target vector containing a neomycin-resistant gene, the vector-introduced embryonic stem cell is cultured in a medium containing G418, and positive selection which selects a homogenous recombinant containing a neomycin-resistant gene can be carried out by selecting a G418-resistant clone. In the case of the target vector containing DT gene, the vector-introduced embryonic stem cell is cultured, and negative selection selecting a DT gene-free homogenous recombinant can be carried out by selecting the grown clone (in the recombinants introduced into a chromosome at random rather than the homogenous recombination, since the DT gene is expressed while integrated in the chromosome, the recombinants cannot grow because of the toxicity of DT). The method for selecting the homogenous recombinant of interest among the selected clones include the Southern hybridization for genomic DNA (*Molecular Cloning*, Second Edition), PCR [*PCR Protocols*, Academic Press (1990)] and the like.

[0200] When the embryonic stem cell is introduced into a fertilized egg by using an aggregation chimera method, in general, a fertilized egg at the development stage before 8-cell stage is preferably used. When the embryonic stem cell is introduced into a fertilized egg by using an injection chimera method, in general, it is preferred that a fertilized egg at the development stage from 8-cell stage to blastocyst stage is preferably used.

[0201] When the fertilized egg is transplanted into a female mouse, it is preferred that a fertilized egg obtained from a pseudopregnant female mouse in which fertility is induced by mating with a male non-human mammal which is subjected to vasoligation is artificially transplanted or implanted. Although the pseudopregnant female mouse can be obtained by natural mating, the pseudopregnant female mouse in which fertility is induced can be obtained by mating with a male mouse after administration of a luteinizing hormone-releasing hormone (hereinafter referred to as "LHRH") or its analogue thereof. The analogue of LHRH includes [3,5-DiI-Tyr5]-LHRH, [Gln8]-LHRH, [D-Ala6]-LHRH, des-Gly10-[D-His(Bzl)6]-LHRH ethylamide and the like.

[0202] Also, a fertilized egg cell of the present invention in which the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is decreased or deleted can be prepared by applying the method similar to that in the item 1 to fertilized egg of a non-human animal of interest such as cattle, sheep, goat, pig, horse, mouse, rat, fowl, monkey, rabbit or the like.

[0203] A transgenic non-human animal in which the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is decreased or deleted can be prepared by transplanting the prepared fertilized egg cell into the oviduct or uterus of a pseudopregnant female using the embryo transplantation method described in *Manipulating Mouse Embryo*, Second Edition or the like, followed by childbirth by the animal.

[0204] In a transgenic plant, the callus of the present invention in which the activity of the GDP-fucose synthase or the activity of an enzyme relating to modification of a sugar chain in which 1-position of fucose is bound to 6- or 3-position of N-acetylglucosamine in the reducing end in the N-glycoside-linked complex sugar chain is decreased or deleted can be prepared by applying the method similar to that in the item 1 to a callus or cell of the plant of interest.

[0205] A transgenic plant in which the activity of the GDP-fucose synthase, the α 1,6-fucose modifying enzyme or the GDP-fucose transport protein is decreased or deleted can be prepared by culturing the prepared callus in a medium comprising auxin and cytokinin to redifferentiate it in accordance with conventional methods [*Tissue Culture (Soshiki Baiyo)*, 20 (1994); *Tissue Cultur (Soshiki Baiyo)e*, 21 (1995); *Trends in Biotechnology*, 15, 45 (1997)].

3. Method for producing antibody composition

[0206] The antibody composition can be obtained by expressing it in a host cell by using the methods described in *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*; *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988 (hereinafter sometimes referred to as "*Antibodies*"); *Monoclonal Antibodies: Principles and Practice*, Third Edition, Acad. Press, 1996 (hereinafter sometimes referred to as "*Monoclonal Antibodies*"); and *Antibody Engineering, A Practical Approach*, IRL Press at Oxford University Press, 1996 (hereinafter sometimes referred to as "*Antibody Engineering*"), for example, as follows.

[0207] A full length cDNA of an antibody molecule is prepared, and a DNA fragment of an appropriate length comprising a DNA encoding the antibody molecule is prepared.

[0208] A recombinant vector is prepared by inserting the DNA fragment or the full length cDNA into downstream of the promoter of an appropriate expression vector.

5 [0209] A transformant which produces the antibody molecule can be obtained by introducing the recombinant vector into a host cell suitable for the expression vector.

[0210] As the host cell, the host cell of any cell such as yeast, an animal cell, an insect cell, a plant cell or the like can be used, so long as it can express the gene of interest.

10 [0211] A cell such as yeast, an animal cell, an insect cell, a plant cell or the like into which an enzyme relating to the modification of an *N*-glycoside-linked sugar chain which binds to the Fc region of the antibody molecule is introduced by a genetic engineering technique can also be used as the host cell.

[0212] As the expression vector, a vector which is autonomously replicable in the host cell or can be integrated into the chromosome and comprises a promoter at such a position that the DNA encoding the antibody molecule of interest can be transferred is used.

15 [0213] The cDNA can be prepared from a human or non-human tissue or cell by using a probe primer specific for the antibody molecule of interest and the like according to the methods described in "Preparation of DNA" in the item 1(1)(a).

[0214] When yeast is used as the host cell, the expression vector includes YEP 13 (ATCC 37115), YE24 (ATCC 37051), YCp50 (ATCC 37419) and the like.

20 [0215] As the promoter, any promoter can be used so long as it can function in yeast. Examples include a promoter of a gene relating to the glycolytic pathway such as a hexose kinase, PHO5 promoter, PGK promoter, GAP promoter, ADH promoter, gal 1 promoter, gal 10 promoter, heat shock protein promoter, MF α 1 promoter, CUP 1 promoter and the like.

25 [0216] The host cell includes yeast belonging to the genus *Saccharomyces*, the genus *Schizosaccharomyces*, the genus *Kluyveromyces*, the genus *Trichosporon*, the genus *Schwanniomyces* and the like, such as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Trichosporon pullulans* and *Schwanniomyces alluvius*.

30 [0217] As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into yeast. Examples include electroporation [*Methods in Enzymology*, 194, 182 (1990)], spheroplast method [*Proc. Natl. Acad. Sci. USA*, 84, 1929 (1978)], lithium acetate method [*J. Bacteriol.*, 153, 163 (1983)], a method described in *Proc. Natl. Acad. Sci. USA*, 75, 1929 (1978) and the like.

35 [0218] When an animal cell is used as the host cell, the expression vector includes pcDNA1, pcDM8 (available from Funakoshi), pAGE107 [Japanese Published Unexamined Patent Application No. 22979/91; *Cytotechnology*, 3, 133 (1990)], pAS3-3 [Japanese Published Unexamined Patent Application No. 227075/90], pCDM8 [*Nature*, 329, 840 (1987)], pcDNA1/Amp (manufactured by Invitrogen), pREP4 (manufactured by Invitrogen), pAGE103 [*J. Biochemistry*, 101, 1307 (1987)], pAGE210 and the like.

40 [0219] As the promoter, any promoter can be used, so long as it can function in an animal cell. Examples include a promoter of IE (immediate early) gene of cytomegalovirus (CMV), an early promoter of SV40, a promoter of retrovirus, a promoter of metallothionein, a heat shock promoter, an SR α promoter and the like. Also, an enhancer of the IE gene of human CMV may be used together with the promoter.

[0220] The host cell includes a human cell such as Namalwa cell, a monkey cell such as COS cell, a Chinese hamster cell such as CHO cell or HBT5637 [Japanese Published Unexamined Patent Application No. 299/88], a rat myeloma cell, a mouse myeloma cell, a cell derived from syrian hamster kidney, an embryonic stem cell, a fertilized egg cell and the like.

45 [0221] As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into an animal cell. Examples include electroporation [*Cytotechnology*, 3, 133 (1990)], the calcium phosphate method [Japanese Published Unexamined Patent Application No. 227075/90], the lipofection method [*Proc. Natl. Acad. Sci. USA*, 84, 7413 (1987)], the injection method [*Manipulating the Mouse Embryo, A Laboratory Manual*], a method using particle gun (gene gun) [Japanese Patent No. 2606856, Japanese Patent No. 2517813], the DEAE-dextran method [*Biomaterial Series 4-Gene Transfer and Expression Analysis* (Yodo-sha), edited by Takashi Yokota and Kenichi Arai (1994)], the virus vector method [*Manipulating Mouse Embryo, Second Edition*] and the like.

50 [0222] When an insect cell is used as the host cell, the protein can be expressed by the method described in *Current Protocols in Molecular Biology, Baculovirus Expression Vectors, A Laboratory Manual*, W.H. Freeman and Company, New York (1992), *Bio/Technology*, 6, 47 (1988) or the like.

55 [0223] That is, the protein can be expressed by co-introducing a recombinant gene-introducing vector and a baculovirus into an insect cell to obtain a recombinant virus in an insect cell culture supernatant and then infecting the insect cell with the recombinant virus.

[0224] The gene introducing vector used in the method includes pVL1392, pVL1393, pBlueBacIII (all manufactured

by Invitrogen) and the like.

[0225] The baculovirus includes *Autographa californica* nuclear polyhedrosis virus which is infected by an insect of the family *Barathra*.

[0226] The insect cell includes *Spodoptera frugiperda* oocytes Sf9 and Sf21 [*Current Protocols in Molecular Biology, Baculovirus Expression Vectors, A Laboratory Manual*, W.H. Freeman and Company, New York (1992)], a *Trichoplusia ni* oocyte High 5 (manufactured by Invitrogen) and the like.

[0227] The method for co-introducing the above recombinant gene-introducing vector and the baculovirus to the insect cells for preparing the above recombinant virus includes the calcium phosphate method (Japanese Published Unexamined Patent Application No. 227075/90), the lipofection method [*Proc. Natl. Acad. Sci. USA*, 84, 7413 (1987)] and the like.

[0228] When a plant cell is used as the host cell, the expression vector includes Ti plasmid, tobacco mosaic virus and the like.

[0229] As the promoter, any promoter can be used, so long as it can function in a plant cell. Examples include cauliflower mosaic virus (CaMV) 35S promoter, rice actin 1 promoter and the like.

[0230] The host cell includes plant cells of tobacco, potato, tomato, carrot, soybean, rape, alfalfa, rice, wheat, barley and the like.

[0231] As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into a plant cell. Examples include a method using *Agrobacterium* (Japanese Published Unexamined Patent Application No. 140885/84, Japanese Published Unexamined Patent Application No. 70080/85, WO94/00977), electroporation (Japanese Published Unexamined Patent Application No. 251887/85), a method using a particle gun (gene gun) (Japanese Patent No. 2606856, Japanese Patent No. 2517813) and the like.

[0232] As the method for expressing an antibody gene, secretion production, expression of a fusion protein of the Fc region with other protein and the like can be carried out in accordance with the method described in *Molecular Cloning*, Second Edition or the like, in addition to the direct expression.

[0233] When a gene is expressed by a microorganism, yeast, an animal cell, an insect cell or a plant cell into which a gene relating to the synthesis of a sugar chain is introduced, an antibody molecule to which a sugar or a sugar chain is added by the introduced gene can be obtained.

[0234] An antibody composition can be produced by culturing the obtained transformant in a medium to produce and accumulate the antibody molecule in the culture and then recovering it from the resulting culture. The method for culturing the transformant in a medium can be carried out in accordance with a general method which is used for the culturing of host cells.

[0235] As the medium for culturing a transformant obtained using yeast as the host cell, the medium may be either a natural medium or a synthetic medium, so long as it comprises materials such as a carbon source, a nitrogen source and an inorganic salt which can be assimilated by the organism and culturing of the transformant can be efficiently carried out.

[0236] As the carbon source, those which can be assimilated by the organism can be used. Examples include carbohydrates such as glucose, fructose, sucrose, molasses thereof, starch and starch hydrolysate; organic acids such as acetic acid and propionic acid; alcohols such as ethanol and propanol; and the like.

[0237] The nitrogen source includes ammonia; ammonium salts of inorganic acid or organic acid such as ammonium chloride, ammonium sulfate, ammonium acetate and ammonium phosphate; other nitrogen-containing compounds; peptone; meat extract; yeast extract; corn steep liquor; casein hydrolysate; soybean meal; soybean meal hydrolysate; various fermented cells and hydrolysates thereof; and the like.

[0238] The inorganic salt includes potassium dihydrogen phosphate, dipotassium hydrogen phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate, calcium carbonate, and the like.

[0239] The culturing is carried out generally under aerobic conditions such as a shaking culture or submerged-aeration stirring culture. The culturing temperature is preferably at 15 to 40°C, and the culturing time is generally 16 hours to 7 days. During the culturing, the pH is maintained at 3.0 to 9.0. The pH is adjusted using an inorganic or organic acid, an alkali solution, urea, calcium carbonate, ammonia or the like.

[0240] Furthermore, if necessary, an antibiotic such as ampicillin or tetracycline can be added to the medium during the culturing.

[0241] When yeast transformed with a recombinant vector obtained by using an inducible promoter as the promoter is cultured, an inducer can be added to the medium, if necessary. For example, when yeast transformed with a recombinant vector obtained by using *lac* promoter is cultured, isopropyl- β -D-thiogalactopyranoside and the like can be added to the medium, and when yeast transformed with a recombinant vector obtained by using *trp* promoter is cultured, indoleacrylic acid and the like can be added to the medium.

[0242] When a transformant obtained by using an animal cell as the host is cultured, the medium includes generally used RPMI 1640 medium [*The Journal of the American Medical Association*, 199, 519 (1967)], Eagle's MEM medium

[*Science*, 122, 501 (1952)], Dulbecco's modified MEM medium [*Virology*, 8, 396 (1959)], 199 medium [*Proceeding of the Society for the Biological Medicine*, 73, 1 (1950)] and Whitten's medium [*Developmental Engineering Experimentation Manual-Preparation of Transgenic Mice* (Kodan-sha), edited by M. Katsuki (1987)], the media to which fetal calf serum, etc. are added, and the like.

[0243] The culturing is carried out generally at conditions under pH of 6 to 8 and 30 to 40°C for 1 to 7 days in the presence of 5% CO₂ and the like, for 1 to 7 days.

[0244] Furthermore, if necessary, an antibiotic such as kanamycin or penicillin can be added to the medium during the culturing.

[0245] The medium for culturing a transformant obtained by using an insect cell as the host includes generally used TNM-FH medium (manufactured by Pharmingen), Sf-900 II SFM medium (manufactured by Life Technologies), ExCell 400 and ExCell 405 (both manufactured by JRH Biosciences), Grace's Insect Medium [*Nature*, 195, 788 (1962)] and the like.

[0246] The culturing is carried out generally at conditions under pH of 6 to 7 and 25 to 30°C and the like, for 1 to 5 days.

[0247] Furthermore, if necessary, an antibiotic such as gentamicin can be added to the medium during the culturing.

[0248] A transformant obtained by using a plant cell as the host can be cultured as a cell or by differentiating it into a plant cell or organ. The medium for culturing the transformant includes generally used Murashige and Skoog (MS) medium and White medium, the media to which a plant hormone such as auxin or cytokinin is added, and the like.

[0249] The culturing is carried out generally at conditions under pH of 5 to 9 and 20 to 40°C for 3 to 60 days.

[0250] Furthermore, if necessary, an antibiotic such as kanamycin or hygromycin can be added to the medium during the culturing.

[0251] As described above, an antibody composition can be produced by culturing a transformant derived from yeast, an animal cell, an insect cell or a plant cell, which comprises a recombinant vector into which a DNA encoding an antibody molecule is inserted, in accordance with a general culturing method, to thereby produce and accumulate the antibody composition, and then recovering the antibody composition from the culture.

[0252] As the method for expressing the gene encoding an antibody, secretion production, expression of a fusion protein and the like can be carried out in accordance with the method described in *Molecular Cloning*, Second Edition in addition to the direct expression.

[0253] The method for producing an antibody composition includes a method of intracellular expression in a host cell, a method of extracellular secretion from a host cell, and a method of production on a host cell membrane outer envelope. The method can be selected by changing the host cell used or the structure of the antibody composition produced.

[0254] The method for producing an antibody composition includes a method of intracellular expression in a host cell, a method of extracellular secretion from a host cell, and a method of production on a host cell membrane outer envelope. The method can be selected by changing the host cell used or the structure of the antibody composition produced.

[0255] When the antibody composition is produced in a host cell or on a host cell membrane outer envelope, it can be positively secreted extracellularly in accordance with the method of Paulson *et al.* [*J. Biol. Chem.*, 264, 17619 (1989)], the method of Lowe *et al.* [*Proc. Natl. Acad. Sci. USA*, 86, 8227 (1989), *Genes Develop.*, 4, 1288 (1990)], the methods described in Japanese Published Unexamined Patent Application No. 336963/93 and Japanese Published Unexamined Patent Application No. 823021/94 and the like.

[0256] That is, an antibody molecule of interest can be positively secreted extracellularly from a host cell by inserting a DNA encoding the antibody molecule and a DNA encoding a signal peptide suitable for the expression of the antibody molecule into an expression vector according to a gene recombination technique and then expressing the antibody molecule.

[0257] Also, its production amount can be increased in accordance with the method described in Japanese Published Unexamined Patent Application No. 227075/90 according to a gene amplification system using a dihydrofolate reductase gene.

[0258] In addition, the antibody composition can also be produced by using a gene-introduced animal individual (transgenic non-human animal) or a plant individual (transgenic plant) which is constructed by the redifferentiation of an animal or plant cell into which the gene is introduced.

[0259] When the transformant is an animal individual or a plant individual, an antibody composition can be produced in accordance with a general method by rearing or cultivating it to thereby produce and accumulate the antibody composition and then recovering the antibody composition from the animal or plant individual.

[0260] The method for producing an antibody composition using an animal individual includes a method in which the antibody composition of interest is produced in an animal constructed by introducing a gene in accordance with a known method [*American Journal of Clinical Nutrition*, 63, 639S (1996); *American Journal of Clinical Nutrition*, 63, 627S (1996); *BioTechnology*, 9, 830 (1991)].

[0261] In the case of an animal individual, an antibody composition can be produced, for example, by rearing a

transgenic non-human animal into which a DNA encoding an antibody molecule is introduced to thereby produce and accumulate the antibody composition in the animal, and then recovering the antibody composition from the animal. The place of the animal where the composition is produced and accumulated includes milk (Japanese Published Un-
 5 examined Patent Application No. 309192/88) and eggs of the animal. As the promoter used in this case, any promoter can be used, so long as it can function in an animal. Preferred examples include mammary gland cell-specific promoters such as α casein promoter, β casein promoter, β lactoglobulin promoter, whey acidic protein promoter and the like.

[0262] The method for producing an antibody composition using a plant individual includes a method in which an antibody composition is produced by cultivating a transgenic plant into which a DNA encoding an antibody molecule is introduced by a known method [*Tissue Culture (Soshiki Baiyo)*, 20 (1994); *Tissue Culture (Soshiki Baiyo)*, 21 (1995);
 10 *Trends in Biotechnology*, 15, 45 (1997)] to produce and accumulate the antibody composition in the plant, and then recovering the antibody composition from the plant.

[0263] Regarding purification of an antibody composition produced by a transformant into which a gene encoding an antibody molecule is introduced, for example, when the antibody composition is intracellularly expressed in a dissolved state, the cells after culturing are recovered by centrifugation, suspended in an aqueous buffer and then disrupted by using ultrasonic oscillator, French press, Manton Gaulin homogenizer, dynamill or the like to obtain a cell-
 15 free extract. A purified product of the antibody composition can be obtained from a supernatant obtained by centrifuging the cell-free extract according to a general enzyme isolation purification techniques such as solvent extraction; salting out or desalting with ammonium sulfate; precipitation with an organic solvent; anion exchange chromatography using a resin such as diethylaminoethyl (DEAE)-Sepharose or DIAION HPA-75 (manufactured by Mitsubishi Chemical); cation exchange chromatography using a resin such as S-Sepharose FF (manufactured by Pharmacia);, hydrophobic chromatography using a resin such as butyl-Sepharose or phenyl-Sepharose, gel filtration using a molecular sieve; affinity chromatography; chromatofocusing; electrophoresis such as isoelectric focusing; and the like which may be used alone or in combination.

[0264] Also, when the antibody composition is expressed intracellularly by forming an insoluble body, the cells are recovered, disrupted and centrifuged in the same manner, and the insoluble body of the antibody composition is recovered as a precipitation fraction. The recovered insoluble body of the antibody composition is solubilized by using a protein denaturing agent. The antibody composition is made into a normal three-dimensional structure by diluting or dialyzing the solubilized solution, and then a purified product of the antibody composition is obtained by the same isolation purification method.

[0265] When the antibody composition is secreted extracellularly, the antibody composition or derivatives thereof can be recovered from the culture supernatant. That is, the culture is treated according to a technique such as centrifugation as described above to obtain a soluble fraction, and a purified preparation of the antibody composition can be obtained from the soluble fraction by the same isolation purification method as described above.

[0266] The thus obtained antibody composition includes an antibody, the fragment of the antibody, a fusion protein comprising the Fc region of the antibody, and the like.

[0267] As an example for obtaining the antibody composition, a method for producing a composition of a humanized antibody and Fc fusion protein is described below in detail, but other antibody compositions can also be obtained in a manner similar to the method.

40 A. Preparation of humanized antibody composition

(1) Construction of vector for expression of humanized antibody

[0268] A vector for expression of humanized antibody is an expression vector for animal cell into which genes encoding CH and CL of a human antibody are inserted, which can be constructed by cloning each of genes encoding CH and CL of a human antibody into an expression vector for animal cell.

[0269] The C regions of a human antibody may be CH and CL of any human antibody. Examples include the C region belonging to IgG1 subclass in the H chain of a human antibody (hereinafter referred to as "hC γ 1"), the C region belonging to κ class in the L chain of a human antibody (hereinafter referred to as "hC κ "), and the like.

[0270] As the genes encoding CH and CL of a human antibody, a chromosomal DNA comprising an exon and an intron can be used, and a cDNA can also be used.

[0271] As the expression vector for animal cell, any vector can be used, so long as a gene encoding the C region of a human antibody can be inserted therein and expressed therein. Examples include pAGE107 [*Cytotechnology*, 3, 133 (1990)], pAGE103 [*J. Biochem.*, 101, 1307 (1987)], pHSG274 [*Gene*, 27, 223 (1984)], pKCR [*Proc. Natl. Acad. Sci. USA*, 78, 1527 (1981)], pSG1 β d2-4 [*Cytotechnology*, 4, 173 (1990)] and the like. The promoter and enhancer in the expression vector for animal cell includes SV40 early promoter and enhancer [*J. Biochem.*, 101, 1307 (1987)], Moloney mouse leukemia virus LTR promoter [*Biochem. Biophys. Res. Commun.*, 149, 960 (1987)], immunoglobulin H chain promoter [*Cell*, 41, 479 (1985)] and enhancer [*Cell*, 33, 717 (1983)], and the like.

[0272] The vector for expression of humanized antibody may be either of a type in which genes encoding the H chain and L chain of an antibody exist on separate vectors or of a type in which both genes exist on the same vector (hereinafter referred to "tandem type"). In respect of easiness of construction of a vector for expression of humanized antibody, easiness of introduction into animal cells, and balance between the expression amounts of the H and L chains consisting of an antibody in animal cells, a tandem type of the vector for humanized antibody expression is more preferred [J. Immunol. Methods, 167, 271 (1994)].

[0273] The constructed vector for expression of humanized antibody can be used for expression of a human chimeric antibody and a human CDR-grafted antibody in animal cells.

(2) Preparation method of cDNA encoding V region of non-human animal antibody

[0274] cDNAs encoding VH and VL of a non-human animal antibody such as mouse antibody can be obtained in the following manner.

[0275] A cDNA is synthesized from mRNA extracted from a hybridoma cell which produces the mouse antibody of interest. The synthesized cDNA is cloned into a vector such as a phage or a plasmid to obtain a cDNA library. Each of a recombinant phage or recombinant plasmid comprising a cDNA encoding VH and a recombinant phage or recombinant plasmid comprising a cDNA encoding VL is isolated from the library by using a C region part or a V region part of an existing mouse antibody as the probe. Full nucleotide sequences of VH and VL of the mouse antibody of interest on the recombinant phage or recombinant plasmid are determined, and full length amino acid sequences of VH and VL are deduced from the nucleotide sequences.

[0276] As the non-human animal, any animal such as mouse, rat, hamster or rabbit can be used, so long as a hybridoma cell can be produced therefrom.

[0277] The method for preparing a total RNA from a hybridoma cell includes the guanidine thiocyanate-caesium trifluoroacetate method [Methods in Enzymology, 154, 3 (1987)] and the like, and the method for preparing mRNA from total RNA includes an oligo(dT)-immobilized cellulose column method (Molecular Cloning, Second Edition) and the like. In addition, a kit for preparing mRNA from a hybridoma cell includes Fast Track mRNA Isolation Kit (manufactured by Invitrogen), Quick Prep mRNA Purification Kit (manufactured by Pharmacia) and the like.

[0278] The method for synthesizing a cDNA and preparing a cDNA library includes the usual methods (Molecular Cloning, Second Edition, Current Protocols in Molecular Biology, Supplement 1-34), methods using a commercially available kit such as SuperScript™, Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by GIBCO BRL) or ZAP-cDNA Synthesis Kit (manufactured by Stratagene), and the like.

[0279] In preparing the cDNA library, the vector into which a cDNA synthesized by using mRNA extracted from a hybridoma cell as the template is inserted may be any vector, so long as the cDNA can be inserted. Examples include ZAP Express [Strategies, 5, 58 (1992)], pBluescript II SK(+) [Nucleic Acids Research, 17, 9494 (1989)], λzapII (manufactured by Stratagene), λgt10 and λgt11 [DNA Cloning, A Practical Approach, 1, 49 (1985)], Lambda BlueMid (manufactured by Clontech), λExCell, pT7T3 18U (manufactured by Pharmacia), pcD2 [Mol. Cell. Biol., 3, 280 (1983)], pUC18 [Gene, 33, 103 (1985)] and the like.

[0280] As Escherichia coli into which the cDNA library constructed from a phage or plasmid vector is introduced, any Escherichia coli can be used, so long as the cDNA library can be introduced, expressed and maintained. Examples include XL1-Blue MRF' [Strategies, 5, 81 (1992)], C600 [Genetics, 39, 440 (1954)], Y1088 and Y1090 [Science, 222, 778 (1983)], NM522 [J. Mol. Biol., 166, 1 (1983)], K802 [J. Mol. Biol., 16, 118 (1966)], JM105 [Gene, 38, 275 (1985)] and the like.

[0281] As the method for selecting a cDNA clone encoding VH and VL of a non-human animal antibody from the cDNA library, a colony hybridization or a plaque hybridization using an isotope- or fluorescence-labeled probe can be used (Molecular Cloning, Second Edition). The cDNA encoding VH and VL can also be prepared by preparing primers and carrying out polymerase chain reaction (hereinafter referred to as "PCR"; Molecular Cloning, Second Edition; Current Protocols in Molecular Biology, Supplement 1-34) using a cDNA synthesized from mRNA or a cDNA library as the template.

[0282] The nucleotide sequences of the cDNAs can be determined by digesting the selected cDNAs with appropriate restriction enzymes, cloning the fragments into a plasmid such as pBluescript SK(-) (manufactured by Stratagene), carrying out the reaction of a generally used nucleotide sequence analyzing method such as the dideoxy method of Sanger et al. [Proc. Natl. Acad. Sci., USA, 74, 5463 (1977)], and then analyzing the clones using an automatic nucleotide sequence analyzer such as A.L.F. DNA Sequencer (manufactured by Pharmacia).

[0283] Whether or not the obtained cDNAs encode the full length amino acid sequences of VH and VL of the antibody comprising a secretory signal sequence can be confirmed by deducing the full length amino acid sequences of VH and VL from the determined nucleotide sequence and comparing them with the full length amino acid sequences of VH and VL of known antibodies [Sequences of Proteins of Immunological Interest, US Dep. Health and Human Services (1991), hereinafter referred to as "Sequences of Proteins of Immunological Interest"].

(3) Analysis of amino acid sequence of V region of non-human animal antibody

[0284] Regarding the full length amino acid sequences of VH and VL of the antibody comprising a secretory signal sequence, the length of the secretory signal sequence and the N-terminal amino acid sequences can be deduced and subgroups to which they belong can also be found, by comparing them with the full length amino acid sequences of VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*). In addition, the amino acid sequences of each CDR of VH and VL can also be found by comparing them with the amino acid sequences of VH and VL of known antibodies (*Sequences of Proteins of Immunological Interest*).

(4) Construction of human chimeric antibody expression vector

[0285] A human chimeric antibody expression vector can be constructed by cloning cDNAs encoding VH and VL of a non-human animal antibody into upstream of genes encoding CH and CL of a human antibody in the vector for expression of humanized antibody constructed described in the item 3(1). For example, a human chimeric antibody expression vector can be constructed by linking each of cDNAs encoding VH and VL of a non-human animal antibody to a synthetic DNA comprising nucleotide sequences at the 3'-terminals of VH and VL of a non-human animal antibody and nucleotide sequences at the 5'-terminals of CH and CL of a human antibody and also having a recognition sequence of an appropriate restriction enzyme at both terminals, and by cloning them into upstream of genes encoding CH and CL of a human antibody contained in the vector for expression of humanized antibody constructed described in the item 3(1) in such a manner that they can be expressed in a suitable form.

(5) Construction of cDNA encoding V region of human CDR-grafted antibody

[0286] cDNAs encoding VH and VL of a human CDR-grafted antibody can be obtained as follows. First, amino acid sequences of the frameworks (hereinafter referred to as "FR") of VH and VL of a human antibody for grafting CDR of VH and VL of a non-human animal antibody is selected. As the amino acid sequences of FRs of VH and VL of a human antibody, any amino acid sequences can be used so long as they are derived from a human antibody. Examples include amino acid sequences of FRs of VH and VL of human antibodies registered at databases such as Protein Data Bank, amino acid sequences common in each subgroup of FRs of VH and VL of human antibodies (*Sequences of Proteins of Immunological Interest*) and the like. In order to produce a human CDR-grafted antibody having enough activities, it is preferred to select an amino acid sequence having homology as high as possible (at least 60% or more) with amino acid sequences of VH and VL of a non-human animal antibody of interest.

[0287] Next, the amino acid sequences of CDRs of VH and VL of the non-human animal antibody of interest are grafted to the selected amino acid sequences of FRs of VH and VL of a human antibody to design amino acid sequences of VH and VL of the human CDR-grafted antibody. The designed amino acid sequences are converted into DNA sequences by considering the frequency of codon usage found in nucleotide sequences of antibody genes (*Sequences of Proteins of Immunological Interest*), and the DNA sequences encoding the amino acid sequences of VH and VL of the human CDR-grafted antibody are designed. Based on the designed DNA sequences, several synthetic DNAs having a length of about 100 bases are synthesized, and PCR is carried out by using them. In this case, it is preferred in each of the H chain and the L chain that 6 synthetic DNAs are designed in view of the reaction efficiency of PCR and the lengths of DNAs which can be synthesized.

[0288] Also, they can be easily cloned into the vector for expression of humanized antibody described in the item 3 (1) by introducing recognition sequences of an appropriate restriction enzyme into the 5'-terminals of the synthetic DNA on both terminals. After the PCR, the amplified product is cloned into a plasmid such as pBluescript SK(-) (manufactured by Stratagene) and the nucleotide sequences are determined by the method in the item 3(2) to thereby obtain a plasmid having DNA sequences encoding the amino acid sequences of VH and VL of the desired human CDR-grafted antibody.

(6) Construction of human CDR-grafted antibody expression vector

[0289] A human CDR-grafted antibody expression vector can be constructed by cloning the cDNAs encoding VH and VL of the human CDR-grafted antibody constructed in the item 3(5) into upstream of the gene encoding CH and CL of a human antibody in the vector for expression of humanized antibody described in the item 3(1). For example, recognizing sequences of an appropriate restriction enzyme are introduced into the 5'-terminals of both terminals of a synthetic DNA fragment, among the synthetic DNA fragments which are used in the item 3(5) for constructing the VH and VL of the human CDR-grafted antibody, so that they are cloned into upstream of the genes encoding CH and CL of a human antibody in the vector for expression of humanized antibody described in the item 3(1) in such a manner that they can be expressed in a suitable form, to thereby construct the human CDR-grafted antibody expression vector.

(7) Stable production of humanized antibody

[0290] A transformant capable of stably producing a human chimeric antibody and a human CDR-grafted antibody (both hereinafter referred to as "humanized antibody") can be obtained by introducing the vectors for humanized antibody expression described in the items 3(4) and (6) into an appropriate animal cell.

[0291] The method for introducing a humanized antibody expression vector into an animal cell includes electroporation [Japanese Published Unexamined Patent Application No. 257891/90, *Cytotechnology*, 3, 133 (1990)] and the like.

[0292] As the animal cell into which a humanized antibody expression vector is introduced, any cell can be used so long as it is an animal cell which can produce the humanized antibody.

[0293] Examples include mouse myeloma cells such as NSO cell and SP2/0 cell, Chinese hamster ovary cells such as CHO/*dhfr* cell and CHO/DG44 cell, rat myeloma such as YB2/0 cell and IR983F cell, BHK cell derived from a syrian hamster kidney, a human myeloma cell such as Namalwa cell, and the like, and a Chinese hamster ovary cell CHO/DG44 cell, a rat myeloma YB2/0 cell and the host cells of the present invention described in the item 5 are preferred.

[0294] After introduction of the humanized antibody expression vector, a transformant capable of stably producing the humanized antibody can be selected by using a medium for animal cell culture comprising an agent such as G418 sulfate (hereinafter referred to as "G418"; manufactured by SIGMA) and the like in accordance with the method described in Japanese Published Unexamined Patent Application No. 257891/90. The medium to culture animal cells includes RPMI 1640 medium (manufactured by Nissui Pharmaceutical), GIT medium (manufactured by Nihon Pharmaceutical), EX-CELL 302 medium (manufactured by JRH), IMDM medium (manufactured by GIBCO BRL), Hybridoma-SFM medium (manufactured by GIBCO BRL), media obtained by adding various additives such as fetal bovine serum (hereinafter referred to as "FBS") to these media, and the like. The humanized antibody can be produced and accumulated in the culture supernatant by culturing the obtained transformant in a medium. The amount of production and antigen binding activity of the humanized antibody in the culture supernatant can be measured by a method such as enzyme-linked immunosorbent assay (hereinafter referred to as "ELISA"; *Antibodies, Monoclonal Antibodies*) or the like. Also, the amount of the humanized antibody produced by the transformant can be increased by using a DHFR gene amplification system in accordance with the method described in Japanese Published Unexamined Patent Application No. 257891/90.

[0295] The humanized antibody can be purified from a culture supernatant culturing the transformant by using a protein A column (*Antibodies, Chapter 8; Monoclonal Antibodies*). In addition, purification methods generally used for the purification of proteins can also be used. For example, the purification can be carried out through the combination of gel filtration, ion exchange chromatography and ultrafiltration. The molecular weight of the H chain, L chain or antibody molecule as a whole of the purified humanized antibody can be measured, e.g., by polyacrylamide gel electrophoresis [hereinafter referred to as "SDS-PAGE"; *Nature*, 227, 680 (1970)], Western blotting (*Antibodies, Monoclonal Antibodies*) or the like.

B. Preparation of Fc fusion protein

(1) Construction of Fc fusion protein expression vector

[0296] An Fc fusion protein expression vector is an expression vector for animal cells into which genes encoding the Fc region of a human antibody and a protein to be fused are inserted, which can be constructed by cloning each of genes encoding the Fc region of a human antibody and the protein to be fused into an expression vector for animal cell.

[0297] The Fc region of a human antibody includes those containing a part of a hinge region and/or CH1 in addition to regions containing CH2 and CH3 regions. Also, it can be any Fc region, so long as at least one amino acid of CH2 or CH3 may be deleted, substituted, added or inserted, and substantially has the binding activity to the Fc γ receptor.

[0298] As the genes encoding the Fc region of a human antibody and the protein to be fused, a chromosomal DNA comprising an exon and an intron can be used, and a cDNA can also be used. The method for linking the genes and the Fc region includes PCR using each of the gene sequences as the template (*Molecular Cloning, Second Edition; Current Protocols in Molecular Biology, Supplement 1-34*).

[0299] As the expression vector for animal cell, any vector can be used, so long as a gene encoding the C region of a human antibody can be inserted therein and expressed therein. Examples include pAGE107 [*Cytotechnology*, 3, 133 (1990)], pAGE103 [*J. Biochem.*, 101, 1307 (1987)], pHSG274 [*Gene*, 27, 223 (1984)], pKCR [*Proc. Natl. Acad. Sci. USA*, 78, 1527 (1981)], pSGI β d2-4 [*Cytotechnology*, 4, 173 (1990)] and the like. The promoter and enhancer in the expression vector for animal cell include SV40 early promoter and enhancer [*J. Biochem.*, 101, 1307 (1987)], Moloney mouse leukemia virus LTR [*Biochem. Biophys. Res. Commun.*, 149, 960 (1987)], immunoglobulin H chain promoter [*Cell*, 41, 479 (1985)] and enhancer [*Cell*, 33, 717 (1983)], and the like.

(2) Preparation of DNA encoding Fc region of human antibody and protein to be fused

[0300] A DNA encoding the Fc region of a human antibody and the protein to be fused can be obtained in the following manner.

[0301] A cDNA is synthesized from mRNA extracted from a cell or tissue which expresses the protein of interest to be fused with Fc. The synthesized cDNA is cloned into a vector such as a phage or a plasmid to obtain a cDNA library. A recombinant phage or recombinant plasmid comprising cDNA encoding the protein of interest is isolated from the library using the gene sequence part of the protein of interest as the probe. A full nucleotide sequence of the antibody of interest on the recombinant phage or recombinant plasmid is determined, and a full length amino acid sequence is deduced from the nucleotide sequence.

[0302] As the non-human animal, any animal such as mouse, rat, hamster or rabbit can be used, so long as a cell or tissue can be removed therefrom.

[0303] The method for preparing a total RNA from a cell or tissue includes the guanidine thiocyanate-caesium trifluoroacetate method [*Methods in Enzymology*, 154, 3 (1987)] and the like, and the method for preparing mRNA from total RNA includes an oligo (dT)-immobilized cellulose column method (*Molecular Cloning*, Second Edition) and the like. In addition, a kit for preparing mRNA from a cell or tissue includes Fast Track mRNA Isolation Kit (manufactured by Invitrogen), Quick Prep mRNA Purification Kit (manufactured by Pharmacia) and the like.

[0304] The method for synthesizing a cDNA and preparing a cDNA library includes the usual methods (*Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*, Supplement 1-34); methods using a commercially available kit such as SuperScript™, Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by GIBCO BRL) or ZAP-cDNA Synthesis Kit (manufactured by Stratagene); and the like.

[0305] In preparing the cDNA library, the vector into which a cDNA synthesized by using mRNA extracted from a cell or tissue as the template is inserted may be any vector so long as the cDNA can be inserted. Examples include ZAP Express [*Strategies*, 5, 58 (1992)], pBluescript II SK(+) [*Nucleic Acids Research*, 17, 9494 (1989)], λzapII (manufactured by Stratagene), λgt10 and λgt11 [*DNA Cloning, A Practical Approach*, 1, 49 (1985)], Lambda BlueMid (manufactured by Clontech), λExCell, pT7T3 18U (manufactured by Pharmacia), pcD2 [*Mol. Cell. Biol.*, 3, 280 (1983)], pUC18 [*Gene*, 33, 103 (1985)] and the like.

[0306] As *Escherichia coli* into which the cDNA library constructed from a phage or plasmid vector is introduced, any *Escherichia coli* can be used, so long as the cDNA library can be introduced, expressed and maintained. Examples include XL1-Blue MRF' [*Strategies*, 5, 81 (1992)], C600 [*Genetics*, 39, 440 (1954)], Y1088 and Y1090 [*Science*, 222, 778 (1983)], NM522 [*J. Mol. Biol.*, 166, 1 (1983)], K802 [*J. Mol. Biol.*, 16, 118 (1966)], JM105 [*Gene*, 38, 275 (1985)] and the like.

[0307] As the method for selecting a cDNA clone encoding the protein of interest from the cDNA library, a colony hybridization or a plaque hybridization using an isotope- or fluorescence-labeled probe can be used (*Molecular Cloning*, Second Edition). The cDNA encoding the protein of interest can also be prepared by preparing primers and using a cDNA synthesized from mRNA or a cDNA library as the template according to PCR.

[0308] The method for fusing the protein of interest with the Fc region of a human antibody includes PCR. For example, synthesized oligo DNAs (primers) are designed at the 5'-terminal and 3'-terminal of the gene sequence encoding the protein of interest, and PCR is carried out to prepare a PCR product. In the same manner, primers are designed for the gene sequence encoding the Fc region of a human antibody to be fused to prepare a PCR product. At this time, the primers are designed in such a manner that the same restriction enzyme site or the same gene sequence is present between the 3'-terminal of the PCR product of the protein to be fused and the 5'-terminal of the PCR product of the Fc region. When it is necessary to modify the amino acids around the linked site, mutation is introduced by using the primer into which the mutation is introduced. PCR is further carried out by using the two kinds of the obtained PCR fragments to link the genes. Also, they can be linked by carrying out ligation after treatment with the same restriction enzyme.

[0309] The nucleotide sequence of the DNA can be determined by digesting the gene sequence linked by the above method with appropriate restriction enzymes, cloning the fragments into a plasmid such as pBluescript SK(-) (manufactured by Stratagene), carrying out analysis by using a generally used nucleotide sequence analyzing method such as the dideoxy method of Sanger *et al.* [*Proc. Natl. Acad. Sci. USA*, 74, 5463 (1977)] or an automatic nucleotide sequence analyzer such as ABI PRISM 377 DNA Sequencer (manufactured by Pharmacia).

[0310] Whether or not the obtained cDNA encodes the full length amino acid sequences of the Fc fusion protein containing a secretory signal sequence can be confirmed by deducing the full length amino acid sequence of the Fc fusion protein from the determined nucleotide sequence and comparing it with the amino acid sequence of interest.

(3) Stable production of Fc fusion protein

[0311] A transformant capable of stably producing an Fc fusion protein can be obtained by introducing the Fc fusion

protein expression vector described in the item (1) into an appropriate animal cell.

[0312] The method for introducing the Fc fusion protein expression vector into an animal cell include electroporation [Japanese Published Unexamined Patent Application No. 257891/90, *Cytotechnology*, 3, 133 (1990)] and the like.

[0313] As the animal cell into which the Fc fusion protein expression vector is introduced, any cell can be used, so long as it is an animal cell which can produce the Fc fusion protein.

[0314] Examples include mouse myeloma cells such as NS0 cell and SP2/0 cell, Chinese hamster ovary cells such as CHO/*dhfr* cell and CHO/DG44 cell, rat myeloma such as YB2/0 cell and IR983F cell, BHK cell derived from a syrian hamster kidney, a human myeloma cell such as Namalwa cell, and the like, and preferred are a Chinese hamster ovary cell CHO/DG44 cell, a rat myeloma YB2/0 cell and the host cells used in the method of the present invention described in the item 1.

[0315] After introduction of the Fc fusion protein expression vector, a transformant capable of stably producing the Fc fusion protein expression vector can be selected by using a medium for animal cell culture comprising an agent such as G418 and the like in accordance with the method described in Japanese Published Unexamined Patent Application No. 257891/90. The medium to culture animal cells includes RPMI 1640 medium (manufactured by Nissui Pharmaceutical), GIT medium (manufactured by Nihon Pharmaceutical), EX-CELL 302 medium (manufactured by JRH), IMDM medium (manufactured by GIBCO BRL), Hybridoma-SFM medium (manufactured by GIBCO BRL), media obtained by adding various additives such as fetal bovine serum to these media, and the like. The Fc fusion protein can be produced and accumulated in the culture medium by culturing the obtained transformant in a medium. The production amount and antigen binding activity of the Fc fusion protein in the culture medium can be measured by a method such as ELISA. Also, the amount of the Fc fusion protein produced by the transformant can be increased by using a *dhfr* gene amplification system in accordance with the method described in Japanese Published Unexamined Patent Application No. 257891/90.

[0316] The Fc fusion protein can be purified from a culture supernatant culturing the transformant using a protein A column (*Antibodies*, Chapter 8; *Monoclonal Antibodies*). In addition, purification methods generally used for purifying proteins can also be used. For example, the purification can be carried out through the combination of a gel filtration, an ion exchange chromatography and an ultrafiltration. The molecular weight as a whole of the purified Fc fusion protein molecule can be measured by SDS-PAGE [*Nature*, 227, 680 (1970)], Western blotting (*Antibodies*, Chapter 12; *Monoclonal Antibodies*) or the like.

[0317] Thus, methods for producing an antibody composition using an animal cell as the host cell have been described, but, as described above, it can also be produced by yeast, an insect cell, a plant cell, an animal individual or a plant individual by the same methods on the animal cell.

[0318] When the host cell is capable of preparing the antibody molecule, the antibody composition of the present invention can be prepared by culturing the cell capable of expressing an antibody molecule according to the method described in the above item 1, culturing the cell, and recovering the antibody composition of interest.

4. Measurement of binding activity to human FcγRIIIa

[0319] Binding activity of the antibody composition to FcγRIIIa can be measured by the following technique.

(1) Preparation of human FcγRIIIa

[0320] FcγRIIIa which can be used includes FcγRIIIa present on the cell surface of peripheral blood lymphocyte of a human or non-human animal, FcγRIIIa obtained by preparing a gene encoding FcγRIIIa and introducing the gene into a host cell and expressing the FcγR on the cell surface, FcγRIIIa secreted from the cell, and the like.

[0321] A method for preparing a gene encoding FcγRIIIa, introducing the gene into a host cell and expressing the FcγRIIIa on the cell surface, and a method obtaining FcγRIIIa by secreting it from the cell are described below.

[0322] A total RNA or mRNA is prepared from human or non-human animal tissues or cells.

[0323] A commercially available product (e.g., manufactured by Clontech) can be used as the mRNA of human or non-human animal tissues or cells, or it may be prepared from human or non-human animal tissues or cells as follows. The method for preparing a total RNA from human or non-human animal tissues or cells includes the guanidine thiocyanate-cesium trifluoroacetate method [*Methods in Enzymology*, 154, 3 (1987).], the acidic guanidine thiocyanate phenol chloroform (AGPC) method [*Analytical Biochemistry*, 162, 156 (1987); *Experimental Medicine (Jikken Igaku)*, 9, 193 7 (1991)] and the like.

[0324] Also, the method for preparing mRNA as poly(A)⁺ RNA from a total RNA includes an oligo(dT)-immobilized cellulose column method (*Molecular Cloning*, Second Edition) and the like.

[0325] In addition, mRNA can be prepared by using a kit such as Fast Track mRNA Isolation Kit (manufactured by Invitrogen) or Quick Prep mRNA Purification Kit (manufactured by Pharmacia).

[0326] A cDNA library is prepared from a full RNA or mRNA of the prepared human or non-human animal tissue or cell.

[0327] The method for preparing a cDNA library include methods described in *Molecular Cloning*, Second Edition; *Current Protocols in Molecular Biology*; and the like, methods using a commercially available kit such as Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning (manufactured by Life Technologies) or ZAP-cDNA Synthesis Kit (manufactured by STRATAGENE), and the like.

[0328] As the cloning vector for the preparation of the cDNA library, any vector such as a phage vector or a plasmid vector can be used, so long as it is autonomously replicable in *Escherichia coli* K12. Examples include ZAP Express [manufactured by STRATAGENE, *Strategies*, 5, 58 (1992)], pBluescript II SK(+) [*Nucleic Acids Research*, 17, 9494 (1989)], Lambda ZAP II (manufactured by STRATAGENE), λ gt10 and λ gt11 [DNA Cloning, A Practical Approach, 1, 49 (1985)], λ TriplEx (manufactured by Clontech), λ ExCell (manufactured by Pharmacia), pcD2 [*Mol. Cell. Biol.*, 3, 280 (1983)], pUC18 [*Gene*, 33, 103 (1985)] and the like.

[0329] Any microorganism can be used as the host microorganism, and *Escherichia coli* is preferably used. Examples include *Escherichia coli* XL1-Blue MRF' [manufactured by STRATAGENE, *Strategies*, 5, 81 (1992)], *Escherichia coli* C600 [*Genetics*, 39, 440 (1954)], *Escherichia coli* Y1088 [*Science*, 222, 778 (1983)], *Escherichia coli* Y1090 [*Science*, 222, 778 (1983)], *Escherichia coli* NM522 [*J. Mol. Biol.*, 166, 1 (1983)], *Escherichia coli* K802 [*J. Mol. Biol.*, 16, 118 (1966)], *Escherichia coli* JM105 [*Gene*, 38, 275 (1985)] and the like.

[0330] The cDNA library may be used as such in the succeeding analysis, and in order to obtain a full length cDNA as efficient as possible by decreasing the ratio of an infull length cDNA, a cDNA library prepared using the oligo cap method developed by Sugano *et al.* [*Gene*, 138, 171 (1994); *Gene*, 200, 149 (1997); *Protein, Nucleic Acid and Enzyme*, 41, 603 (1996); *Experimental Medicine*, 11, 2491 (1993); *cDNA Cloning* (Yodo-sha) (1996); *Methods for Preparing Gene Libraries* (Yodo-sha) (1994)] may be used in the following analysis.

[0331] A gene encoding Fc γ R can be obtained by preparing primers specific for 5'-terminal and 3'-terminal nucleotide sequences based on the nucleotide sequences of various Fc γ RIIIa, and amplifying DNA by PCR [*PCR Protocols*, Academic Press (1990)] using a prepared cDNA library as the template.

[0332] Whether the thus obtained gene is a DNA encoding Fc γ RIIIa can be confirmed by analyzing it according to the generally used nucleotide sequence analyzing method such as the dideoxy method of Sanger *et al.* [*Proc. Natl. Acad. Sci. U.S.A.*, 74, 5463 (1977)] or by using a nucleotide sequence analyzer such as ABI PRISM 377 DNA Sequencer (manufactured by PE Biosystems).

[0333] The nucleotide sequence of a gene encoding Fc γ RIIIa obtained by the above method includes the nucleotide sequence of Fc γ RIIIa represented by SEQ ID NO:27.

[0334] The gene encoding Fc γ RIIIa can also be obtained based on the determined DNA nucleotide sequence by carrying out chemical synthesis by a DNA synthesizer such as DNA Synthesizer Model 392 manufactured by Perkin Elmer using a phosphoramidite method.

[0335] A recombinant vector is prepared by inserting the thus obtained cDNA encoding Fc γ RIIIa into downstream of the promoter of an appropriate expression vector.

[0336] A transformant which produces an antibody molecule can be obtained by introducing the recombinant vector into a host cell suitable for the expression vector.

[0337] As the host cell, any of yeast, an animal cell, an insect cell, a plant cell or the like can be used, so long as it can express the gene of interest.

[0338] As the expression vector, a vector which is autonomously replicable in the above host cell or can be integrated into the chromosome and comprises a promoter at such a position that the DNA encoding the Fc γ RIIIa of interest can be transferred is used.

[0339] When a yeast is used as the host cell, the expression vector includes YEP 13 (ATCC 37115), YEp24 (ATCC 37051), YCp50 (ATCC 37419) and the like.

[0340] Any promoter can be used, so long as it can function in yeast. Examples include a promoter of a gene of the glycolytic pathway such as a hexose kinase gene, PHO5 promoter, PGK promoter, GAP promoter, ADH promoter, gal 1 promoter, gal 10 promoter, heat shock protein promoter, MF α 1 promoter, CUP 1 promoter and the like.

[0341] The host cell includes microorganisms belonging to the genus *Saccharomyces*, the genus *Schizosaccharomyces*, the genus *Kluyveromyces*, the genus *Trichosporon*, the genus *Schwanniomyces* and the like, such as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Trichosporon pullulans* and *Schwanniomyces alluvius*.

[0342] As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into yeast. Examples include electroporation [*Methods in Enzymology*, 194, 182 (1990)], spheroplast method [*Proc. Natl. Acad. Sci. USA*, 84, 1929 (1978)], lithium acetate method [*J. Bacteriol.*, 153, 163 (1983)], a method described in *Proc. Natl. Acad. Sci. USA*, 75, 1929 (1978) and the like.

[0343] When an animal cell is used as the host, the expression vector includes pcDNA1, pcDM8 (available from Funakoshi), pAGE107 [Japanese Published Unexamined Patent Application No. 22979/91; *Cytotechnology*, 3, 133 (1990)], pAS3-3 [Japanese Published Unexamined Patent Application No. 227075/90], pCDM8 [*Nature*, 329, 840 (1987)], pcDNA1/Amp (manufactured by Invitrogen), pREP4 (manufactured by Invitrogen), pAGE103 [*J. Biochemistry*,

101, 1307 (1987)], pAGE210 and the like.

[0344] Any promoter can be used, so long as it can function in an animal cell. Examples include a promoter of IE (immediate early) gene of cytomegalovirus (CMV), an early promoter of SV40, a promoter of retrovirus, a promoter of metallothionein, a heat shock promoter, an SR α promoter and the like. Also, an enhancer of the IE gene of human CMV may be used together with the promoter.

[0345] The host cell includes a human cell such as Namalwa cell, a monkey cell such as COS cell, a Chinese hamster cell such as CHO cell or HBT5637 (Japanese Published Unexamined Patent Application No. 299/88), a rat myeloma cell, a mouse myeloma cell, a cell derived from syrian hamster kidney, an embryonic stem cell, a fertilized egg cell and the like.

[0346] As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into an animal cell. Examples include electroporation [*Cytotechnology*, 3, 133 (1990)], the calcium phosphate method (Japanese Published Unexamined Patent Application No. 227075/90), the lipofection method (*Proc. Natl. Acad. Sci. USA*, 84, 7413 (1987)), the injection method (*Manipulating the Mouse Embryo, A Laboratory Manual*), a method using particle gun (gene gun) (Japanese Patent No. 2606856, Japanese Patent No. 2517813), the DEAE-dextran method [*Biomaterial Series 4-Gene Transfer and Expression Analysis* (Yodo-sha), edited by Takashi Yokota and Kenichi Arai (1994)], the virus vector method (*Manipulating Mouse Embryo*, Second Edition) and the like.

[0347] When an insect cell is used as the host, the protein can be expressed by the method described in *Current Protocols in Molecular Biology, Baculovirus Expression Vectors, A Laboratory Manual*, W.H. Freeman and Company, New York (1992), *BioTechnology*, 6, 47 (1988) or the like.

[0348] That is, the protein can be expressed by co-introducing a recombinant gene-introducing vector and a baculovirus into an insect cell to obtain a recombinant virus in an insect cell culture supernatant and then infecting the insect cell with the recombinant virus.

[0349] The gene introducing vector used in the method includes pVL1392, pVL1393, pBlueBacIII (all manufactured by Invitrogen) and the like.

[0350] The baculovirus includes *Arctographa californica* nuclear polyhedrosis virus which is infected by an insect of the family *Barathra*.

[0351] The insect cell includes *Spodoptera frugiperda* oocytes Sf9 and Sf21 [*Current Protocols in Molecular Biology, Baculovirus Expression Vectors, A Laboratory Manual*, W.H. Freeman and Company, New York (1992)], a *Trichoplusia ni* oocyte High 5 (manufactured by Invitrogen) and the like.

[0352] The method for co-introducing the recombinant gene-introducing vector and the baculovirus for preparing the recombinant virus includes the calcium phosphate method (Japanese Published Unexamined Patent Application No. 227075/90), the lipofection method [*Proc. Natl. Acad. Sci. USA*, 84, 7413 (1987)] and the like.

[0353] When a plant cell is used as the host, the expression vector includes Ti plasmid, tobacco mosaic virus and the like.

[0354] As the promoter, any promoter can be used, so long as it can function in a plant cell. Examples include cauliflower mosaic virus (CaMV) 35S promoter, rice actin 1 promoter and the like.

[0355] The host cell includes plant cells of tobacco, potato, tomato, carrot, soybean, rape, alfalfa, rice, wheat, barley and the like.

[0356] As the method for introducing the recombinant vector, any method can be used, so long as it can introduce DNA into a plant cell. Examples include a method using *Agrobacterium* (Japanese Published Unexamined Patent Application No. 140885/84, Japanese Published Unexamined Patent Application No. 70080/85, WO94/00977), electroporation (Japanese Published Unexamined Patent Application No. 251887/85), a method using a particle gun (gene gun) (Japanese Patent No. 2606856, Japanese Patent No. 2517813) and the like.

[0357] As the method for expressing a gene, secretion production, expression of a fusion protein with other protein and the like can be carried out in accordance with the method described in *Molecular Cloning*, Second Edition or the like, in addition to the direct expression.

[0358] Fc γ R1IIa can be produced by culturing the thus obtained transformant in a medium to produce and accumulate Fc γ R1IIa in the culture and then recovering it from the resulting culture. The method for culturing the transformant using a medium can be carried out in accordance with a general method which is used for culturing host cells.

[0359] As the medium for culturing a transformant obtained by using yeast as the host, the medium may be either a natural medium or a synthetic medium, so long as it comprises materials such as a carbon source, a nitrogen source and an inorganic salt which can be assimilated by the yeast and culturing of the transformant can be efficiently carried out.

[0360] As the carbon source, those which can be assimilated by the yeast can be used. Examples include carbohydrates such as glucose, fructose, sucrose, molasses containing them, starch and starch hydrolysate; organic acids such as acetic acid and propionic acid; alcohols such as ethanol and propanol; and the like.

[0361] The nitrogen source includes ammonia; ammonium salts of inorganic acid or organic acid such as ammonium chloride, ammonium sulfate, ammonium acetate and ammonium phosphate; other nitrogen-containing compounds;

peptone; meat extract; yeast extract; corn steep liquor; casein hydrolysate; soybean meal; soybean meal hydrolysate; various fermented cells and hydrolysates thereof; and the like.

[0362] The inorganic salt includes potassium dihydrogen phosphate, dipotassium hydrogen phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate, calcium carbonate, and the like.

[0363] The culturing is carried out generally under aerobic conditions such as shaking culture or submerged-aeration stirring culture. The culturing temperature is preferably 15 to 40°C, and the culturing time is generally 16 hours to 7 days. During the culturing, the pH is maintained at 3.0 to 9.0. The pH is adjusted with an inorganic or organic acid, an alkali solution, urea, calcium carbonate, ammonia or the like.

[0364] If necessary, an antibiotic such as ampicillin or tetracycline can be added to the medium during the culturing.

[0365] When yeast transformed with a recombinant vector obtained by using an inducible promoter as the promoter is cultured, an inducer can be added to the medium, if necessary. For example, when yeast transformed with a recombinant vector obtained using *lac* promoter is cultured, isopropyl-β-D-thiogalactopyranoside can be added to the medium, and when yeast transformed with a recombinant vector obtained using *trp* promoter is cultured, indoleacrylic acid and the like can be added to the medium.

[0366] When a transformant obtained by using an animal cell as the host cell is cultured, the medium includes generally used RPMI 1640 medium [*The Journal of the American Medical Association*, 199, 519 (1967)], Eagle's MEM medium [*Science*, 122, 501 (1952)], Dulbecco's modified MEM medium [*Virology*, 8, 396 (1959)], 199 medium [*Proceeding of the Society for the Biological Medicine*, 73, 1 (1950)] and Whitten's medium [*Developmental Engineering Experimentation Manual (Hasei Kogaku Jikken Manual)-Preparation of Transgenic Mice* (Kodan-sha), edited by M. Katsuki (1987)], the media to which fetal calf serum, etc, is added, and the like.

[0367] The culturing is carried out generally at a pH of 6.0 to 8.0 and 30 to 40°C for 1 to 7 days in the presence of 5% CO₂. If necessary, an antibiotic such as kanamycin or penicillin can be added to the medium during the culturing.

[0368] The medium for the culturing of a transformant obtained by using an insect cell as the host includes generally used TNM-FH medium (manufactured by Pharmingen), Sf-900 II SFM medium (manufactured by Life Technologies), ExCell 400 and ExCell 405 (both manufactured by JRH Biosciences), Grace's Insect Medium [*Nature*, 195, 788 (1962)] and the like.

[0369] The culturing is carried out generally at a medium pH of 6.0 to 7.0 and 25 to 30°C for 1 to 5 days.

[0370] In addition, antibiotics such as gentamicin can be added to the medium during the culturing, if necessary.

[0371] A transformant obtained by using a plant cell as the host cell can be cultured as a cell or by differentiating it into a plant cell or organ. The medium for culturing the transformant includes generally used Murashige and Skoog (MS) medium and White medium, the media to which a plant hormone such as auxin or cytokinin is added, and the like.

[0372] The culturing is carried out generally at a pH of 5.0 to 9.0 and 20 to 40°C for 3 to 60 days.

[0373] If necessary, an antibiotic such as kanamycin or hygromycin can be added to the medium during the culturing.

[0374] As discussed above, FcγRIIIa can be produced by culturing a transformant derived from a microorganism, an animal cell, an insect cell or a plant cell which comprises a recombinant vector into which a DNA encoding FcγRIIIa is inserted, in accordance with the general culturing method to thereby produce and accumulate FcγRIIIa, and then recovering FcγRIIIa from the culture.

[0375] As the method for expressing FcγRIIIa, secretion production, expression of a fusion protein and the like can be carried out in accordance with the method described in *Molecular Cloning*, Second Edition in addition to the direct expression.

[0376] The method for producing FcγRIIIa includes a method of intracellular expression in a host cell, a method of extracellular secretion from a host cell, and a method of production on a host cell membrane outer envelope. The method can be selected by changing the host cell used or the structure of FcγRIIIa produced.

[0377] When FcγRIIIa is produced in a host cell or on a host cell membrane outer envelope, it can be positively secreted extracellularly in accordance with the method of Paulson *et al.* [*J. Biol. Chem.*, 264, 17619 (1989)], the method of Lowe *et al.* [*Proc. Natl. Acad. Sci. USA*, 86, 8227 (1989)], *Genes Develop.*, 4, 1288 (1990)], the methods described in Japanese Published Unexamined Patent Application No. 336963/93 and Japanese Published Unexamined Patent Application No. 823021/94 and the like.

[0378] That is, FcγRIIIa of interest can be positively secreted extracellularly from a host cell by inserting a DNA encoding FcγRIIIa and a signal peptide suitable for the expression of FcγRIIIa into an expression vector by using a gene recombination technique, and then expressing the vector.

[0379] Also, its production amount can be increased in accordance with the method described in Japanese Published Unexamined Patent Application No. 227075/90 based on a gene amplification system using a dihydrofolate reductase gene.

[0380] In addition, FcγR can also be produced by using a gene-introduced animal individual (transgenic non-human animal) or a plant individual (transgenic plant) which is constructed by the redifferentiation of an animal or plant cell into which the gene is introduced.

[0381] When the transformant is an animal individual or a plant individual, FcγRIIIa can be produced in accordance with a general method by rearing or cultivating it to thereby produce and accumulate FcγR and then recovering FcγRIIIa from the animal or plant individual.

[0382] The method for producing FcγRIIIa by using an animal individual includes a method in which FcγRIIIa of interest is produced in an animal constructed by introducing a gene in accordance with a known method [*American Journal of Clinical Nutrition*, 63, 639 (1996); *American Journal of Clinical Nutrition*, 63, 627S (1996); *Bio Technology*, 9, 830 (1991)].

[0383] In the case of an animal individual, FcγRIIIa can be produced by rearing a transgenic non-human animal into which a DNA encoding FcγRIIIa is introduced to thereby produce and accumulate FcγRIIIa in the animal, and then recovering FcγRIIIa from the animal. The place of the animal where FcγRIIIa is produced and accumulated includes milk (Japanese Published Unexamined Patent Application No. 309192/88) and eggs of the animal. As the promoter used in this case, any promoter can be used, so long as it can function in an animal. Preferred examples include mammary gland cell-specific promoters such as α casein promoter, β casein promoter, β lactoglobulin promoter and whey acidic protein promoter.

[0384] The method for producing FcγRIIIa by using a plant individual includes a method in which FcγRIIIa is produced by cultivating a transgenic plant into which a DNA encoding FcγR is introduced by a known method [*Tissue Culture (Soshiki Baiyo)*, 20 (1994); *Tissue Culture (Soshiki Baiyo)*, 21 (1995); *Trends in Biotechnology*, 15, 45 (1997)] to produce and accumulate FcγR in the plant, and then recovering FcγRIIIa from the plant.

[0385] Regarding purification of FcγRIIIa produced by a transformant into which a gene encoding FcγRIIIa is introduced, for example, when FcγRIIIa is intracellularly expressed in a dissolved state, the cells after culturing are recovered by centrifugation, suspended in an aqueous buffer and then disrupted using ultrasonic oscillator, French press, Manton Gaulin homogenizer, dynamill or the like to obtain a cell-free extract. A purified product of FcγRIIIa can be obtained from a supernatant obtained by centrifuging the cell-free extract, by using an ordinary enzyme isolation purification technique such as solvent extraction; salting out and desalting with ammonium sulfate, *etc.*; precipitation with an organic solvent; anion exchange chromatography using a resin such as diethylaminoethyl (DEAE)-Sepharose or DIAION HPA-75 (manufactured by Mitsubishi Chemical); cation exchange chromatography using a resin such as S-Sepharose FF (manufactured by Pharmacia); hydrophobic chromatography using a resin such as butyl-Sepharose or phenyl-Sepharose; gel filtration using a molecular sieve; affinity chromatography; chromatofocusing; electrophoresis such as isoelectric focusing; and the like which may be used alone or in combination.

[0386] Also, when FcγRIIIa is expressed intracellularly by forming an insoluble body, the cells are recovered, disrupted and centrifuged in the same manner, and the insoluble body of FcγRIIIa is recovered as a precipitation fraction. The recovered insoluble body of FcγRIIIa is solubilized by using a protein denaturing agent. FcγRIIIa is made into a normal three-dimensional structure by diluting or dialyzing the solubilized solution, and then a purified product of FcγRIIIa is obtained by the same isolation purification method.

[0387] When FcγRIIIa is secreted extracellularly, FcγRIIIa or derivatives thereof can be recovered from the culture supernatant. That is, the culture is treated by a similar technique such as centrifugation to obtain a soluble fraction, and a purified preparation of FcγRIIIa can be obtained from the soluble fraction by the same isolation purification method.

(2) Measurement of binding activity to FcγRIIIa

[0388] Binding activity of the antibody composition to FcγRIIIa expressed on the cell membrane can be measured by the immunofluorescent method [*Cancer Immunol. Immunother.*, 36, 373 (1993)] or the like. Also, binding activity to the purified FcγRIIIa prepared by the method described in the item 4(1) can be measured according to the immunological determination method such as Western staining described in literatures [*Monoclonal Antibodies: Principles and Applications*, Wiley-Liss, Inc., (1995); *Enzyme Immunoassay (Koso Men-eki Sokutei Ho)*, 3rd edition, Igaku Shoin (1987), reversed edition; *Enzyme Antibody Method (Koso Kotai Ho)*, Gakusai Kikaku (1985)], RIA (radioimmunoassay), VIA (viroimmunoassay), EIA (enzyme immunoassay), FIA (fluoroimmunoassay) or MIA (metalloimmunoassay), for example, as follows.

[0389] FcγRIIIa is immobilized on a plastic plate for EIA and is allowed to react with a sample containing an antibody composition. Next, an amount of the bound antibody composition is measured by using an appropriate secondary antibody.

[0390] In addition, binding activity to the purified FcγRIIIa can also be measured by a measuring method using biosensor [e.g., BIAcore (manufactured by BIACORE)] [*J. Immunol. Methods*, 200, 121 (1997)], isothermal titration calorimetry [*Proc. Natl. Acad. Sci. U.S.A.*, 97, 9026 (2000)] or the like.

5. Activity evaluation of antibody composition

[0391] As the method for measuring the amount of the purified antibody composition, its binding activity to an antigen, its binding activity to FcγRIIIa and its effector function, the known method described in *Monoclonal Antibodies, Antibody Engineering* and the like can be used.

[0392] For example, when the antibody composition is a humanized antibody, the binding activity to an antigen, binding activity to an antigen-positive cultured clone and binding activity to FcγRIIIa can be measured by methods such as ELISA and the immunofluorescent method [*Cancer Immunol. Immunother.*, 36, 373 (1993)], measurement using biosensor [for example, using BIAcore (manufactured by BIACORE)] [*J. Immunol. Methods*, 200, 121 (1997)], isothermal titration calorimetry method [*Proc. Natl. Acad. Sci. USA.*, 97, 9026 (2000)] and the like. Among the effector functions, the cytotoxic activity against an antigen-positive cultured clone can be evaluated by measuring CDC activity, ADCC activity [*Cancer Immunol. Immunother.*, 36, 373 (1993)] and the like.

6. Analysis of sugar chains of antibody molecule expressed in various cells

[0393] The sugar chain structure binding to an antibody molecule expressed in various cells can be analyzed in accordance with the general analysis of the sugar chain structure of a glycoprotein. For example, the sugar chain which is bound to IgG molecule comprises a neutral sugar such as galactose, mannose, fucose, an amino sugar such as *N*-acetylglucosamine and an acidic sugar such as sialic acid, and can be analyzed by a method such as a sugar chain structure analysis using sugar composition analysis, two dimensional sugar chain mapping or the like.

(1) Analysis of neutral sugar and amino sugar compositions

[0394] The composition analysis of the sugar chain of an antibody molecule can be carried out by acid hydrolysis of sugar chains with trifluoroacetic acid or the like to release a neutral sugar or an amino sugar and measuring the composition ratio.

[0395] Examples include a method using a sugar composition analyzer (BioLC) manufactured by Dionex. The BioLC is an apparatus which analyzes a sugar composition by HPAEC-PAD (high performance anion-exchange chromatography-pulsed amperometric detection) [*J. Liq. Chromatogr.*, 6, 1577 (1983)].

[0396] The composition ratio can also be analyzed by a fluorescence labeling method using 2-aminopyridine. Specifically, the composition ratio can be calculated in accordance with a known method [*Agric. Biol. Chem.*, 55(1), 283 (1991)] by labeling an acid-hydrolyzed sample with a fluorescence by 2-aminopyridylation and then analyzing the composition by HPLC.

(2) Analysis of sugar chain structure

[0397] The sugar chain structure binding to an antibody molecule can be analyzed by the two dimensional sugar chain mapping method [*Anal. Biochem.*, 171, 73 (1988), *Biochemical Experimentation Methods 23 - Methods for Studying Glycoprotein Sugar Chains* (Japan Scientific Societies Press) edited by Reiko Takahashi (1989)]. The two dimensional sugar chain mapping method is a method for deducing a sugar chain structure by, e.g., plotting the retention time or elution position of a sugar chain by reverse phase chromatography as the X axis and the retention time or elution position of the sugar chain by normal phase chromatography as the Y axis, respectively, and comparing them with those of known sugar chains.

[0398] Specifically, sugar chains are released from an antibody by subjecting the antibody to hydrazinolysis, and the released sugar chains are subjected to fluorescence labeling with 2-aminopyridine (hereinafter referred to as "PA") [*J. Biochem.*, 95, 197 (1984)], and then the sugar chains are separated from an excess PA-treating reagent by gel filtration, and subjected to reverse phase chromatography. Thereafter, each peak of the separated sugar chains are subjected to normal phase chromatography. From these results, the sugar chain structure can be deduced by plotting the results on a two dimensional sugar chain map and comparing them with the spots of a sugar chain standard (manufactured by Takara Shuzo) or a literature [*Anal. Biochem.*, 171, 73 (1988)].

[0399] The structure deduced by the two dimensional sugar chain mapping method can be confirmed by further carrying out mass spectrometry such as MALDI-TOF-MS of each sugar chain.

7. Immunological determination method for identifying sugar chain structure of antibody molecule

[0400] An antibody composition comprises an antibody molecule in which sugar chains binding to the Fc region of the antibody are different in structure. The antibody composition comprising a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among the total complex *N*-glycoside-linked sugar chains

binding to the Fc region in the antibody composition reducing end can be identified by using the method for analyzing the sugar chain structure binding to an antibody molecule described in the item 6. Also, it can also be identified by an immunological determination method using a lectin.

[0401] The sugar chain structure binding to an antibody molecule can be identified by the immunological determination method using a lectin in accordance with the known immunological determination method such as Western staining, IRA (radioimmunoassay), VIA (viroimmunoassay), EIA (enzymimmunoassay), FIA (fluoroimmunoassay) or MIA (metalloimmunoassay) described in literatures [*Monoclonal Antibodies: Principles and Applications*, Wiley-Liss, Inc. (1995); *Immunoassay (Koso Meneki Sokuteiho)*, 3rd Ed., Igakushoin (1987); Revised Edition, *Enzyme Antibody Method (Koso Kotaiho)*, Gakusai Kikaku (1985)] and the like. A lectin which recognizes the sugar chain structure binding to an antibody molecule comprised in an antibody composition is labeled, and the labeled lectin is allowed to react with a sample antibody composition. Then, the amount of the complex of the labeled lectin with the antibody molecule is measured.

[0402] The lectin used for identifying the sugar chain structure binding to an antibody molecule includes WGA (wheat-germ agglutinin derived from *T. vulgaris*), ConA (concanavalin A derived from *C. ensiformis*), RIC (toxin derived from *R. communis*), L-PHA (leucoagglutinin derived from *P. vulgaris*), LCA (lentil agglutinin derived from *L. culinaris*), PSA (pea lectin derived from *P. sativum*), AAL (*Aleuria aurantia* lectin), ACL (*Amaranthus caudatus* lectin), BPL (*Bauhinia purpurea* lectin), DSL (*Datura stramonium* lectin), DBA (*Dolichos biflorus* agglutinin), EBL (elderberry balk lectin), ECL (*Erythrina cristagalli* lectin), EEL (*Euonymus europaeus* lectin), GNL (*Galanthus nivalis* lectin), GSL (*Griffonia simplicifolia* lectin), HPA (*Helix pomatia* agglutinin), HHL (*Hippeastrum* hybrid lectin), Jacalin, LTL (*Lotus tetragonolobus* lectin), LEL (*Lycopersicon esculentum* lectin), MAL (*Maackia amurensis* lectin), MPL (*Maclura pomifera* lectin), NPL (*Narcissus pseudonarcissus* lectin), PNA (peanut agglutinin), E-PHA (*Phaseolus vulgaris* erythroagglutinin), PTL (*Psophocarpus tetragonolobus* lectin), RCA (*Ricinus communis* agglutinin), STL (*Solanum tuberosum* lectin), SJA (*Sophora japonica* agglutinin), SBA (soybean agglutinin), UEA (*Ulex europaeus* agglutinin), VVL (*Vicia villosa* lectin) and WFA (*Wisteria floribunda* agglutinin).

[0403] It is preferable to use a lectin which specifically recognizes a sugar chain structure wherein fucose binds to the *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain. Examples include *Lens culinaris* lectin LCA (lentil agglutinin derived from *Lens culinaris*), pea lectin PSA (pea lectin derived from *Pisum sativum*), broad bean lectin VFA (agglutinin derived from *Vicia faba*) and *Aleuria aurantia* lectin AAL (lectin derived from *Aleuria aurantia*).

8. Method for measuring binding activity of antibody composition to FcγRIIIa

[0404] The present invention relates to a method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in an antibody composition by using a measuring method which comprises reacting an antigen with a tested antibody composition to form a complex of the antigen and the antibody composition; contacting the complex with an Fcγ receptor IIIa. Furthermore, the present invention relates to a method for detecting the antibody-dependent cell-mediated cytotoxic activity.

[0405] The measuring method used in the present invention is described below in detail.

[0406] First, an antigen is fixed on a plate, and a sample of antibody composition is allowed to react with it. The resulting complex after the antigen-antibody reaction is allowed to react with human FcγRIIIa.

[0407] The human FcγRIIIa to be allowed to react is labeled with a label such as an enzyme, a radioisotope or an fluorescent, and the binding activity of the antibody bound to the antigen can be measured by an immunological measuring method.

[0408] The immunological measuring method includes any method which uses an antigen-antibody reaction such as an immunoassay, an immunoblotting, a coagulation reaction, a complement binding reaction, a hemolysis reaction, a precipitation reaction, a colloidal gold method, a chromatography or an immune staining method. Among these, the immunoassay is preferred.

[0409] Also, human FcγRIIIa having a tag can be obtained by ligating a nucleotide sequence encoding a short peptide to a gene encoding the human FcγRIIIa, and expressing the product by genetic engineering techniques. The tag includes histidine and the like.

[0410] Accordingly, when the above reaction is carried out by using the human FcγRIIIa having a tag, an immunoassay having high sensitivity can be carried out by applying an antibody against the tag after the reaction and labeling the antibody against the tag as described above, or using a labeled antibody which binds to the antibody against the tag.

[0411] Also, human FcγRIIIa having a tag can be obtained by ligating a nucleotide sequence encoding a short peptide to a gene encoding the human FcγRIIIa, and expressing the product by genetic engineering techniques. The tag includes histidine and the like.

[0412] Accordingly, when the above reaction is carried out by using the human FcγRIIIa having a tag, an immunoassay having high sensitivity can be carried out by applying an antibody against the tag after the reaction and labeling

the antibody against the tag as described above, or using a labeled antibody which binds to the antibody against the tag.

[0413] The detection method of the present invention can also be carried out by directly contacting a sample of antibody composition with FcγRIIIa, without reacting an antigen with the sample of antibody component. For example, a labeled antibody which recognizes the human Fc region can be detected by solid-phasing an antibody against the tag and reacting the antibody with tag-containing FcγRIIIa, followed by reaction with a tested antibody composition.

[0414] The method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end of an antibody composition can be carried out according to the following method.

[0415] First, antibody compositions (standards) necessary for the preparation of a standard curve having a different ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the sugar chain reducing end are prepared and the sugar chain analysis of an antibody composition is carried out. In this case, the concentrations of the antibody compositions are adjusted to the same. Each of the binding activities of the prepared antibody composition samples to FcγRIIIa is measured, and a standard curve between the ratio of sugar chain and the binding activity to FcγRIIIa is prepared.

[0416] Based on the above standard curve, the binding activity of the antibody composition sample to be measured to FcγRIIIa is measured while keeping the definite concentration of the antibody composition according to a measuring method similar to the above,

[0417] Based on the above standard curve, the binding activity of the antibody composition to FcγRIIIa is measured with the same concentration of the sample to be measured according to the measuring method similar to the above, so that the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the sugar chain reducing end in the antibody composition sample can be obtained.

[0418] Furthermore, detection of ADCC activity can be carried out by the following method.

[0419] ADCC activities of the standards used for the preparation of the standard curve in the method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the sugar chain reducing end of the above antibody composition are measured. The measuring method includes the method for measuring ADCC described below. Each of binding activities of the prepared antibody composition samples is measured by the above measuring method, and a standard curve between the ADCC activity and the binding activity to FcγRIIIa is prepared.

[0420] Based on the above standard curve, the binding activity of the antibody composition sample to be measured to FcγRIIIa is measured while keeping the definite concentration of the antibody composition according to the measuring method used above, so that the ADCC activity can be obtained. 8 Screening method for antibody composition

[0421] The present invention relates to a method for screening an antibody composition having a higher binding activity to FcRIIIa, which comprises reacting an antigen with a tested antibody, followed by treatment with FcγRIIIa.

[0422] The screening method of the present invention is described below in detail.

[0423] An antigen is fixed on a plate and allowed to react with an antibody to be tested. Human FcγRIIIa is allowed to react with the complex of the antigen-antibody reaction.

[0424] The human FcγRIIIa to be allowed to react is labeled with a label such as an enzyme, a radioisotope or an fluorescent, and the binding activity of the antibody bound to the antigen can be measured by an immunological measuring method.

[0425] The immunological measuring method includes any method which uses an antigen-antibody reaction such as an immunoassay, an immunoblotting, a coagulation reaction, a complement binding reaction, a hemolysis reaction, a precipitation reaction, a colloidal gold method, a chromatography or an immune staining method. Among these, the immunoassay is preferred.

[0426] Also, human FcγRIIIa having a tag can be obtained by ligating a nucleotide sequence encoding a short peptide to a gene encoding the human FcγRIIIa, and expressing the product by genetic engineering techniques. The tag includes histidine and the like.

[0427] Accordingly, when the above reaction is carried out by using the human FcγRIIIa having a tag, an immunoassay having high sensitivity can be carried out by applying an antibody against the tag after the above reaction and labeling the antibody against the tag as described above, or using a labeled antibody which binds to the antibody against the tag.

10. Application of antibody composition of the present invention

[0428] The antibody composition obtained by the screening method of the present invention has high ADCC activity.

[0429] An antibody having high ADCC activity is useful for preventing and treating various diseases including cancers, inflammatory diseases, immune diseases such as autoimmune diseases and allergies, cardiovascular diseases and viral or bacterial infections.

[0430] In the case of cancers, namely malignant tumors, cancer cells grow. General anti-tumor agents are characterized by inhibiting the growth of cancer cells. In contrast, an antibody having high ADCC activity can treat cancers by injuring cancer cells through its cell killing effect, and therefore, it is more effective as a therapeutic agent than the

general anti-tumor agents. At present, in regard to therapeutic agents for cancers, an anti-tumor effect of an antibody medicament alone is insufficient, so that it has been taken to chemotherapy [*Science*, 280, 1197 (1998)]. If higher anti-tumor effect is found by the antibody composition of the present invention alone, the dependency on chemotherapy will be decreased and side effects will be reduced.

[0431] In immune diseases such as inflammatory diseases, autoimmune diseases and allergies, *in vivo* reactions of the diseases are induced by the release of a mediator molecule by immunocytes, so that the allergy reaction can be inhibited by eliminating immunocytes using an antibody having high ADCC activity.

[0432] The cardiovascular diseases include arteriosclerosis and the like. The arteriosclerosis is treated by using balloon catheter at present, but cardiovascular diseases can be prevented and treated by using an antibody having high ADCC activity because growth of arterial cells in restricture after above treatment can be inhibited by using the antibody.

[0433] Various diseases including viral and bacterial infections can be prevented and treated by inhibiting proliferation of cells infected with a virus or bacterium using an antibody having high ADCC activity.

[0434] Specific examples of an antibody which recognizes a tumor-related antigen, an antibody which recognizes an allergy- or inflammation-related antigen, an antibody which recognizes cardiovascular disease-related antigen and an antibody which recognizes a viral or bacterial infection-related antigen are described below.

[0435] The antibody which recognizes a tumor-related antigen includes anti-GD2 antibody [*Anticancer Res.*, 13, 331 (1993)], anti-GD3 antibody [*Cancer Immunol. Immunother.*, 36, 260 (1993)], anti-GM2 antibody [*Cancer Res.*, 54, 1511 (1994)], anti-HER2 antibody [*Proc. Natl. Acad. Sci. USA*, 89, 4285 (1992)], anti-CD52 antibody [*Proc. Natl. Acad. Sci. USA*, 89, 4285 (1992)], anti-MAGE antibody [*British J. Cancer*, 83, 493 (2000)], anti-HM1.24 antibody [*Molecular Immunol.*, 36, 387 (1999)], anti-parathyroid hormone-related protein (PTHrP) antibody [*Cancer*, 88, 2909 (2000)], anti-basic fibroblast growth factor antibody and anti-FGF8 antibody [*Proc. Natl. Acad. Sci. USA*, 86, 9911 (1989)], anti-basic fibroblast growth factor receptor antibody and anti-FGF8 receptor antibody [*J. Biol. Chem.*, 265, 16455 (1990)], anti-insulin-like growth factor antibody [*J. Neurosci. Res.*, 40, 647 (1995)], anti-insulin-like growth factor receptor antibody [*J. Neurosci. Res.*, 40, 647 (1995)], anti-PMSA antibody [*J. Urology*, 160, 2396 (1998)], anti-vascular endothelial cell growth factor antibody [*Cancer Res.*, 57, 4593 (1997)], anti-vascular endothelial cell growth factor receptor antibody [*Oncogene*, 19, 2138 (2000)] and the like.

[0436] The antibody which recognizes an allergy- or inflammation-related antigen includes anti-interleukin 6 antibody [*Immunol. Rev.*, 127, 5 (1992)], anti-interleukin 6 receptor antibody [*Molecular Immunol.*, 31, 371 (1994)], anti-interleukin 5 antibody [*Immunol. Rev.*, 127, 5 (1992)], anti-interleukin 5 receptor antibody and anti-interleukin 4 antibody [*Cytokine*, 3, 562 (1991)], anti-interleukin 4 receptor antibody [*J. Immunol. Methods*, 217, 41 (1998)], anti-tumor necrosis factor antibody [*Hybridoma*, 13, 183 (1994)], anti-tumor necrosis factor receptor antibody [*Molecular Pharmacol.*, 58, 237 (2000)], anti-CCR4 antibody [*Nature*, 400, 776 (1999)], anti-chemokine antibody [*J. Immuno. Meth.*, 174, 249 (1994)], anti-chemokine receptor antibody [*J. Exp. Med.*, 186, 1373 (1997)] and the like. The antibody which recognizes a cardiovascular disease-related antigen includes anti-GpIIb/IIIa antibody [*J. Immunol.*, 152, 2968 (1994)], anti-platelet-derived growth factor antibody [*Science*, 253, 1129 (1991)], anti-platelet-derived growth factor receptor antibody [*J. Biol. Chem.*, 272, 17400 (1997)] and anti-blood coagulation factor antibody [*Circulation*, 101, 1158 (2000)] and the like.

[0437] The antibody which recognizes a viral or bacterial infection-related antigen includes anti-gp120 antibody [*Structure*, 8, 385 (2000)], anti-CD4 antibody [*J. Rheumatology*, 25, 2065 (1998)], anti-CCR5 antibody and anti-Vero toxin antibody [*J. Clin. Microbiol.*, 37, 396 (1999)] and the like.

[0438] These antibodies can be obtained from public organizations such as ATCC (The American Type Culture Collection), RIKEN Gene Bank at The Institute of Physical and Chemical Research and National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, or private reagent sales companies such as Dainippon Pharmaceutical, R & D SYSTEMS, PharMingen, Cosmo Bio and Funakoshi Co., Ltd.

[0439] The antibody composition obtained by the process of the present invention can be administered as various therapeutic agents alone, but generally, it is preferable to provide it as a pharmaceutical formulation produced by an appropriate method well known in the technical field of pharmaceutical, by mixing it with one or more pharmaceutically acceptable carriers.

[0440] It is preferable to select a route of administration which is most effective in treatment. Examples include oral administration and parenteral administration, such as buccal, tracheal, rectal, subcutaneous, intramuscular and intravenous. In the case of an antibody preparation, intravenous administration is preferred.

[0441] The dosage form includes sprays, capsules, tablets, granules, syrups, emulsions, suppositories, injections, ointments, tapes and the like.

[0442] The pharmaceutical preparation suitable for oral administration include emulsions, syrups, capsules, tablets, powders, granules and the like.

[0443] Liquid preparations such as emulsions and syrups can be produced by using, as additives, water; sugars such as sucrose, sorbitol and fructose; glycols such as polyethylene glycol and propylene glycol; oils such as sesame oil, olive oil and soybean oil; antiseptics such as p-hydroxybenzoic acid esters; flavors such as strawberry flavor and

peppermint; and the like.

[0444] Capsules, tablets, powders, granules and the like can be prepared by using, as additives, excipients such as lactose, glucose, sucrose and mannitol; disintegrating agents such as starch and sodium alginate; lubricants such as magnesium stearate and talc; binders such as polyvinyl alcohol, hydroxypropylcellulose and gelatin; surfactants such as fatty acid ester; plasticizers such as glycerine; and the like.

[0445] The pharmaceutical preparation suitable for parenteral administration includes injections, suppositories, sprays and the like.

[0446] Injections may be prepared by using a carrier such as a salt solution, a glucose solution or a mixture thereof. Also, powdered injections can be prepared by freeze-drying the antibody composition in the usual way and adding sodium chloride thereto.

[0447] Suppositories may be prepared by using a carrier such as cacao butter, hydrogenated fat or carboxylic acid.

[0448] Also, sprays may be prepared by using the antibody composition as such or using a carrier, etc. which do not stimulate the buccal or airway mucous membrane of the patient and can facilitate absorption of the antibody composition by dispersing it as fine particles.

[0449] The carrier includes lactose, glycerine and the like. Depending on the properties of the antibody composition and the carrier, it is possible to produce pharmaceutical preparations such as aerosols and dry powders. In addition, the components exemplified as additives for oral preparations can also be added to the parenteral preparations.

[0450] Although the clinical dose or the frequency of administration varies depending on the objective therapeutic effect, administration method, treating period, age, body weight and the like, it is usually 10 µg/kg to 20 mg/kg per day and per adult.

[0451] Also, as the method for examining antitumor effect of the antibody composition against various tumor cells, *in vitro* tests include CDC activity measuring method, ADCC activity measuring method and the like, and *in vivo* tests include antitumor experiments using a tumor system in an experimental animal such as a mouse, and the like.

[0452] CDC activity and ADCC activity measurements and antitumor experiments can be carried out in accordance with the methods described in *Cancer Immunology Immunotherapy*, 36, 373 (1993); *Cancer Research*, 54, 1511 (1994) and the like.

[0453] The present invention will be described below in detail based on Examples; however, Examples are only simple illustrations, and the scope of the present invention is not limited thereto.

BRIEF DESCRIPTION OF THE DRAWINGS

[0454]

Fig. 1 shows photographs of electrophoresis patterns of SDS-PAGE of five kinds of purified anti-GD3 chimeric antibodies (using gradient gel from 4 to 15%). Fig. 1A and Fig. 1B show results of the electrophoresis under non-reducing conditions and under reducing conditions, respectively. Lanes 1 to 7 show electrophoresis patterns of high molecular weight markers, YB2/0-GD3 chimeric antibody, CHO/DG44-GD3 chimeric antibody, SP2/0-GD3 chimeric antibody, NS0-GD3 chimeric antibody (302), NS0-GD3 chimeric antibody (GIT) and low molecular weight markers, respectively.

Fig. 2 shows binding activities of five kinds of purified anti-GD3 chimeric antibodies to GD3, measured by changing the antibody concentration. The ordinate and the abscissa show the binding activity to GD3 and the antibody concentration, respectively. "o", "•", "□", "■" and "Δ" show the activities of YB2/0-GD3 chimeric antibody, CHO/DG44-GD3 chimeric antibody, SP2/0-GD3 chimeric antibody, NSO-GD3 chimeric antibody (302) and NSO-GD3 chimeric antibody (GIT), respectively.

Fig. 3 shows ADCC activities of five kinds of purified anti-GD3 chimeric antibodies to a human melanoma cell line G-361. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "o", "•", "□", "■" and "Δ" show the activities of YB2/0-GD3 chimeric antibody, CHO/DG44-GD3 chimeric antibody, SP2/0-GD3 chimeric antibody, NSO-GD3 chimeric antibody (302) and NSO-GD3 chimeric antibody (GIT), respectively.

Fig. 4 shows elution patterns of PA-treated sugar chains prepared from the anti-GD3 chimeric antibody of lot 2, obtained by analyzing them with reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively.

Fig. 5 shows binding activities of six kinds of anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond to GD3, measured by changing the antibody concentration. The ordinate and the abscissa show the binding activity to GD3 and the antibody concentration, respectively. "•", "□", "■", "Δ", "▲" and "×" show the activities of anti-GD3 chimeric antibody (50%), anti-GD3 chimeric antibody (45%), anti-GD3 chimeric antibody (29%), anti-GD3 chimeric antibody (24%), anti-GD3 chimeric antibody (13%) and anti-GD3 chimeric antibody (7%), respectively.

Fig. 6 shows results of ADCC activities using an effector cell of each doner. Fig. 6A and Fig. 6B show ADCC activities of six kinds of anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond against a human melanoma cell line G-361, using effector cells of the doner A and the doner B, respectively. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "•", "□", "■", "Δ", "▲" and "×" show the activities of anti-GD3 chimeric antibody (50%), anti-GD3 chimeric antibody (45%), anti-GD3 chimeric antibody (29%), anti-GD3 chimeric antibody (24%), anti-GD3 chimeric antibody (13%) and anti-GD3 chimeric antibody (7%), respectively.

Fig. 7 shows the relationship between sugar chain contents in which 1-fucose is not bound to *N*-acetylglucosamine at the reduced end in the doner A and the doner B and the ADCC activities.

Fig. 8 shows elution patterns of PA-treated sugar chains prepared from six kinds of anti-GD3 chimeric antibodies, obtained by analyzing them with reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively.

Fig. 9 shows binding activities of six kinds of anti-CCR4 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond measured by changing the antibody concentration to CCR4. The ordinate and the abscissa show the binding activity to CCR4 and the antibody concentration, respectively. "■", "□", "▲", "Δ", "•" and "o" show the activities of anti-CCR4 chimeric antibody (46%), anti-CCR4 chimeric antibody (39%), anti-CCR4 chimeric antibody (27%), anti-CCR4 chimeric antibody (18%), anti-CCR4 chimeric antibody (9%) and anti-CCR4 chimeric antibody (8%), respectively.

Fig. 10 shows ADCC activities of anti-CCR4 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond against CCR4/EL-4 cell, using an effector cell of the doner A. As effector cells, the effector cells derived from doner A were used. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "■", "□", "▲", "Δ", "•" and "o" show the activities of anti-CCR4 chimeric antibody (46%), anti-CCR4 chimeric antibody (39%), anti-CCR4 chimeric antibody (27%), anti-CCR4 chimeric antibody (18%), anti-CCR4 chimeric antibody (9%) and anti-CCR4 chimeric antibody (8%), respectively.

Fig. 11 shows ADCC activities of anti-CCR4 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond against CCR4/EL-4 cell, using an effector cell of the doner B. As effector cells, the effector cells derived from doner B were used. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "■", "□", "▲", "Δ", "•" and "o" show the activities of anti-CCR4 chimeric antibody (46%), anti-CCR4 chimeric antibody (39%), anti-CCR4 chimeric antibody (27%), anti-CCR4 chimeric antibody (18%), anti-CCR4 chimeric antibody (9%) and anti-CCR4 chimeric antibody (8%), respectively.

Fig. 12 shows construction of plasmids CHFT8-pCR2.1 and YBFT8-pCR2.1.

Fig. 13 shows construction of plasmids CHAc-pBS and YBAc-pBS.

Fig. 14 shows construction of plasmids CHFT8d-pCR2.1 and YBFT8d-pCR2.1.

Fig. 15 shows construction of plasmids CHAc-d-pBS and YBAc-d-pBS.

Fig. 16 shows results of determination of an α 1,6-fucosyltransferase (FUT8) transcription product in each host cell line using competitive RT-PCR. Amounts of the FUT8 transcription product in each host cell line when rat FUT8 sequence was used as the standard and internal control are shown. "■" and "□" show results when CHO cell line and YB2/0 cell line, respectively, were used as the host cell.

Fig. 17 shows results of evaluation of ADCC activities of anti-CCR4 human chimeric antibodies produced by lectin-resistant cell lines. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "□", "■", "♦" and "▲" show the activities of antibodies produced by the clone 5-03, the clone CHO/CCR4-LCA, the clone CHO/CCR4-AAL and the clone CHO/CCR4-PHA, respectively.

Fig. 18 shows results of evaluation of ADCC activities of anti-CCR4 human chimeric antibodies produced by lectin-resistant clones. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively. "□", "Δ" and "•" show activities of antibodies produced by the clone YB2/0 (KM2760#58-35-16), the clone 5-03 and the clone CHO/CCR4-LCA, respectively.

Fig. 19 shows elution patterns of PA-treated sugar chains prepared from purified anti-CCR4 human chimeric antibodies, obtained by analyzing them with reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively. Fig. 27A, Fig. 27B, Fig. 27C and Fig. 27D show results of analyses of antibodies produced by the clone 5-03, the clone CHO/CCR4-LCA, the clone CHO/CCR4-AAL and the clone CHOLCCR4-PHA, respectively.

Fig. 20 shows the 1st step of construction of an expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 21 shows the 2nd step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 22 shows the 3rd step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 23 shows the 4th step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 24 shows the 5th step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 25 shows the 6th step of construction of the expression vector of CHO cell-derived GMD (6 steps in total).

Fig. 26 shows resistance of GMD-expressed clone CHO/CCR4-LCA for LCA lectin. The measurement was carried out twice by defining the survival rate of a group of cells cultured without adding LCA lectin as 100%. In the drawing, "249" shows the survival rate of the clone CHO/CCR4-LCA introduced with an expression vector to LCA lectin. GMD shows resistance of the clone CHO/CCR4-LCA introduced with a GMD expression vector pAGE249GMD to LCA lectin.

Fig. 27 shows ADCC activities of an anti-CCR4 chimeric antibody produced by cells of GMD-expressed clone CHO/CCR4-LCA. The ordinate and the abscissa show the cytotoxic activity and the antibody concentration, respectively.

Fig. 28 shows elution patterns of PA-treated sugar chains prepared from an anti-CCR4 human chimeric antibody purified from GMD gene-expressed clone CHO/CCR4-LCA, obtained by analyzing them with reverse phase HPLC. The ordinate and the abscissa show the relative fluorescence intensity and the elution time, respectively.

Fig. 29 shows a photograph of SDS-PAGE (using 4 to 15% gradient gel) electrophoresis pattern of purified shFc γ R1IIa under reduced conditions. Lane 1 and lane M show electrophoresis patterns of shFc γ R1IIa and molecular weight markers, respectively.

Fig. 30 shows binding activities of various anti-GD3 chimeric antibodies to shFc γ R1IIa. The ordinate and abscissa show the cytotoxic activity and the antibody concentration, respectively. "o" and "*" show the activities of anti-GD3 chimeric antibody (45%) and anti-GD3 chimeric antibody (7%), respectively.

Fig. 31 shows binding activities of various anti-fibroblast growth factor-8 (FGF-8) chimeric antibodies to shFc γ R1IIa. The ordinate and abscissa show the cytotoxic activity and the antibody concentration, respectively. "o" and "*" show the activities of anti-FGF-8 chimeric antibody (58%) and anti-FGF-8 chimeric antibody (13%), respectively.

Fig. 32 shows binding activities of various anti-CCR4 chimeric antibodies to shFc γ R1IIa. In Fig. 32A, the ordinate and abscissa show the cytotoxic activity and the antibody concentration, respectively. "o", "*", "□", "■", "△", "▲" and "×" show the activities of anti-CCR4 chimeric antibody (87%), anti-CCR4 chimeric antibody (46%), anti-CCR4 chimeric antibody (39%), anti-CCR4 chimeric antibody (27%), anti-CCR4 chimeric antibody (18%), anti-CCR4 chimeric antibody (9%) and anti-CCR4 chimeric antibody (8%), respectively. In Fig. 32B, the ordinate and abscissa show the cytotoxic activity and the ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond, respectively. "*" and "o" show the activities at 40 μ g/mL and 4 μ g/mL, respectively.

Fig. 33 shows binding activities of various anti-CCR4 chimeric antibodies to shFc γ R1IIa. The ordinate and abscissa show the cytotoxic activity and the antibody concentration, respectively. "o", "△" and "*" show the activities of anti-CCR4 chimeric antibody (87%), anti-CCR4 chimeric antibody (48%) and anti-CCR4 chimeric antibody (8%), respectively.

Fig. 34 shows ADCC activities of various anti-GD3 chimeric antibodies to human myeloma cell line G-361. The ordinate and abscissa show the cytotoxic activity and the antibody concentration, respectively. "▲" and "*" show the activities of anti-GD3 chimeric antibody (42%) and anti-GD3 chimeric antibody (7%), respectively.

Fig. 35 shows results of measurement of binding activities of FGF-8/Fc fusion proteins to KM1334. The ordinate and abscissa show the binding activity and the antibody concentration, respectively. "■" and "o" show the activities of FGF-8/Fc fusion proteins produced by the cell line YB2/0 and the cell line CHO/DG44, respectively, as the host cell.

Fig. 36 shows results of measurement of binding activities of various FGF-8/Fc fusion proteins to shFc γ R1IIa(V). The ordinate and abscissa show the binding activity and the antibody concentration, respectively. "■" and "o" show the activities of FGF8/Fc fusion proteins produced by the cell line YB2/0 and the cell line CHO/DG44, respectively, as the host cell.

Fig. 37 show a construction step of a plasmid CHO-GMD in which the 5'-terminal of a clone 34-2 is introduced into the 5'-terminal of a CHO cell-derived GMD cDNA clone 22-8.

Fig. 38 shows construction steps of plasmid pKANTEX1334H and pKANTEX 1334.

Best Mode for Carrying Out the Invention

Example 1

5 Preparation of anti-ganglioside GD3 human chimeric antibody:

1. Construction of tandem expression vector pChi641LHGM4 for anti-ganglioside GD3 human chimeric antibody

10 **[0455]** A plasmid pChi641LGM40 was constructed by ligating a fragment of about 4.03 kb containing an L chain cDNA, obtained by digesting an L chain expression vector pChi641LGM4 [*J. Immunol. Methods*, 167, 271 (1994)] for anti-ganglioside GD3 human chimeric antibody (hereinafter referred to as "anti-GD3 chimeric antibody") with restriction enzymes *MluI* (manufactured by Takara Shuzo) and *SalI* (manufactured by Takara Shuzo) with a fragment of about 3.40 kb containing a G418-resistant gene and a splicing signal, obtained by digesting an expression vector pAGE107 [*Cytotechnology*, 3, 133 (1990)] for animal cell with restriction enzymes *MluI* (manufactured by Takara Shuzo) and *SalI* (manufactured by Takara Shuzo) using DNA Ligation Kit (manufactured by Takara Shuzo), and then transforming *E. coli* HB101 (*Molecular Cloning*, Second Edition) with the ligated product.

15 **[0456]** Next, a fragment of about 5.68 kb containing an L chain cDNA, obtained by digesting the constructed plasmid pChi641LGM40 with a restriction enzyme *ClaI* (manufactured by Takara Shuzo), changing it to blunt-end using DNA Blunting Kit (manufactured by Takara Shuzo) and further digesting it with *MluI* (manufactured by Takara Shuzo), was ligated with a fragment of about 8.40 kb containing an H chain cDNA, obtained by digesting an anti-GD3 chimeric antibody H chain expression vector pChi641HGM4 [*J. Immunol. Methods*, 167, 271 (1994)] with a restriction enzyme *XhoI* (manufactured by Takara Shuzo), changing it to blunt-end using DNA Blunting Kit (manufactured by Takara Shuzo) and further digesting it with *MluI* (manufactured by Takara Shuzo) using DNA Ligation Kit (manufactured by Takara Shuzo), and then *E. coli* HB101 (*Molecular Cloning*, Second Edition) was transformed with the ligated product to thereby
25 construct a tandem expression vector pChi641LHGM4 for anti-GD3 chimeric antibody.

2. Preparation of cell stably producing anti-GD3 chimeric antibody

30 **[0457]** Cells capable of stably producing an anti-GD3 chimeric antibody were prepared by introducing the tandem expression vector pChi641LHGM4 for anti-GD3 chimeric antibody constructed in the item 1 of Example 1 and selecting suitable clones, as described below.

(1) Preparation of producing cell using rat myeloma YB2/0 cell

35 **[0458]** After introducing 5 µg of the anti-GD3 chimeric antibody expression vector pChi641LHGM4 into 4×10^6 cells of rat myeloma YB2/0 [ATCC CRL-1662, *J. Cell. Biol.*, 93, 576 (1982)] by electroporation [*Cytotechnology*, 3, 133 (1990)], the cells were suspended in 40 ml of RPMI1640-FBS(10) (RPMI1640 medium comprising 10% (fetal bovine serum (hereinafter referred to as "FBS") (manufactured by GIBCO BRL)) and dispensed at 200 µl/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418
40 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of transformants showing G418 resistance were formed and growth of colonies was observed, and the antigen binding activity of the anti-GD3 chimeric antibody in the supernatant was measured by the ELISA shown in the item 3 of Example 1.

45 **[0459]** Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, in order to increase the amount of the antibody production using a DHFR gene amplification system, each of them was suspended in the RPMI1640-FBS(10) medium comprising 0.5 mg/ml G418 and 50 nmol/L DHFR inhibitor, methotrexate (hereinafter referred to as "MTX"; manufactured by SIGMA) to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed at 2 ml into each well of a 24 well plate (manufactured by Greiner). Transformants showing 50 nmol/L MTX resistance were induced by culturing at 37°C for 1 to 2 weeks in a 5% CO₂
50 incubator. The antigen binding activity of the anti-GD3 chimeric antibody in culture supernatants in wells in which growth of transformants was observed was measured by the ELISA shown in the item 3 of Example 1. Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, the MTX concentration was increased to 100 nmol/L and then to 200 nmol/L, and transformants capable of growing in the RPMI1640-FBS(10) medium comprising 0.5 mg/ml G418 and 200 nmol/L MTX and capable of producing the anti-GD3
55 chimeric antibody in a large amount were finally obtained by the same method as described above. Among the obtained transformants, suitable clones were selected and were made into a single cell (cloning) by limiting dilution twice.

[0460] The obtained anti-GD3 chimeric antibody-producing transformed cell clone 7-9-51 has been deposited on April 5, 1999, as FERM BP-6691 in National Institute of Bioscience and Human Technology, Agency of Industrial Sci-

ence and Technology (Higashi 1-1-3, Tsukuba, Ibaraki, Japan) (present name: International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1, Higashi 1-Chome Tsukuba-shi, Ibaraki-ken, Japan))

(2) Preparation of producing cell using CHO/DG44 cell

[0461] After introducing 4 µg of the anti-GD3 chimeric antibody expression vector pChi641LHGM4 into 1.6×10^6 cells of CHO/DG44 cell [*Proc. Natl. Acad. Sci. USA*, 77, 4216 (1980)] by electroporation [*Cytotechnology*, 3, 133 (1990)], the cells were suspended in 10 ml of IMDM-FBS(10)-HT(1) [IMDM medium comprising 10% FBS and 1 x concentration of HT supplement (manufactured by GIBCO BRL)] and dispensed at 200 µl/well into a 96 well culture plate (manufactured by Iwaki Glass). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of transformants showing G418 resistance were formed and growth of colonies was observed, and the antigen binding activity of the anti-GD3 chimeric antibody in the supernatant was measured by the ELISA shown in the item 3 of Example 1.

[0462] Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, in order to increase the amount of the antibody production using a DHFR gene amplification system, each of them was suspended in an IMDM-dFBS(10) medium [IMDM medium comprising 10% dialyzed fetal bovine serum (hereinafter referred to as "dFBS"; manufactured by GIBCO BRL)] comprising 0.5 mg/ml G418 and 10 nmol/L MTX to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed at 0.5 ml into each well of a 24 well plate (manufactured by Iwaki Glass). Transformants showing 10 nmol/L MTX resistance were induced by culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator. Regarding the transformants in wells in which their growth was observed, the MTX concentration was increased to 100 nmol/L, and transformants capable of growing in the IMDM-dFBS(10) medium comprising 0.5 mg/ml G418 and 100 nmol/L MTX and of producing the anti-GD3 chimeric antibody in a large amount were finally obtained by the same method as described above. Among the obtained transformants, suitable clones were selected and were made into a single cell (cloning) by limiting dilution twice.

(3) Preparation of producing cell using mouse myeloma NS0 cell

[0463] After introducing 5 µg of the anti-GD3 chimeric antibody expression vector pChi641LHGM4 into 4×10^6 cells of mouse myeloma NS0 by electroporation [*Cytotechnology*, 3, 133 (1990)], the cells were suspended in 40 ml of EX-CELL302-FBS(10) [EX-CELL302 medium comprising 10% FBS and 2 mmol/L L-glutamine (hereinafter referred to as "L-Gln"; manufactured by GIBCO BRL)] and dispensed at 200 µl/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of transformants showing G418 resistance were formed and growth of colonies was observed, and the antigen binding activity of the anti-GD3 chimeric antibody in the supernatant was measured by the ELISA shown in the item 3 of Example 1.

[0464] Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, in order to increase the amount of the antibody production using a DHFR gene amplification system, each of them was suspended in an EX-CELL302-dFBS(10) medium (EX-CELL302 medium comprising 10% dFBS and 2 mmol/L L-Gln) comprising 0.5 mg/ml G418 and 50 nmol/L MTX to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed at 2 ml into each well of a 24 well plate (manufactured by Greiner). Transformants showing 50 nmol/L MTX resistance were induced by culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator. The antigen binding activity of the anti-GD3 chimeric antibody in culture supernatants in wells in which growth of transformants was observed was measured by the ELISA shown in the item 3 of Example 1. Regarding the transformants in wells in which production of the anti-GD3 chimeric antibody was observed in culture supernatants, the MTX concentration was increased to 100 nmol/L and then to 200 nmol/L, and transformants capable of growing in the EX-CELL302-dFBS(10) medium comprising 0.5 mg/ml G418 and 200 nmol/L MTX and of producing the anti-GD3 chimeric antibody in a large amount was finally obtained by the same method as described above. Among the obtained transformants, suitable clones were selected and were made into a single cell (cloning) by limiting dilution twice. Also, using the method for determining the transcription product of an α1,6-fucosyltransferase gene shown in Example 9, a clone producing a relatively small amount of the transcription product was selected and used as a suitable clone.

3. Measurement of binding activity of antibody to GD3 (ELISA)

[0465] The binding activity of the antibody to GD3 was measured as described below.

[0466] In 2 ml of an ethanol solution containing 10 µg of dipalmitoylphosphatidylcholine (manufactured by SIGMA)

and 5 µg of cholesterol (manufactured by SIGMA), 4 nmol of GD3 (manufactured by Snow Brand Milk Products) was dissolved. Into each well of a 96 well plate for ELISA (manufactured by Greiner), 20 µl of the solution (40 pmol/well in final concentration) was dispensed, followed by air-drying, 1% bovine serum albumin (hereinafter referred to as "BSA"; manufactured by SIGMA)-containing PBS (hereinafter referred to as "1% BSA-PBS") was dispensed at 100 µl/well, and then the reaction was carried out at room temperature for 1 hour to block remaining active groups. After discarding 1% BSA-PBS, a culture supernatant of a transformant or a diluted solution of a human chimeric antibody was dispensed at 50 µl/well to carry out the reaction at room temperature for 1 hour. After the reaction, each well was washed with 0.05% Tween 20 (manufactured by Wako Pure Chemical Industries)-containing PBS (hereinafter referred to as "Tween-PBS"), a peroxidase-labeled goat anti-human IgG (H & L) antibody solution (manufactured by American Qualex) diluted 3,000 times with 1% BSA-PBS was dispensed at 50 µl/well as a secondary antibody solution, and then the reaction was carried out at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, ABTS substrate solution [solution prepared by dissolving 0.55 g of 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) ammonium salt in 1 liter of 0.1 mol/L citrate buffer (pH 4.2) and adding 1 µl/ml of hydrogen peroxide to the solution just before use (hereinafter the same solution was used)] was dispensed at 50 µl/well for color development, and then absorbance at 415 nm (hereinafter referred to as "OD415") was measured.

4. Purification of anti-GD3 chimeric antibody

(1) Culturing of producing cell derived from YB2/0 cell and purification of antibody

[0467] The anti-GD3 chimeric antibody-producing transformed cell clone obtained in the item 2(1) of Example 1 was suspended in the Hybridoma-SFM medium comprising 0.2% BSA, 200 nmol/L MTX and 100 nmol/L triiodothyronine (hereinafter referred to as "T3"; manufactured by SIGMA) to give a density of 3×10^5 cells/ml and cultured in a 2.0 liter bottle (manufactured by Iwaki Glass) under stirring at a rate of 50 rpm. After culturing at 37°C for 10 days in a temperature-controlling room, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named YB2/0-GD3 chimeric antibody.

(2) Culturing of producing cell derived from CHO/DG44 cell and purification of antibody

[0468] The anti-GD3 chimeric antibody-producing transformed cell clone obtained in the item 2(2) of Example 1 was suspended in the EX-CELL302 medium comprising 3 mmol/L L-Gln, 0.5% fatty acid concentrated solution (hereinafter referred to as "CDLC"; manufactured by GIBCO BRL) and 0.3% Pluronic F68 (hereinafter referred to as "PF68"; manufactured by GIBCO BRL) to give a density of 1×10^6 cells/ml, and the suspension was dispensed at 50 ml into 175 mm² flasks (manufactured by Greiner). After culturing at 37°C for 4 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named CHO/DG44-GD3 chimeric antibody.

(3) Culturing of producing cell derived from NS0 cell and purification of antibody

[0469] The anti-GD3 chimeric antibody-producing transformed cell clone obtained in the item 2(3) of Example 1 was suspended in the EX-CELL302 medium comprising 2 mmol/L L-Gln, 0.5 mg/ml G418, 200 nmol/L MTX and 1% FBS, to give a density of 1×10^6 cells/ml, and the suspension was dispensed at 200 ml into 175 mm² flasks (manufactured by Greiner). After culturing at 37°C for 4 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named NS0-GD3 chimeric antibody (302).

[0470] Also, the transformed cell clone was suspended in the GIT medium comprising 0.5 mg/ml G418 and 200 nmol/L MTX to give a density of 3×10^5 cells/ml, and the suspension was dispensed at 200 ml into 175 mm² flasks (manufactured by Greiner). After culturing at 37°C for 10 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named NS0-GD3 chimeric antibody (GIT).

(4) Culturing of producing cell derived from SP2/0 cell and purification of antibody

[0471] The anti-GD3 chimeric antibody-producing transformed cell clone (KM-871 (FERM BP-3512)) described in

Japanese Published Unexamined Patent Application No. 304989/93 (EP 533199) was suspended in the GIT medium comprising 0.5 mg/ml G418 and 200 nmol/L MTX to give a density of 3×10^5 cells/ml, and the suspension was dispensed at 200 ml into 175 mm² flasks (manufactured by Greiner). After culturing at 37°C for 8 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-GD3 chimeric antibody was purified from the culture supernatant using a Prosep-A (manufactured by Bioprocessing) column in accordance with the manufacture's instructions. The purified anti-GD3 chimeric antibody was named SP2/0-GD3 chimeric antibody.

5. Analysis of purified anti-GD3 chimeric antibody

[0472] In accordance with a known method [*Nature*, 227, 680 (1970)], 4 µg of each of the five kinds of the anti-GD3 chimeric antibodies produced by and purified from respective animal cells, obtained in the item 4 of Example 1, was subjected to SDS-PAGE to analyze the molecular weight and purity. The results are shown in Fig. 1. As shown in Fig. 1, a single band of about 150 kilodaltons (hereinafter referred to as "Kd") in molecular weight was found under non-reducing conditions, and two bands of about 50 Kd and about 25 Kd under reducing conditions, in each of the purified anti-GD3 chimeric antibodies. The molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of H chain and L chain of the antibody (H chain: about 49 Kd, L chain: about 23 Kd, whole molecule: about 144 Kd), and also coincided with the reports stating that the IgG antibody has a molecular weight of about 150 Kd under non-reducing conditions and is degraded into H chains having a molecular weight of about 50 Kd and L chains having a molecular weight of about 25 Kd under reducing conditions due to cutting of the disulfide bond (hereinafter referred to as "S-S bond") in the molecule (*Antibodies*, Chapter 14; *Monoclonal Antibodies*), so that it was confirmed that each anti-GD3 chimeric antibody was expressed and purified as an antibody molecule having the true structure.

Example 2

Activity evaluation of anti-GD3 chimeric antibody:

1. Binding activity of anti-GD3 chimeric antibody to GD3 (ELISA)

[0473] Binding activities of the five kinds of the purified anti-GD3 chimeric antibodies obtained in the item 4 of Example 1 to GD3 were measured by the ELISA shown in the item 3 of Example 1. Fig. 2 shows results of the examination of the binding activity measured by changing the concentration of the anti-GD3 chimeric antibody to be added. As shown in Fig. 2, the five kinds of the anti-GD3 chimeric antibodies showed almost the same binding activity to GD3. The result shows that antigen binding activities of these antibodies are constant independently of the antibody-producing animal cells and their culturing methods. Also, it was suggested from the comparison of the NSO-GD3 chimeric antibody (302) with the NSO-GD3 chimeric antibody (GIT) that the antigen binding activities are constant independently of the media used in the culturing.

2. ADCC activity of anti-GD3 chimeric antibody

[0474] ADCC activities of the five kinds of the purified anti-GD3 chimeric antibodies obtained in the item 4 of Example 1 were measured in accordance with the following method.

(1) Preparation of target cell solution

[0475] A human melanoma cell line G-361 (ATCC CRL 1424) was cultured in the RPMI1640-FBS(10) medium to prepare 1×10^6 cells, and the cells were radioisotope-labeled by reacting them with 3.7 MBq equivalents of a radioactive substance Na₂⁵¹CrO₄ at 37°C for 1 hour. After the reaction, the cells were washed three times through their suspension in the RPMI1640-FBS(10) medium and centrifugation, resuspended in the medium and then allowed to react at 4°C for 30 minutes on ice for spontaneous dissolution of the radioactive substance. After centrifugation, the precipitate was adjusted to 2×10^5 cells/ml by adding 5 ml of the RPMI1640-FBS(10) medium and used as the target cell solution.

(2) Preparation of effector cell solution

[0476] From a healthy donor, 50 ml of venous blood was collected, and gently mixed with 0.5 ml of heparin sodium (manufactured by Takeda Pharmaceutical). The mixture was centrifuged to isolate a mononuclear cell layer using Lymphoprep (manufactured by Nycomed Pharma AS) in accordance with the manufacture's instructions. After washing with the RPMI1640-FBS(10) medium by centrifugation three times, the resulting precipitate was re-suspended to give

a density of 2×10^6 cells/ml by using the medium and used as the effector cell solution.

(3) Measurement of ADCC activity

[0477] Into each well of a 96 well U-shaped bottom plate (manufactured by Falcon), 50 μ l of the target cell solution prepared in the above (1) (1×10^4 cells/well) was dispensed. Next, 100 μ l of the effector cell solution prepared in the above (2) was added thereto (2×10^5 cells/well, the ratio of effector cells to target cells becomes 20:1). Subsequently, each of the anti-GD3 chimeric antibodies was added at various concentrations, followed by reaction at 37°C for 4 hours. After the reaction, the plate was centrifuged, and the amount of ^{51}Cr in the supernatant was measured with a γ -counter. The amount of spontaneously released ^{51}Cr was calculated by the same operation using only the medium instead of the effector cell solution and the antibody solution, and measuring the amount of ^{51}Cr in the supernatant. The amount of total released ^{51}Cr was calculated by the same operation as above using only the medium instead of the antibody solution and adding 1 N hydrochloric acid instead of the effector cell solution, and measuring the amount of ^{51}Cr in the supernatant. The ADCC activity was calculated from the following equation (1):

$$\text{ADCC activity (\%)} = \frac{{}^{51}\text{Cr in sample supernatant} - \text{spontaneously released } {}^{51}\text{Cr}}{\text{total released } {}^{51}\text{Cr} - \text{spontaneously released } {}^{51}\text{Cr}} \times 100 \quad (1)$$

[0478] The results are shown in Fig. 3. As shown in Fig. 3, among the five kinds of the anti-GD3 chimeric antibodies, the YB2/0-GD3 chimeric antibody showed the highest ADCC activity, followed by the SP2/0-GD3 chimeric antibody, NSO-GD3 chimeric antibody and CHO-GD3 chimeric antibody in that order. No difference in the ADCC activity was found between the NSO-GD3 chimeric antibody (302) and NSO-GD3 chimeric antibody (GIT) prepared by using different media in the culturing. The above results show that the ADCC activity of antibodies greatly varies depending on the kind of the animal cells to be used in their production. As its mechanism, since their antigen binding activities were equal, it was considered that ADCC activity depends on a difference in the structure of the Fc region of the antibody.

Example 3

[0479] Activity evaluation of anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end:

1. Preparation of anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond

[0480] In accordance with the method described in the item 2(1) of Example 1, some transformed clones derived from YB2/0 cell capable of producing an anti-GD3 chimeric antibody was obtained. Antibodies were prepared from the transformed clones derived from YB2/0 cell and named lot 1, lot 2 and lot 3. Sugar chain analysis of the anti-GD3 chimeric antibodies of lot 1, lot 2 and lot 3 was carried out by the following method.

[0481] The solution of each purified antibody was exchanged to 10 mmol/L KH_2PO_4 using Ultra Free 0.5-10K (manufactured by Millipore). The exchange was carried out in such a manner that the exchanging ratio became 80-fold or more.



[0482] Into Hydraclub S-204 test tube, 100 μ g of each antibody was put and dried with a centrifugal evaporator. The dried sample was subjected to hydrazinolysis using Hydraclub manufactured by Hohnen. The sample was allowed to react with hydrazine at 110°C for 1 hour using a hydrazinolysis reagent manufactured by Hohnen [*Method of Enzymology*, 83, 263 (1982)]. After the reaction, hydrazine was evaporated under a reduced pressure, and the reaction tube was returned to room temperature by allowing it to stand for 30 minutes. Next, 250 μ l of an acetylation reagent manufactured by Hohnen and 25 μ l of acetic anhydride were added thereto, followed by thoroughly stirred for reaction at room temperature for 30 minutes. Then, 250 μ l of the acetylation reagent and 25 μ l of acetic anhydride were further added thereto, followed by thoroughly stirring for reaction at room temperature for 1 hour. The sample was frozen at -80°C in a freezer and freeze-dried for about 17 hours. Sugar chains were recovered from the freeze-dried sample by using Cellulose Cartridge Glycan Preparation Kit manufactured by Takara Shuzo. The sample sugar chain solution was dried with a centrifugal evaporator and then subjected to fluorescence labeling with 2-aminopyridine [*J. Biochem.*, 95, 197 (1984)]. The 2-aminopyridine solution was prepared by adding 760 μ l of HCl per 1 g of 2-aminopyridine (1 \times PA solution) and diluting the solution 10-fold with reverse osmosis purified water (10-fold diluted PA solution). The sodium cyanoborohydride solution was prepared by adding 20 μ l of 1 \times PA solution and 430 μ l of reverse osmosis purified water per 10 mg of sodium cyanoborohydride. To the sample, 67 μ l of a 10 fold-diluted PA solution was added, followed by reaction at 100°C for 15 minutes and spontaneously cooled, and 2 μ l of sodium cyanoborohydride was

[0483] Next, reverse phase HPLC analysis of the purified PA-treated sugar chains was carried out by using CLC-ODS column (manufactured by Shimadzu, ϕ 6.0 nm \times 159 nm). The step was carried out at a column temperature of 55°C and at a flow rate of 1 ml/min and using a fluorescence detector of 320 nm excitation wavelength and 400 nm fluorescence wavelength. The column was equilibrated with a 10 mmol/L sodium phosphate buffer (pH 3.8) and elution was carried out for 80 minutes by a 0.5% 1-butanol linear density gradient. Fig. 4 shows elution patterns of the purified PA-treated sugar chains of the anti-GD3 antibody of lot 2. Each of the PA-treated sugar chain was identified by post source decay analysis of each peak of the separated PA-treated sugar chains using matrix-assisted laser ionization type of flight mass spectrometry (MALDI-TOF-MS analysis), comparison of elution positions with standards of PA-treated sugar chain manufactured by Takara Shuzo, and reverse phase HPLC analysis after digestion of each PA-treated sugar chain using various enzymes.

[0484] The sugar chain content was calculated from each of the peak area of PA-treated sugar chain by reverse HPLC analysis. A PA-treated sugar chain whose reducing end is not *N*-acetylglucosamine was excluded from the peak area calculation, because it is an impurity or a by-product during preparation of PA-treated sugar chain. Peaks (i) to (ix) in the figure show the following structures (1) to (9), respectively.



[illegible]



[0485] GlcNAc, Gal, Man, Fuc and PA indicate *N*-acetylglucosamine, galactose, mannose, fucose and a pyridylamino group, respectively. In Fig. 4, the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was calculated from the area occupied by the peaks (i) to (iv) among (i) to (ix), and the ratio of a sugar chain in which 1-position of fucose was bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was calculated from the area occupied by the peaks (v) to (ix) among (i) to (ix). Each ratio of a sugar chain was shown as an average value of the result of two sugar chain analyses.

[0486] As a result, the ratios of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond were 50%, 45% and 29% in lot 1, lot 2 and lot 3, respectively. Herein, these samples were named anti-GD3 chimeric antibody (50%), anti-GD3 chimeric antibody (45%) and anti-GD3 chimeric antibody (29%).

[0487] Also, sugar chains of the anti-GD3 chimeric antibody derived from the CHO/DG44 cell prepared in the item 2(2) of Example 1 were analyzed in accordance with the above-described method, and it was found that the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was 7%. Herein, the sample was named anti-GD3 chimeric antibody (7%).

[0488] Further, the anti-GD3 chimeric antibody (45%) and anti-GD3 chimeric antibody (7%) were mixed at a ratio of anti-GD3 chimeric antibody (45%) : anti-GD3 chimeric antibody (7%) = 5:3 and 1:7, respectively. Sugar chains of the samples were analyzed in accordance with the above-described method, and the ratios of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond were 24% and 13%. Herein, they were named anti-GD3 chimeric antibody (24%) and anti-GD3 chimeric antibody (13%).

2. Evaluation of binding activity to GD3 (ELISA)

[0489] The binding activities of the six kinds of the anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond prepared in the item 1 of Example 3 to GD3 were measured by the ELISA shown in the item 3 of Example 1. As a result, all of the six kinds of the anti-GD3 chimeric antibodies showed almost the same GD3-binding activity as shown in Fig. 5, and it was found that the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond does not have influence on the antigen binding activity of the antibody.

3. Evaluation of ADCC activity on human melanoma cell line

[0490] ADCC activities of six kinds of the anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond prepared in the item 1 of Example 3 to human melanoma cell line G-361 (ATCC CRL1424) were measured according to the method described in the item 2 of Example 2.

[0491] Figs. 6 and 7 show results of the measurement of ADCC activity of the six kinds of the anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond at various concentrations (0.0005 to 5 μ g/ml) using effector cells of two healthy donors (A and B). As shown in Figs. 6 and 7, the ADCC activity of the anti-GD3 chimeric antibodies showed a tendency to increase in proportion to the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond at each antibody concentration. The ADCC activity decreases when the antibody concentration is low.

[0492] At an antibody concentration of 0.05 μ g/ml, the antibody in which the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was 24%, 29%, 45% or 50% showed almost the same high ADCC activity, but the antibody (13%) or (7%) in which the ratio of a sugar chain in which 1-position of fucose was bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is less than 20%, 13% or 7% showed the low ADCC activity. These results did not change even if the effector cell having doner was different.

Example 4

[0493] Activity evaluation of anti-CCR4 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond:

1. Preparation of cells stably producing anti-CCR4 chimeric antibody

[0494] Cells which are capable of stably producing an anti-CCR4 chimeric antibody were prepared as follows by using a tandem type expression vector pKANTEX2160 for an anti-CCR4 chimeric antibody described in WO01/64754.

(1) Preparation of producing cell using rat myeloma YB2/0 cell

[0495] After introducing 10 μ g of the anti-CCR4 chimeric antibody expression vector pKANTEX2160 into 4×10^6 cells of rat myeloma YB2/0 cell (ATCC CRL 1662) by electroporation [*Cytotechnology*, 3, 133 (1990)], the cells were suspended in 40 ml of Hybridoma-SFM-FBS(5) [Hybridoma-SFM medium (manufactured by Invitrogen) comprising 5% FBS (manufactured by PAA Laboratories)] and dispensed at 200 μ l/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 1 mg/ml, followed by culturing for 1 to 2 weeks. Culture supernatant was recovered from wells in which growth of transformants showing G418 resistance was observed by the formation of colonies, and antigen binding activity of the anti-CCR4 chimeric antibody in the supernatant was measured by the ELISA described in the item 2 of Example 4.

[0496] Regarding the transformants in wells in which production of the anti-CCR4 chimeric antibody was observed in culture supernatants, in order to increase an amount of the antibody production using a DHFR gene amplification system, each of them was suspended in the Hybridoma-SFM-FBS(5) medium comprising 1 mg/ml G418 and 50 nmol/L DHFR inhibitor MTX (manufactured by SIGMA) to give a density of 1 to 2×10^5 cells/ml, and the suspension was dispensed at 1 ml into each well of a 24 well plate (manufactured by Greiner). After culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nmol/L MTX resistance were induced. Antigen binding activity of the anti-CCR4 chimeric antibody in culture supernatants in wells in which growth of transformants was observed was measured by the ELISA described in the item 2 of Example 4.

[0497] Regarding the transformants in wells in which production of the anti-CCR4 chimeric antibody was observed in culture supernatants, the MTX concentration was increased by the same method, and a transformant capable of growing in the Hybridoma-SFM-FBS(5) medium comprising 200 nmol/L MTX and of producing the anti-CCR4 chimeric antibody in a large amount was finally obtained. The obtained transformant was made into a single cell (cloning) by limiting dilution twice, and the obtained clone was named KM2760#58-35-16.

(2) Preparation of producing cell using CHO/DG44 cell

[0498] After introducing 4 μ g of the anti-CCR4 chimeric antibody expression vector pKANTEX2160 into 1.6×10^6 cells of CHO/DG44 cell by electroporation [*Cytotechnology*, 3, 133 (1990)], the cells were suspended in 10 ml of IMDM-

dFBS(10)-HT(1) (IMDM medium (manufactured by Invitrogen) comprising 10% dFBS (manufactured by Invitrogen) and 1× concentration of HT supplement (manufactured by Invitrogen)] and dispensed at 100 µl/well into a 96 well culture plate (manufactured by Iwaki Glass). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, the medium was changed to IMDM-dFBS(10) (IMDM medium comprising 10% of dialyzed FBS), followed by culturing for 1 to 2 weeks. Culture supernatant was recovered from wells in which the growth was observed due to formation of a transformant showing HT-independent growth, and an expression amount of the anti-CCR4 chimeric antibody in the supernatant was measured by the ELISA described in the item 2 of Example 4.

[0499] Regarding the transformants in wells in which production of the anti-CCR4 chimeric antibody was observed in culture supernatants, in order to increase an amount of the antibody production using a DHFR gene amplification system, each of them was suspended in the IMDM-dFBS(10) medium comprising 50 nmol/L MTX to give a density of 1 to 2×10⁵ cells/ml, and the suspension was dispensed at 0.5 ml into each well of a 24 well plate (manufactured by Iwaki Glass). After culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nmol/L MTX resistance were induced. Regarding the transformants in wells in which the growth was observed, the MTX concentration was increased to 200 nmol/L by the similar method as above, and a transformant capable of growing in the IMDM-dFBS(10) medium comprising 200 nmol/L MTX and of producing the anti-CCR4 chimeric antibody in a large amount was finally obtained. The obtained transformant was named clone 5-03.

2. Binding activity of antibody to CCR4 partial peptide (ELISA)

[0500] Compound 1 (SEQ ID NO:25) was selected as a human CCR4 extracellular region peptide capable of reacting with the anti-CCR4 chimeric antibody. In order to use it in the activity measurement by ELISA, a conjugate with BSA (manufactured by Nacalai Tesque) was prepared by the following method and used as the antigen. That is, 100 ml of a DMSO solution comprising 25 mg/ml SMCC [4-(N-maleimidomethyl)-cyclohexane-1-carboxylic acid *N*-hydroxysuccinimide ester] (manufactured by Sigma) was added dropwise to 900 ml of a 10 mg BSA-containing PBS solution under stirring with a vortex, followed by gently stirring for 30 minutes. To a gel filtration column such as NAP-10 column equilibrated with 25 ml of PBS, 1 ml of the reaction solution was applied and then eluted with 1.5 ml of PBS and the resulting eluate was used as a BSA-SMCC solution (BSA concentration was calculated based on A₂₈₀ measurement). Next, 250 ml of PBS was added to 0.5 mg of Compound 1 and then completely dissolved by adding 250 ml of DMF, and the BSA-SMCC solution was added thereto under vortex, followed by gently stirring for 3 hours. The reaction solution was dialyzed against PBS at 4°C overnight, sodium azide was added thereto to give a final concentration of 0.05%, and the mixture was filtered through a 0.22 µm filter to be used as a BSA-compound 1 solution.

[0501] The prepared conjugate was dispensed at 0.05 µg/ml and 50 µl/well into a 96 well EIA plate (manufactured by Greiner) and incubated for adhesion at 4°C overnight. After washing each well with PBS, 1% BSA-PBS was added thereto in 100 µl/well and allowed to react at room temperature to block the remaining active groups. After washing each well with Tween-PBS, a culture supernatant of a transformant was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction, each well was washed with Tween-PBS, and then a peroxidase-labeled goat anti-human IgG(γ) antibody solution (manufactured by American Qualex) diluted 6000 times with 1% BSA-PBS as the secondary antibody solution was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was added at 50 µl/well for color development, and 20 minutes thereafter, the reaction was stopped by adding a 5% SDS solution at 50 µl/well. Thereafter, the absorbance at OD₄₁₅ was measured. The anti-CCR4 chimeric antibody obtained in the item 1 of Example 4 showed the binding activity to CCR4.

3. Purification of anti-CCR4 chimeric antibody

(1) Culturing of producing cell derived from YB2/O cell and purification of antibody

[0502] The anti-CCR4 chimeric antibody-expressing transformant cell clone KM2760#58-35-16 obtained in the item 1(1) of Example 4 was suspended in Hybridoma-SFM (manufactured by Invitrogen) medium comprising 200 nmol/L MTX and 5% of Daigo's GF21 (manufactured by Wako Pure Chemical Industries) to give a density of 2×10⁵ cells/ml and subjected to fed-batch shaking culturing with a spinner bottle (manufactured by Iwaki Glass) in a constant temperature chamber of 37°C. After culturing for 8 to 10 days and recovering the culture supernatant, the anti-CCR4 chimeric antibody was purified using Prosep-A (manufactured by Millipore) column and gel filtration. The purified anti-CCR4 chimeric antibody was named KM2760-1.

(2) Culturing of producing cell derived from CHO-DG44 cell and purification of antibody

[0503] The anti-CCR4 chimeric antibody-producing transformant clone 5-03 obtained in the item 1(2) of Example 4

was cultured at 37°C in a 5% CO₂ incubator using IMDM-dFBS(10) medium in a 182 cm² flask (manufactured by Greiner). When the cell density reached confluent after several days, the culture supernatant was discarded, and the cells were washed with 25 ml of PBS buffer and then mixed with 35 ml of EXCELL 301 medium (manufactured by JRH). After culturing at 37°C for 7 days in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-CCR4 chimeric antibody was purified from the culture supernatant using Prosep-A (manufactured by Millipore) column in accordance with the manufacture's instructions. The purified anti-CCR4 chimeric antibody was named KM3060.

[0504] When the binding activity to CCR4 of KM2760-1 and KM3060 was measured by the ELISA described in the item 2 of Example 4, they showed equivalent binding activity.

4. Analysis of purified anti-CCR4 chimeric antibodies

[0505] Each 4 µg of the two kinds of the anti-CCR4 chimeric antibodies produced by and purified from various animal cells, obtained in the item 3 of Example 4 was subjected to SDS-PAGE in accordance with a known method [*Nature*, 227, 680 (1970)], and the molecular weight and purity were analyzed. In each of the purified anti-CCR4 chimeric antibodies, a single band corresponding to the molecular weight of about 150 Kd was found under non-reducing conditions, and two bands of about 50 Kd and about 25 Kd were found under reducing conditions. The molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of antibody H chain and L chain (H chain: about 49 Kd, L chain: about 23 Kd, whole molecule: about 144 Kd) and further coincided with reports stating that an IgG type antibody has a molecular weight of about 150 Kd under non-reducing conditions and is degraded into H chain having a molecular weight of about 50 Kd and L chain having a molecular weight of about 25 Kd under reducing conditions caused by cutting an S-S bond in the molecule (*Antibodies*, Chapter 14 (1988), *Monoclonal Antibodies*), thus confirming that the anti-CCR4 chimeric antibody was expressed and purified as an antibody molecule having a correct structure.

5. Preparation of anti-CCR4 chimeric antibody having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond

[0506] Sugar chains of the anti-CCR4 chimeric antibody KM2760-1 derived from YB2/0 cell and the anti-CCR4 chimeric antibody KM3060 derived from CHO/DG44 cell prepared in the item 3 of Example 4 were analyzed in accordance with the method in the item 1 of Example 3. The ratios of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond were 87% and 8% in KM2760-1 and KM3060, respectively. Herein, the samples were named anti-CCR4 chimeric antibody (87%) and anti-CCR4 chimeric antibody (8%).

[0507] Furthermore, the anti-CCR4 chimeric antibody (87%) and anti-CCR4 chimeric antibody (8%) were mixed at a ratio of anti-CCR4 chimeric antibody (87%) : anti-CCR4 chimeric antibody (8%) = 1 : 39, 16 : 67, 22 : 57, 32 : 47 and 42 : 37, respectively. Sugar chains of these samples were analyzed in accordance with the method of the item 1 of Example 3. The ratios of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond were 9%, 18%, 27%, 39% and 46%, respectively. Herein, these samples were named anti-CCR4 chimeric antibody (9%), anti-CCR4 chimeric antibody (18%), anti-CCR4 chimeric antibody (27%), anti-CCR4 chimeric antibody (39%) and anti-CCR4 chimeric antibody (46%).

[0508] Results of the sugar chain analysis of each of the samples are shown in Fig. 8. The ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond was shown as an average value of the result of two sugar chain analyses.

6. Evaluation of binding activity of antibody to CCR4 partial peptide (ELISA)

[0509] Binding activities of the six kinds of the anti-CCR4 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond prepared in the item 5 of Example 4 to a CCR4 partial peptide were measured in accordance with the method described in the item 2 of Example 4.

[0510] As a result, as shown in Fig. 9, the six kinds of the anti-CCR4 chimeric antibodies showed almost the same CCR4-binding activity, and it was found that the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond does not have influence on the antigen-binding activity of the antibody.

7. Evaluation of ADCC activity on human CCR4-high expressing clone

[0511] The ADCC activity of the anti-CCR4 chimeric antibodies against a human CCR4-high highly expressing cell

was measured as follows.

(1) Preparation of target cell suspension

[0512] Cells (1.5×10^6) of a human CCR4-highly expressing cell, CCR4/EL-4 cell, described in WO01/64754 were prepared and a 5.55 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ was added thereto, followed by reaction at 37°C for 1.5 hours to thereby label the cells with a radioisotope. After the reaction, the cells were washed three times by suspension in a medium and subsequent centrifugation, resuspended in the medium and then incubated at 4°C for 30 minutes on ice for spontaneous dissociation of the radioactive substance. After centrifugation, the cells were adjusted to give a density of 2×10^5 cells/ml by adding 7.5 ml of the medium and used as a target cell suspension.

(2) Preparation of human effector cell suspension

[0513] From a healthy donor, 60 ml of peripheral blood was collected, 0.6 ml of heparin sodium (manufactured by Shimizu Pharmaceutical) was added thereto, followed by gently mixing. The mixture was centrifuged (800 g, 20 minutes) to isolate a mononuclear cell layer using Lymphoprep (manufactured by AXIS SHIELD) in accordance with the manufacture's instructions. The cells were washed by centrifuging (1,400 rpm, 5 minutes) three times in a medium and then re-suspended in the medium to give a density of 5×10^6 cells/ml and used as a human effector cell suspension.

(3) Measurement of ADCC activity

[0514] The target cell suspension prepared in the above item (1) was dispensed at 50 μl (1×10^4 cells/well) into each well of a 96 well U-bottom plate (manufactured by Falcon). Next, 100 μl of the effector cell suspension prepared in the above item (2) was added thereto (5×10^5 cells/well, ratio of the human effector cells to the target cells was 50 : 1). Furthermore, each of the anti-CCR4 chimeric antibodies was added thereto to give a final concentration of 0.0001 to 10 $\mu\text{g/ml}$, followed by reaction at 37°C for 4 hours. After the reaction, the plate was centrifuged and the amount of ^{51}Cr in the supernatant was measured with a γ -counter. An amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure as above using the medium alone instead of the human effector cell suspension and antibody solution, and measuring the amount of ^{51}Cr in the supernatant. An amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure as above using a 1 mol/L hydrochloric acid solution instead of the antibody solution and human effector cell suspension, and measuring the amount of ^{51}Cr in the supernatant. The ADCC activity (%) was calculated based on the above-mentioned equation (I).

[0515] Figs. 10 and 11 show results of the measurement of ADCC activity of the anti-CCR4 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond at various concentrations (0.001 to 10 $\mu\text{g/ml}$) using effector cells of two healthy donors (A and B), respectively. As shown in Figs. 10 and 11, the ADCC activity of the anti-CCR4 chimeric antibodies showed a tendency to increase in proportion to the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond at each antibody concentration. The ADCC activity decreases when the antibody concentration is low. At an antibody concentration of 0.01 $\mu\text{g/ml}$, the antibody in which the ratios of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was 27%, 39% and 46%, respectively, showed almost the same high ADCC activity but the ADCC activity was low in the antibody in which the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is less than 20%. These results did not change even if the effector cell having donor was different.

Example 5

[0516] Determination of transcription product of $\alpha 1,6$ -fucosyltransferase (FUT8) gene in host clone:

1. Preparation of single-stranded cDNA from various clones

[0517] Single-stranded cDNAs were prepared from *dhfr*-deleted CHO/DG44 cells and rat myeloma YB2/0 cells by the following procedure.

[0518] The CHO/DG44 cells were suspended in IMDM medium (manufactured by Life Technologies) supplemented with 10% FBS (manufactured by Life Technologies) and $1 \times$ concentration HT supplement (manufactured by Life Technologies), and 15 ml of the suspension was inoculated into T75 flask for adhesion cell culture use (manufactured by Greiner) at a density of 2×10^5 cells/ml. Also, the YB2/0 cells were suspended in RPMI 1640 medium (manufactured by Life Technologies) supplemented with 10% FBS (manufactured by Life Technologies) and 4 mmol/l L-GLN (manu-

factured by Life Technologies), and 15 ml of the suspension was inoculated into T75 flask for suspension cell culture (manufactured by Greiner) at a density of 2×10^5 cells/ml. They were cultured at 37°C in a 5% CO₂ incubator, and 1×10^7 of respective host cells were recovered on the 1st, 2nd, 3rd, 4th and 5th days of the culturing to extract total RNA using RNAeasy (manufactured by QIAGEN) in accordance with the manufacture's instructions.

[0519] The total RNA was dissolved in 45 µl of sterile water, 1 µl of RQ 1 RNase-Free DNase (manufactured by Promega), 5 µl of the attached 10 × DNase buffer and 0.5 µl of RNasin Ribonuclease Inhibitor (manufactured by Promega) were added thereto, followed by reaction at 37°C for 30 minutes to degrade genomic DNA contaminated in the sample. After the reaction, the total RNA was purified again using RNAeasy (manufactured by QIAGEN) and dissolved in 50 µl of sterile water.

[0520] In a 20 µl of the reaction mixture using oligo(dT) as a primer, single-stranded cDNA was synthesized from 3 µg of each of the obtained total RNA samples by reverse transcription reaction using SUPERScript™ Preamplification System for First Strand cDNA Synthesis (manufactured by Life Technologies) in accordance with the manufacture's instructions. For the cloning of FUT8 and β-actin derived from respective host cells, 1 × concentration solution of the reaction solution was used, and for the determination of each gene transcription amount by competitive PCR, 50 fold-diluted aqueous solution of the reaction solution were used. The solutions were stored at -80°C until use.

2. Preparation method of cDNA partial fragments of Chinese hamster FUT8 and rat FUT8

[0521] Each cDNA partial fragment of Chinese hamster FUT8 and rat FUT8 was prepared by the following procedure (Fig. 12).

[0522] First, primers (represented by SEQ ID NOs:4 and 5) specific for nucleotide sequences common to human FUT8 cDNA [*J. Biochem.*, 121, 626 (1997)] and swine FUT8 cDNA [*J. Biol. Chem.*, 271, 27810 (1995)] were designed.

[0523] Next, 25 µl of a reaction solution [ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs and 0.5 µmol/l of the above gene-specific primers (SEQ ID NOs:4 and 5)] containing 1 µl of each of the cDNA prepared from CHO cell and cDNA prepared from YB2/0 cell, both obtained in the item 1 of Example 5 on the second day after culturing, and polymerase chain reaction (hereinafter referred to as "PCR") was carried out by using a DNA polymerase ExTaq (manufactured by Takara Shuzo). The PCR was carried out by heating at 94°C for 1 minute, subsequent 30 cycles of heating at 94°C for 30 seconds, 55°C for 30 seconds and 72°C for 2 minutes as one cycle, and further heating at 72°C for 10 minutes.

[0524] After the PCR, the reaction solution was subjected to 0.8% agarose gel electrophoresis, and a specific amplified fragment of 979 bp was purified using GENE CLEAN Spin Kit (manufactured by BIO 101) and eluted with 10 µl of sterile water (hereinafter, the method was used for the purification of DNA fragments from agarose gel). Into a plasmid pCR2.1, 4 µl of the amplified fragment was employed to insert in accordance with the manufacture's instructions of TOPO TA Cloning Kit (manufactured by Invitrogen), and *E. coli* XL1-Blue was transformed with the reaction solution by the method of Cohen et al. [*Proc. Natl. Acad. Sci. USA*, 69, 2110 (1972)] (hereinafter, the method was used for the transformation of *E. coli*). Plasmid DNA samples were isolated in accordance with a known method [*Nucleic Acids Research*, 7, 1513 (1979)] (hereinafter, the method was used for the isolation of plasmid) from cDNA-inserted six clones among the obtained kanamycin-resistant colonies.

[0525] The nucleotide sequences of cDNAs inserted into the plasmids were determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method of the manufacture's instructions. It was confirmed that all of the inserted cDNAs of which sequences were determined by the method encode the open reading frame (ORF) partial sequences of Chinese hamster FUT8 and rat FUT8 (represented by SEQ ID NOs:6 and 7). Among these, plasmid DNA samples containing absolutely no reading error by the PCR in the sequences were selected. Herein, these plasmids were named CHFT8-pCR2.1 and YBFT8-pCR2.1.

3. Preparation of Chinese hamster β-actin and rat β-actin cDNA

[0526] Chinese hamster β-actin and rat β-actin were prepared by the following procedure (Fig. 13).

[0527] First, a forward primer specific for a common sequence containing translation initiation codon (represented by SEQ ID NO:8) and reverse primers specific for respective sequences containing translation termination codon (represented by SEQ ID NOs:9 and 10) were designed from Chinese hamster β-actin genomic sequence (GenBank, U20114) and rat β-actin genomic sequence [*Nucleic Acids Research*, 11, 1759 (1983)].

[0528] Next, 25 µl of a reaction solution [1 × concentration KOD buffer #1 (manufactured by Toyobo), 0.2 mmol/l dNTPs, 1 mmol/l MgCl₂, 0.4 µmol/l of the above gene-specific primers (SEQ ID NOs:8 and 9, or SEQ ID NOs:8 and 10) and 5% DMSO] containing 1 µl of each of the cDNA prepared from CHO cell and cDNA prepared from YB2/0 cell, both obtained in the item 1 of Example 5 on the second day after culturing was prepared, and PCR was carried out by using a DNA polymerase KOD (manufactured by Toyobo). The PCR was carried out by heating at 94°C for 4 minutes

and subsequent 25 cycles of heating at 98°C for 15 seconds, 65°C for 2 seconds and 74°C for 30 seconds as one cycle.

[0529] After the PCR, the reaction solution was subjected to 0.8% agarose gel electrophoresis, and a specific amplified fragment of 1128 bp was purified. The DNA fragment was subjected to 5'-terminal phosphorylation using MEG-ALABEL (manufactured by Takara Shuzo) in accordance with the manufacture's instructions. The DNA fragment was recovered from the reaction solution using an ethanol precipitation method and dissolved in 10 µl of sterile water.

[0530] Separately, 3 µg of a plasmid pBluescript II KS(+) (manufactured by Stratagene) was dissolved in 35 µl of NEBuffer 2 (manufactured by New England Biolabs), and 16 units of a restriction enzyme *EcoRV* (manufactured by Takara Shuzo) were added thereto for digestion reaction at 37°C for 3 hours. To the reaction solution, 35 µl of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.5 µl of *E. coli* C15-derived alkaline phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 30 minutes to thereby dephosphorylate the DNA terminus. The reaction solution was extracted with phenol/chloroform, followed by ethanol precipitation, and the recovered DNA fragment was dissolved in 100 µl of sterile water.

[0531] Each 4 µl of the amplified fragment prepared from Chinese hamster cDNA and the amplified fragment (1192 bp) prepared from rat cDNA was mixed with 1 µl of the *EcoRV*-*EcoRV* fragment (about 3.0 Kb) prepared from plasmid pBluescript II KS(+) and 5 µl of Ligation High (manufactured by Toyobo) for ligation reaction at 16°C for 30 minutes. Using the reaction solution, *E. coli* XL1-Blue was transformed, and plasmid DNA samples were isolated respectively from the obtained ampicillin-resistant clones in accordance with a known method.

[0532] The nucleotide sequence of each cDNA inserted into the plasmid was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method of the manufacture's instructions. It was confirmed that all of the inserted cDNAs of which sequences were determined by the method encode the ORF full sequences of Chinese hamster β-actin or rat β-actin. Among these, plasmid DNA samples containing absolutely no reading error of bases by the PCR in the sequences were selected. Herein, the plasmids are called CHAc-pBS and YBac-pBS.

4. Preparation of FUT8 standard and internal control

[0533] In order to measure a transcription amount of mRNA derived from FUT8 gene in each cell, CHFT8-pCR2.1 or YBFT8-pCR2.1, as plasmids in which cDNA partial fragments prepared in the item 2 of Example 5 from Chinese hamster FUT8 or rat FUT8 were inserted into pCR2.1, respectively, were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the standards for the preparation of a calibration curve. CHFT8d-pCR2.1 and YBFT8d-pCR2.1, which were obtained from the CHFT8-pCR2.1 and YBFT8-pCR2.1, by deleting 203 bp between *Scal* and *HindIII*, an inner nucleotide sequence of Chinese hamster FUT8 and rat FUT8, respectively, were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the internal standards for FUT8 amount determination. Details thereof are described below.

[0534] Chinese hamster FUT8 and rat FUT8 standards were prepared as follows. In 40 µl of NEBuffer 2 (manufactured by New England Biolabs), 2 µg of the plasmid CHFT8-pCR2.1 was dissolved, 24 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. Separately, 2 µg of the plasmid YBFT8-pCR2.1 was dissolved in 40 µl of NEBuffer 2 (manufactured by New England Biolabs), and 24 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. By subjecting a part of each of the reaction solutions to 0.8% agarose gel electrophoresis, it was confirmed that an *EcoRI*-*EcoRI* fragment (about 1 Kb) containing each of cDNA partial fragments of Chinese hamster FUT8 and rat FUT8 was separated from the plasmids CHFT8-pCR2.1 and YBFT8-pCR2.1 by the above restriction enzyme digestion reactions. Each of the reaction solutions was diluted with 1 µg/ml of baker's yeast t-RNA (manufactured by SIGMA) to give a concentration of 0.02 fg/µl, 0.2 fg/µl, 1 fg/µl, 2 fg/µl, 10 fg/µl, 20 fg/µl and 100 fg/µl and used as the Chinese hamster FUT8 and rat FUT8 standards.

[0535] Controls of Chinese hamster FUT8 and rat FUT8 were prepared as follows (Fig. 14). By using a DNA polymerase KOD (manufactured by Toyobo), 25 µl of a reaction solution [1× concentration KOD buffer #1 (manufactured by Toyobo), 0.2 mmol/l dNTPs, 1 mmol/l MgCl₂, 0.4 µmol/l gene-specific primers (SEQ ID NOs:11 and 12) and 5% DMSO] containing 5 ng of CHFT8-pCR2.1 or YBFT8-pCR2.1 was prepared, and PCR was carried out. The PCR was carried out by heating at 94°C for 4 minutes and subsequent 25 cycles of heating at 98°C for 15 seconds, 65°C for 2 seconds and 74°C for 30 seconds as one cycle. After the PCR, the reaction solution was subjected to 0.8% agarose gel electrophoresis, and a specific amplified fragment of about 4.7 Kb was purified. The DNA fragment was subjected to 5'-terminal phosphorylation using MEGALABEL (manufactured by Takara Shuzo) in accordance with the manufacture's instructions, and then the DNA fragment was recovered from the reaction solution by ethanol precipitation and dissolved in 50 µl of sterile water. The above obtained DNA fragment (5 µl, about 4.7 kb) and 5 µl of Ligation High (manufactured by Toyobo) were mixed, followed by self-cyclization reaction at 16°C for 30 minutes.

[0536] Using the reaction solution, *E. coli* DH5α was transformed, and plasmid DNA samples were isolated in accordance with a known method from the obtained ampicillin-resistant clones. The nucleotide sequence of each plasmid

DNA was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer), and it was confirmed that a 203 bp inner nucleotide sequence between *Scal* and *HindIII* of Chinese hamster FUT8 or rat FUT8 was deleted. The obtained plasmids were named CHFT8d-pCR2.1 or YBFT8d-pCR2.1, respectively.

[0537] Next, 2 µg of the plasmid CHFT8d-pCR2.1 was dissolved in 40 µl of NEBuffer 2 (manufactured by New England Biolabs), and 24 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. Separately, 2 µg of the plasmid YBFT8d-pCR2.1 was dissolved in 40 µl of NEBuffer 2 (manufactured by New England Biolabs), and 24 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. A part of each of the reaction solutions was subjected to 0.8% agarose gel electrophoresis, and it was confirmed that an *EcoRI*-*EcoRI* fragment (about 800 bp) containing a fragment from which 203 bp of the inner nucleotide sequences of Chinese hamster FUT8 or rat FUT8 partial fragments was deleted was separated from the plasmids CHFT8d-pCR2.1 or YBFT8d-pCR2.1 by the above restriction enzyme digestion reactions. Dilutions of 2 fg/µl were prepared from the reaction solutions using 1 µg/ml baker's yeast t-RNA (manufactured by SIGMA) and used as the Chinese hamster FUT8 or rat FUT8 internal controls.

5. Preparation of β-actin standard and internal control

[0538] In order to measure a transcription amount of mRNA derived from β-actin gene in various host cells, CHAc-pBS and YBac-pBS, as plasmids in which the ORF full length of each cDNA of Chinese hamster β-actin and rat β-actin prepared in the item 3 of Example 5 was inserted into pBluescript II KS(+), respectively, were digested with restriction enzymes *HindIII* and *PstI* and restriction enzymes *HindIII* and *KpnI*, respectively, and the digested linear DNAs were used as the standards for the preparation of a calibration curve. CHAc-pBS and YBac-pBS which were obtained from the CHAc-pBS and YBac-pBS by deleting 180 bp between *DraIII* and *DraIII* of an inner nucleotide sequence of Chinese hamster β-actin and rat β-actin were digested with restriction enzymes *HindIII* and *PstI* and restriction enzymes *HindIII* and *KpnI*, respectively, and the digested linear DNAs were used as the internal controls for β-actin amount determination. Details thereof are described below.

[0539] Chinese hamster β-actin and rat β-actin standards were prepared as follows. In 40 µl of NEBuffer 2 (manufactured by New England Biolabs), 2 µg of the plasmid CHAc-pBS was dissolved, and 25 units of a restriction enzyme *HindIII* (manufactured by Takara Shuzo) and 20 units of *PstI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. Separately, 2 µg of the plasmid YBac-pBS was dissolved in 40 µl of NEBuffer 2 (manufactured by New England Biolabs), and 25 units of a restriction enzyme *HindIII* (manufactured by Takara Shuzo) and 24 units of *KpnI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. A part of each of the reaction solutions was subjected to 0.8% agarose gel electrophoresis, and it was confirmed that a *HindIII*-*PstI* fragment and a *HindIII*-*KpnI* fragment (about 1.2 Kb) containing the full length ORF of each cDNA of Chinese hamster β-actin and rat β-actin were separated from the plasmids CHAc-pBS and YBac-pBS by the above restriction enzyme digestion reactions. Each of the reaction solutions was diluted with 1 µg/ml baker's yeast t-RNA (manufactured by SIGMA) to give a concentration 2 pg/µl, 1 pg/µl, 200 fg/µl, 100 fg/µl and 20 fg/µl and used as the Chinese hamster β-actin or rat β-actin standards.

[0540] Chinese hamster β-actin and rat β-actin internal controls were prepared as follows (Fig. 15). In 100 µl of NEBuffer 3 (manufactured by New England Biolabs) containing 100 ng/µl of BSA (manufactured by New England Biolabs), 2 µg of CHAc-pBS was dissolved, and 10 units of a restriction enzyme *DraIII* (manufactured by New England Biolabs) were added thereto, followed by digestion reaction at 37°C for 3 hours. DNA fragments were recovered from the reaction solution by ethanol precipitation and the DNA termini were changed to blunt-ends using DNA Blunting Kit (manufactured by Takara Shuzo) in accordance with the manufacture's instructions, and then the reaction solution was divided into two equal parts. First, to one part of the reaction solution, 35 µl of 1 mol/l Tris-HCl buffer (pH 8.0) and 3.5 µl of *E. coli* C15-derived alkaline phosphatase (manufactured by Takara Shuzo) were added thereto, followed by reaction at 65°C for 30 minutes for dephosphorylating the DNA termini. The DNA fragment was recovered by carrying out dephosphorylation treatment, phenol/chloroform extraction treatment and ethanol precipitation treatment and then dissolved in 10 µl of sterile water. The remaining part of the reaction solution was subjected to 0.8% agarose gel electrophoresis to purify a DNA fragment of about 1.1 Kb containing the ORF partial fragment of Chinese hamster β-actin.

[0541] The dephosphorylated *DraIII*-*DraIII* fragment (0.5 µl), 4.5 µl of the *DraIII*-*DraIII* fragment of about 1.1 Kb and 5 µl of Ligation High (manufactured by Toyobo) were mixed, followed by ligation reaction at 16°C for 30 minutes. Using the reaction solution, *E. coli* DH5α was transformed, and plasmid DNAs were isolated in accordance with a known method from the obtained ampicillin-resistant colonies. The nucleotide sequence of each plasmid DNA was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer), and it was confirmed that a Chinese hamster β-actin *DraIII*-*DraIII* 180 bp

inserted into the plasmid was deleted. The plasmid was named CHAcD-pBS.

[0542] Also, a plasmid in which rat β -actin *Dralll-Dralll* 180 bp was deleted was prepared via same steps of CHAcD-pBS. The plasmid was named YBAcD-pBS.

[0543] Next, 2 μ g of the plasmid CHAcD-pBS was dissolved in 40 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 25 units of a restriction enzyme *HindIII* (manufactured by Takara Shuzo) and 20 units of *PstI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. Separately, 2 μ g of the plasmid YBAcD-pBS was dissolved in 40 μ l of NEBuffer 2 (manufactured by New England Biolabs), and 25 units of a restriction enzyme *HindIII* (manufactured by Takara Shuzo) and 24 units of *KpnI* (manufactured by Takara Shuzo) were added thereto, followed by digestion reaction at 37°C for 3 hours. A part of each of the reaction solutions was subjected to 0.8% agarose gel electrophoresis, and it was confirmed that an *HindIII-PstI* fragment and *HindIII-KpnI* fragment (about 1.0 Kb) containing a fragment in which 180 bp of the inner nucleotide sequence of the ORF full length of each cDNA of Chinese hamster β -actin and rat β -actin was deleted were separated from the plasmids CHAcD-pBS and YBAcD-pBS by the restriction enzyme digestion reactions. Dilutions of 200 fg/ μ l were prepared from the reaction solutions using 1 μ g/ml baker's yeast t-RNA (manufactured by SIGMA) and used as the Chinese hamster β -actin and rat β -actin internal controls.

6. Determination of transcription amount by competitive PCR

[0544] Competitive PCR was carried out by using the FUT8 internal control DNA prepared in the item 4 of Example 5 and the host cell-derived cDNA obtained in the item 1 of Example 5 as the templates, and the determined value of the FUT8 transcription product in the host clone was calculated from the relative value of the amount of the amplified product derived from each template. On the other hand, since it is considered that the β -actin gene is transcribed constantly in each cell and its transcription amount is approximately the same between cells, transcription amount of the β -actin gene was determined as an indication of the efficiency of synthesis reaction of cDNA derived from each host clone. That is, the PCR was carried out by using the β -actin internal control DNA prepared in the item 5 of Example 5 and the host cell-derived cDNA obtained in the item 1 of Example 5 as the templates, the determined value of the β -actin transcription product in the host clone was calculated from the relative value of the amount of the amplified product derived from each template. Details thereof are described below.

[0545] The FUT8 transcription product was determined by the following procedure. First, a set of sequence-specific primers (represented by SEQ ID NOs:13 and 14) common to the inner sequences of the ORF partial sequences of Chinese hamster FUT8 and rat FUT8 obtained in the item 2 of Example 5 were designed.

[0546] Next, PCR was carried out by using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 μ l in total volume of a reaction solution [1 \times concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 μ mol/l of the above gene-specific primers (SEQ ID NOs:13 and 14) and 5% DMSO] containing 5 μ l of 50 fold-diluted cDNA solution prepared from each of respective host clone in the item 1 of Example 5 and 5 μ l (10 fg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 32 cycles of heating at 94°C for 1 minute, 60°C for 1 minute and 72°C for 1 minute as one cycle.

[0547] Also, PCR was carried out in a series of reaction in which 5 μ l (0.1 fg, 1 fg, 5 fg, 10 fg, 50 fg, 100 fg, 500 fg or 1 pg) of the FUT8 standard plasmid obtained in the item 4 of Example 5 was added instead of the each host clone-derived cDNA, and used in the preparation of a calibration curve for the FUT8 transcription amount.

[0548] The β -actin transcription product was determined by the following procedure. First, two sets of respective gene-specific primers common to the inner sequences of the ORF full lengths of Chinese hamster β -actin and rat β -actin obtained in the item 3 of Example 5 were designed (the former are represented by SEQ ID NOs:15 and 16, and the latter are represented by SEQ ID NOs:17 and 18).

[0549] Next, PCR was carried out by using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 μ l in total volume of a reaction solution [1 \times concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 μ mol/l of the above gene-specific primers (SEQ ID NOs:15 and 16, or SEQ ID NOs:17 and 18) and 5% DMSO] containing 5 μ l of 50 fold-diluted cDNA solution prepared from respective host clone in the item 1 of Example 5 and 5 μ l (1 pg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 17 cycles of heating at 94°C for 30 seconds, 65°C for 1 minute and 72°C for 2 minutes as one cycle.

[0550] Also, PCR was carried out in a series of reaction in which 5 μ l (10 pg, 5 pg, 1 pg, 500 fg or 100 fg) of the β -actin standard plasmid obtained in the item 5 of Example 5 was added instead of the each host clone-derived cDNA, and used in the preparation of a calibration curve for the β -actin transcription amount.

Table 1

| Target gene | Primer set * | Size of PCR amplification product (bp) | |
|-------------------------------------|--|--|------------|
| | | Target | Competitor |
| FUT8 | F: 5'-GTCCATGGTGATCCTGCAGTGTGG-3' R: 5'-CACCAATGATATCTCCAGGTTCC-3' | 638 | 431 |
| β -Actin (Chinese hamster) | F: 5'-GATATCGCTGCGCTCGTTGTCGAC-3' R: 5'-CAGGAAGGAAGGCTGGAAGAGAGC-3' | 789 | 609 |
| β -Actin (Rat) | F: 5'-GATATCGCTGCGCTCGTCGTCGAC-3' R: 5'-CAGGAAGGAAGGCTGGAAGAGAGC-3' | 789 | 609 |

* F: forward primer, R: reverse primer

[0551] By carrying out PCR using the primer set shown in Table 1, a DNA fragment having a size shown in the target column of Table 1 can be amplified from each gene transcription product and each standard, and a DNA fragment having a size shown in the competitor column of Table 1 can be amplified from each internal control.

[0552] After 7 μ l of each of the solutions after PCR was subjected to 1.75% agarose gel electrophoresis, the gel was stained by soaking it for 30 minutes in 1 x concentration SYBR Green I Nucleic Acid Gel Stain (manufactured by Molecular Probes). The amount of the amplified DNA fragment was measured by calculating luminescence intensity of each amplified DNA using a fluoro-imager (FluorImager SI; manufactured by Molecular Dynamics).

[0553] The amount of an amplified product formed by PCR using a standard plasmid as the template was measured by the above-mentioned method, and a calibration curve was prepared by plotting the measured values against the amounts of the standard plasmid. Using the calibration curve, the amount of cDNA of a gene of interest in each clone was calculated from the amount of the amplified product when each expression clone-derived total cDNA was used as the template, and the amount was defined as the mRNA transcription amount in each clone.

[0554] The amount of the FUT8 transcription product in each host clone when a rat FUT8 sequence was used as the standard and internal control is shown in Fig. 16. Throughout the culturing period, the CHO clone showed a transcription amount 10-fold or higher than that of the YB2/0 clone. The tendency was also found when a Chinese hamster FUT8 sequence was used as the standard and internal control.

[0555] Also, the FUT8 transcription amounts are shown in Table 2 as relative values to the amount of the β -actin transcription product. Throughout the culturing period, the FUT8 transcription amount in the YB2/0 clone was around 0.1% of β -actin while it was 0.5% to 2% in the CHO/DG44 clone.

[0556] The results shows that the amount of the FUT8 transcription product in YB2/0 clone was significantly smaller than that in the CHO/DG44 clone.

Table 2

| Clone | Culture days | | | | |
|-------|--------------|------|------|------|------|
| | 1st | 2nd | 3rd | 4th | 5th |
| CHO | 1.95 | 0.90 | 0.57 | 0.52 | 0.54 |
| YB2/0 | 0.12 | 0.11 | 0.14 | 0.08 | 0.07 |

Example 6

[0557] Determination of transcription product of FUT8 gene in anti-GD3 chimeric antibody-producing clone:

1. Preparation of single-stranded cDNA derived from various antibody-producing clones

[0558] Single-stranded cDNA was prepared from anti-GD3 chimeric antibody-producing cell clones DCHI01-20 and 61-33 as follows. The clone DCHI01-20 is a transformant clone derived from the CHO/DG44 cell described in item 2 (2) of Example 1. Also, the clone 61-33 is a clone obtained by carrying out serum-free adaptation of YB2/0-derived transformant cell clone 7-9-51 (FERM BP-6691, International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology) and then carrying out single cell isolation by two limiting dilution.

[0559] The clone DCHI01-20 were suspended in EXCELL 302 medium (manufactured by JRH BIOSCIENCES) sup-

plemented with 3 mmol/l L-GLN (manufactured by Life Technologies), 0.3% PLURONIC F-68 (manufactured by Life Technologies) and 0.5% fatty acid concentrate (manufactured by Life Technologies), and 15 ml of the suspension was inoculated into T75 flask for suspension cell culture (manufactured by Greiner) at a density of 2×10^5 cells/ml. Also, cells of the clone 61-33 were suspended in Hybridoma-SFM medium (manufactured by Life Technologies) supplemented with 0.2% BSA, and 15 ml of the suspension was inoculated into T75 flask for suspension cell culture (manufactured by Greiner) at a density of 2×10^5 cells/ml. They were cultured at 37°C in a 5% CO₂ incubator, and 1, 2, 3, 4 and 5 days after culturing, 1×10^7 of respective host cells were recovered to extract total RNA using RNeasy (manufactured by QIAGEN) in accordance with the manufacture's instructions.

[0560] The total RNA was dissolved in 45 µl of sterile water, and 1 µl of RQ1 RNase-Free DNase (manufactured by Promega), 5 µl of the attached 10× DNase buffer and 0.5 µl of RNasin Ribonuclease Inhibitor (manufactured by Promega) were added thereto, followed by reaction at 37°C for 30 minutes to degrade genomic DNA contaminated in the sample. After the reaction, the total RNA was purified again using RNeasy (manufactured by QIAGEN) and dissolved in 50 µl of sterile water.

[0561] In a 20 µl reaction mixture using oligo(dT) as a primer, single-stranded cDNA was synthesized from 3 µg of each of the obtained total RNA samples by reverse transcription reaction using SUPERScript™ Preamplification System for First Strand cDNA Synthesis (manufactured by Life Technologies) in accordance with the manufacture's instructions. The reaction solution was diluted 50-fold with water and stored at -80°C until use.

2. Determination of transcription amounts of each gene by competitive PCR

[0562] The transcription amount of each of the genes of the cDNA derived from the antibody-producing clone obtained in the item 1 of Example 6 was determined by competitive PCR in accordance with the item 6 of Example 5.

[0563] The FUT8 gene-derived mRNA transcription amount in each antibody-producing clone was determined by the following procedure.

[0564] CHFT8-pCR2.1 and YBFT8-pCR2.1, as plasmids in which cDNA partial fragments prepared in item 2 of Example 5 from Chinese hamster FUT8 and rat FUT8, respectively, were inserted into pCR2.1, were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the standards in the preparation of a calibration curve for determining the FUT8 transcription amount.

[0565] CHFT8d-pCR2.1 and YBFT8d-pCR2.1, which were obtained by deleting 203 bp between *Scal* and *HindIII* of an inner nucleotide sequence of Chinese hamster FUT8 and rat FUT8, respectively, in the item 4 of Example 9 were digested with a restriction enzyme *EcoRI*, and the obtained linear DNAs were used as the internal controls for FUT8 amount determination.

[0566] PCR was carried out by using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 µl in total volume of a reaction solution [1 × concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 µmol/l FUT8 gene-specific primers (SEQ ID NOs:13 and 14) and 5% DMSO] containing 5 µl of 50 fold-diluted cDNA solution derived from each of the antibody-producing clone in the item 1 of Example 6 and 5 µl (10 fg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 32 cycles of heating at 94°C for 1 minute, 60°C for 1 minute and 72°C for 1 minute as one cycle.

[0567] Also, PCR was carried out in a series of reaction in which 5 µl (0.1 fg, 1 fg, 5 fg, 10 fg, 50 fg, 100 fg, 500 fg or 1 pg) of the FUT8 standard plasmid was added instead of the each antibody-producing clone-derived cDNA, and used in the preparation of a calibration curve for the FUT8 transcription amount. In this case, 1 µg/ml of a baker's yeast t-RNA (manufactured by SIGMA) was used for the dilution of the standard plasmid.

[0568] On the other hand, since it is considered that the β-actin gene is transcribed constantly in each cell and its transcription amount is approximately the same between cells, the transcription amount of the β-actin gene was determined as an index of the efficiency of synthesis reaction of cDNA in each antibody-producing clone.

[0569] CHAc-pBS and YBAc-pBS as plasmids in which the ORF full length of each cDNA of Chinese hamster β-actin and rat β-actin prepared in the item 3 of Example 5 were inserted into pBluescript II KS(+), respectively, were digested with restriction enzymes *HindIII* and *KpnI*, and the obtained linear DNAs were used as the standards in the preparation of a calibration curve for determining the β-actin gene transcription amount.

[0570] CHAc-d-pBS and YBAc-d-pBS which were obtained by deleting 180 bp between *DraIII* and *DraIII* of an inner nucleotide sequence of Chinese hamster β-actin and rat β-actin, respectively in the item 5 of Example 5, were digested with restriction enzymes *HindIII* and *KpnI*, and the obtained linear DNAs were used as the internal controls for β-actin determination.

[0571] PCR was carried out by using a DNA polymerase ExTaq (manufactured by Takara Shuzo) in 20 µl in total volume of a reaction solution [1 × concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 0.5 µmol/l β-actin-specific primers (SEQ ID NOs:17 and 18) and 5% DMSO] containing 5 µl of 50 fold-diluted cDNA solution derived from each of the antibody-producing clones and 5 µl (1 pg) of the plasmid for internal control. The PCR was carried out by heating at 94°C for 3 minutes and subsequent 17 cycles of heating at 94°C for 30 seconds, 65°C for 1

minute and 72°C for 2 minutes as one cycle. Also, PCR was carried out in a series of reaction in which 10 pg, 5 pg, 1 pg, 500 fg or 100 fg of the β -actin standard plasmid was added instead of the each antibody-producing clone-derived cDNA, and used in the preparation of a calibration curve for the β -actin transcription amount. In this case, 1 μ g/ml of a baker's yeast t-RNA (manufactured by SIGMA) was used for the dilution of standard plasmid.

[0572] By PCR using the primer set described in Table 1, a DNA fragment having a size shown in the target column of Table 1 can be amplified from each gene transcription product and each standard, and a DNA fragment having a size shown in the competitor column of Table 1 can be amplified from each internal control.

[0573] After 7 μ l of the solutions after PCR was subjected to 1.75% agarose gel electrophoresis, the gel was stained by soaking it for 30 minutes in 1 \times concentration SYBR Green I Nucleic Acid Gel Stain (manufactured by Molecular Probes). The amount of the amplified DNA fragment was measured by calculating luminescence intensity of each amplified DNA using a fluoro-imager (FluorImager SI; manufactured by Molecular Dynamics).

[0574] The amount of the amplified product formed by PCR which used a standard plasmid as the template was measured by the above method, and a calibration curve was prepared by plotting the measured values against the amounts of the standard plasmid. Using the calibration curve, the amount of cDNA of a gene of interest in each clone was calculated from the amount of the amplified product when each antibody-producing clone-derived total cDNA was used as the template, and the value was defined as the mRNA transcription amount in each clone.

[0575] The FUT8 transcription amounts are shown in Table 3 as relative values to the amount of the β -actin transcription product. Throughout the culturing period, the FUT8 transcription amount in the YB2/0 cell-derived antibody-producing clone 61-33 was 0.3% or less of β -actin while the FUT8 transcription amount in the CHO-derived antibody-producing clone DCH101-20 was 0.7% to 1.5% in the CHO cell-derived antibody-producing cell. The results shows that the amount of the FUT8 transcription product in the YB2/0 cell-derived antibody-producing clone was significantly less than that in the CHO cell-derived antibody-producing clone.

Table 3

| Clone | Culture days | | | | |
|-----------|--------------|------|------|------|-------|
| | 1st | 2nd | 3rd | 4th | 5th |
| DCH101-20 | 0.75 | 0.73 | 0.99 | 1.31 | 1.36 |
| 61-33 | 0.16 | 0.19 | 0.24 | 0.30 | <0.10 |

Example 7

Preparation of lectin-resistant CHO/DG44 cell and production of antibody using the cell:

1. Preparation of lectin-resistant CHO/DG44

[0576] CHO/DG44 cells were cultured in a 75 cm² flask for adhesion culture (manufactured by Greiner) in IMDM-FBS (10)-HT(1) medium [IMDM medium comprising 10% of FBS and 1 \times concentration of HT supplement (manufactured by GIBCO BRL)] to grow until they reached a stage of just before confluent. After washing the cells with 5 ml of PBS (manufactured by Invitrogen), 1.5 ml of 0.05% trypsin (manufactured by Invitrogen) diluted with Dulbecco PBS was added thereto and cultured at 37°C for 5 minutes to remove the cells from the flask bottom. The removed cells were recovered by a centrifugation operation generally used in cell culture and suspended in IMDM-FBS(10) medium to give a density of 1 \times 10⁵ cells/ml, and then 0.1 μ g/ml of an alkylating agent *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (hereinafter referred to as "MNNG", manufactured by Sigma) was added or not added thereto. After culturing at 37°C for 3 days in a CO₂ incubator (manufactured by TABAI), the culture supernatant was discarded, and the cells were again washed, removed and recovered by the same operations as described above, suspended in IMDM-FBS(10)-HT (1) medium and then inoculated into an adhesion culture 96 well plate (manufactured by IWAKI Glass) to give a density of 1 \times 10³ cells/well. To each well, as the final concentration in medium, 1 mg/ml *Lens culinaris* agglutinin (hereinafter referred to as "LCA", manufactured by Vector), 1 mg/ml *Aleuria aurantia* agglutinin (*Aleuria aurantia* lectin; hereinafter referred to as "AAL", manufactured by Vector) or 1 mg/ml kidney bean agglutinin (*Phaseolus vulgaris* leucoagglutinin; hereinafter referred to as "L-PHA", manufactured by Vector) was added. After culturing at 37°C for 2 weeks in a CO₂ incubator, the appeared colonies were obtained as lectin-resistant clone CHO/DG44. Regarding the obtained lectin-resistant clone CHO/DG44, an LCA-resistant clone was named clone CHO-LCA, an AAL-resistant clone was named clone CHO-AAL and an L-PHA-resistant clone was named clone CHO-PHA. When the resistance of these clones to various kinds of lectin was examined, it was found that the clone CHO-LCA was also resistant to AAL and the clone CHO-AAL was also resistant LCA. In addition, the clone CHO-LCA and the clone CHO-AAL also showed a resistance to a lectin which recognizes a sugar chain structure identical to the sugar chain structure recognized by LCA and AAL,

namely a lectin which recognizes a sugar chain structure in which 6-position of fucose is bound to 1-position of *N*-acetylglucosamine residue in the reducing end through α -bond in the *N*-glycoside-linked sugar chain. Specifically, it was found that the clone CHO-LCA and the clone CHO-AAL can show resistance and survive even in a medium supplemented with 1 mg/ml at a final concentration of a pea agglutinin (*Pisum sativum* agglutinin; hereinafter referred to as "PSA", manufactured by Vector). In addition, even when the alkylating agent MNNG was not added, it was able to obtain lectin-resistant clones by increasing the number of cells to be treated. Hereinafter, these clones were used in analyses.

2. Preparation of anti-CCR4 chimeric antibody-producing cell

[0577] An anti-CCR4 chimeric antibody expression plasmid pKANTEX2160 was introduced into the three kinds of the lectin-resistant clones obtained in the item 1 of Example 7 by the method described in Example 4, and gene amplification by a drug MTX was carried out to prepare an anti-CCR4 human chimeric antibody-producing clone. By measuring an amount of antibody expression by the ELISA described in the item 2 of Example 4, antibody-expressing transformants were obtained from each of the clone CHO-LCA, the clone CHO-AAL and the clone CHO-PHA. Regarding each of the obtained transformants, a transformant derived from the clone CHO-LCA was named clone CHO/CCR4-LCA, a transformant derived from the clone CHO-AAL was named clone CHO/CCR4-AAL and a transformant derived from the clone CHO-PHA was named clone CHO/CCR4-PHA. Further, the clone CHO/CCR4-LCA, as a name of Nega-13, has been deposited on September 26, 2001, as FERM BP-7756 in International Patent Organism Depositary, National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1, Higashi 1-Chome Tsukuba-shi, Ibarakiken, Japan).

3. Production of high ADCC activity antibody by lectin-resistant CHO cell

[0578] Using the three kinds of the transformants obtained in the item 2 of Example 7, purified antibodies were obtained by the method described in the item 3 of Example 4. The antigen binding activity of the purified anti-CCR4 chimeric antibodies was evaluated by the ELISA described in the item 2 of Example 4. The antibodies produced by all transformants showed an antigen binding activity similar to that of the antibody produced by a recombinant clone (clone 5-03) prepared in Example 4 using normal CHO/DG44 cell as the host. Using these purified antibodies, ADCC activity of each of the anti-CCR4 chimeric antibodies was evaluated in accordance with the method described in the item 7 of Example 4. The results are shown in Fig. 17. In comparison with the antibody produced by the clone 5-03, about 100 fold-increased ADCC activity was observed in the antibodies produced by the clone CHO/CCR4-LCA and the clone CHO/CCR4-AAL. On the other hand, no significant increase in the ADCC activity was observed in the antibody produced by the clone CHO/CCR4-PHA. Also, when ADCC activities of the antibodies produced by the clone CHO/CCR4-LCA and the YB2/0 cell-derived clone were compared in accordance with the method described in the item 7 of Example 4, it was found that the antibody produced by the clone CHO/CCR4-LCA shows higher ADCC activity, similar to the case of the antibody KM2760-1 produced by the YB2/0 cell-derived clone prepared in the item 1 of Example 4 (Fig. 18).

4. Sugar chain analysis of antibody produced by lectin-resistant CHO cell

[0579] Sugar chains of the anti-CCR4 chimeric antibodies purified in the item 3 of Example 7 were analyzed according to the method described in the item 1 of Example 3. Fig. 19 shows elution patterns of the purified PA-treated sugar chains of the various anti-CCR4 chimeric antibodies.

[0580] Table 4 shows the result of ratios (%) of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end obtained by the result of the sugar chain analysis of the anti-CCR4 chimeric antibodies produced by various lectin-resistant clones.

Table 4

| Antibody producing cell | Ratio of α 1,6-Fucose-free sugar chain (%) |
|-------------------------|---|
| Clone 5-03 | 9 |
| Clone CHO/CCR4-LCA | 48 |
| Clone CHO/CCR4-AAL | 27 |
| Clone CHO/CCR4-PHA | 8 |

[0581] In comparison with the antibody produced by the clone 5-03, the ratio of a sugar chain in which 1-position of

fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was increased from 9% to 48% in the antibody produced by the clone CHO/CCR4-LCA. The ratio of α 1,6-fucose-free sugar chains was increased from 9% to 27% in the antibody produced by the clone CHO/CCR4-AAL. On the other hand, changes in the sugar chain pattern and the ratio of α 1,6-fucose-free sugar chains were hardly found in the clone CHO/CCR4-PHA when compared with the clone 5-03.

Example 8

Analysis of lectin-resistant CHO clone:

1. Analysis of expression amount of GMD enzyme in anti-CCR4 chimeric antibody-producing clone CHO/CCR4-LCA

[0582] The expression amount of each of the genes of GMD, GFPP and FX, known as fucose biosynthesis enzymes and FUT8 as a fucosyltransferase, in the anti-CCR4 human chimeric antibody-producing clone CHO/CCR4-LCA obtained in Example 7, was analyzed using RT-PCR method.

(1) Preparation of RNA from various clones

[0583] Each of CHO/DG44 cell, the anti-CCR4 human chimeric antibody-producing clone 5-03 obtained in the item 1(2) of Example 4 and the anti-CCR4 chimeric antibody-producing clone CHO/CCR4-LCA obtained in the item 2 of Example 7 was subcultured at 37°C in a 5% CO₂ incubator and then cultured for 4 days. After culturing, RNA was prepared from 1×10⁷ cells of each clone using RNeasy Protect Mini Kit (manufactured by QIAGEN) in accordance with the manufacture's instructions. Subsequently, single-stranded cDNA was synthesized from 5 µg of each RNA in a 20 µl of a reaction solution using SUPER SCRIPT First-Strand Synthesis System for RT-PCR (manufactured by GIBCO BRL) in accordance with the manufacture's instructions.

(2) Analysis of expression amount of GMD gene using RT-PCR

[0584] In order to amplify GMD cDNA by PCR, a 24 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:32 and a 26 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:33 were prepared based on the CHO cell-derived GMD cDNA sequence shown in the item 1 of Reference Example 2.

[0585] Next, 20 µl of a reaction solution [1× concentration Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 µL of the synthetic DNA primers of SEQ ID NOs:32 and 33] containing 0.5 µl of the single-stranded cDNA derived from each clone in the item 1(1) of Example 8 as the template was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After subjecting 10 µl of the PCR reaction solution to agarose electrophoresis, DNA fragments were stained with Cyber Green (manufactured by BMA) and then the amount of the DNA fragment of about 350 bp was measured by using Fluor Imager SI (manufactured by Molecular Dynamics).

(3) Analysis of expression amount of GFPP gene using RT-PCR

[0586] In order to amplify GFPP cDNA by PCR, a 27 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:34 and a 23 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:35 were prepared based on the CHO cell-derived GFPP cDNA sequence obtained in the item 2 of Reference Example 1.

[0587] Next, 20 µl of a reaction solution [1×Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 µmol/L of the synthetic DNA primers of SEQ ID NOs:34 and 35] containing 0.5 µl of the single-stranded cDNA prepared from each clone in the item 1(1) of Example 8 as the template was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 24 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After subjecting 10 µl of the PCR reaction solution to agarose electrophoresis, DNA fragments were stained with Cyber Green (manufactured by BMA) and then the amount of the DNA fragment of about 600 bp was measured by using Fluor Imager SI (manufactured by Molecular Dynamics).

(4) Analysis of expression amount of FX gene using RT-PCR

[0588] In order to amplify FX cDNA by PCR, a 28 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:36 and a 28 mer synthetic DNA primer having the nucleotide sequence shown by SEQ ID NO:3 were

prepared based on the CHO cell-derived FX cDNA sequence shown in the item 1 of Reference Example 1.

[0589] Next, 20 μ l of a reaction solution [$1 \times$ concentration Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 μ mol/L of the synthetic DNA primers of SEQ ID NO:36 and SEQ ID NO:37] containing 0.5 μ l of the single-stranded cDNA derived from each clone in the item 1(1) of Example 8 as the template was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 22 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After subjecting 10 μ l of the PCR reaction solution to agarose electrophoresis, DNA fragments were stained with Cyber Green (manufactured by BMA) and then the amount of the DNA fragment of about 300 bp was measured by using Fluor Imager SI (manufactured by Molecular Dynamics).

(5) Analysis of expression amount of FUT8 gene using RT-PCR

[0590] In order to amplify FUT8 cDNA by PCR, 20 μ l of a reaction solution [$1 \times$ Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 μ mol/L of the synthetic DNA primers of SEQ ID NOs:13 and 14] containing 0.5 μ l of the single-stranded cDNA derived from each clone in the item 1(1) of Example 8 as the template was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 20 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After subjecting 10 μ l of the PCR reaction solution to agarose electrophoresis, DNA fragments were stained with Cyber Green (manufactured by BMA) and then amount of the DNA fragment of about 600 bp was measured using Fluor Imager SI (manufactured by Molecular Dynamics).

(6) Analysis of expression amount of β -actin gene using RT-PCR

[0591] In order to amplify β -actin cDNA by PCR, 20 μ l of a reaction solution [$1 \times$ Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 μ mol/L of the synthetic DNA primers of SEQ ID NOs:15 and 16] containing 0.5 μ l of the single-stranded cDNA derived from each clone in the item 1(1) of Example 8 as the template was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 14 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After subjecting 10 μ l of the PCR reaction solution to agarose electrophoresis, DNA fragments were stained with Cyber Green (manufactured by BMA) and then the amount of the DNA fragment of about 800 bp was measured using Fluor Imager SI (manufactured by Molecular Dynamics).

(7) Expression amount of GMD, GFPP, FX and FUT8 genes in each clone

[0592] The amount of the PCR-amplified fragment of each gene in the clone 5-03 and the clone CHO/CCR4-LCA was calculated by dividing values of the amounts of PCR-amplified fragments derived from GMD, GFPP, FX and FUT8 cDNA in each clone measured in the items 1(2) to 1(6) of Example 8 by the value of the amount of PCR-amplified fragment derived from β -actin cDNA in each clone, and defining the amount of the PCR-amplified fragments in CHO/DG44 cell as 1. The results are shown in Table 5.

Table 5

| | GMD | GEPP | FX | FUT8 |
|---|-------|-------|-------|-------|
| Clone CHO/DG44 | 1 | 1 | 1 | 1 |
| Clone CHO/DG44 Anti-CCR4 antibody-producing cell Clone 5-03 | 1.107 | 0.793 | 1.093 | 0.901 |
| Derived from clone 5-03 LCA-resistant cell CHO/CCR4-LCA | 0.160 | 0.886 | 0.920 | 0.875 |

[0593] As shown in Table 5, the expression amount of GMD gene in the clone CHO/CCR4-LCA was decreased to about 1/10 in comparison with other clones. In this case, the test was independently carried out twice, and the average value was used.

2. Analysis using anti-CCR4 chimeric antibody-producing clone CHO/CCR4-LCA in which GMD gene was forced to express

(1) Construction of CHO cell-derived GMD gene expression vector pAGE249GMD

[0594] Based on the CHO cell-derived GMD cDNA sequence obtained in the item 1 of Reference Example 2, a 28 mer primer having the nucleotide sequence shown by SEQ ID NO:38 and a 29 mer primer having the nucleotide sequence shown by SEQ ID NO:39 were prepared. Next, 20 µl of a reaction solution [1 × concentration Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 µmol/L of the synthetic DNA primers of SEQ ID NOs:38 and 39] containing 0.5 µl of the CHO cell-derived single-stranded cDNA prepared in the item 1(1) of Example 8 as the template was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequently 8 cycles of heating at 94°C for 1 minute, 58°C for 1 minute and 72°C for 1 minute as one cycle, and then 22 cycles of heating at 94°C for 1 minute and 68°C as one cycle. After completion of the reaction, the PCR reaction solution was fractionated by agarose electrophoresis, and then a DNA fragment of about 600 bp was recovered. The recovered DNA fragment was connected to pT7Blue(R) vector (manufactured by Novagen) by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α (manufactured by Toyobo) was transformed using the obtained recombinant plasmid DNA to obtain a plasmid mt-C (Fig. 20).

[0595] Next, based on the CHO cell-derived GMD cDNA sequence obtained in the item 1 of Reference Example 2, a 45 mer primer having the nucleotide sequence shown by SEQ ID NO:40 and a 31 mer primer having the nucleotide sequence shown by SEQ ID NO:41 were prepared. Next, 20 µl of a reaction solution [1 × Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 µmol/L of the synthetic DNA primers of SEQ ID NOs:40 and 41] containing 0.5 µl of the CHO cell-derived single-stranded cDNA prepared in the item 1(1) of Example 8 as the template was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequently 8 cycles of heating at 94°C for 1 minute, 57°C for 1 minute and 72°C for 1 minute as one cycle, and then 22 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After completion of the reaction, the PCR reaction solution was fractionated by agarose electrophoresis, and then a DNA fragment of about 150 bp was recovered. The recovered DNA fragment was connected to pT7Blue(R) vector (manufactured by Novagen) by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α (manufactured by Toyobo) was transformed using the obtained recombinant plasmid DNA to obtain a plasmid ATG (Fig. 21).

[0596] Next, 3 µg of the plasmid CHO-GMD prepared in the item 1 of Reference Example 2 was allowed to react with a restriction enzyme *SacI* (manufactured by Takara Shuzo) at 37°C for 16 hours, a DNA was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and allowed to react with a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 900 bp was recovered. The plasmid mt-C (1.4 µg) was allowed to react with a restriction enzyme *SacI* (manufactured by Takara Shuzo) at 37°C for 16 hours, DNA was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and allowed to react with a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 3.1 kbp was recovered. The recovered DNA fragments were ligated by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α was transformed using the obtained recombinant plasmid DNA to obtain a plasmid WT-N(-) (Fig. 22).

[0597] Next, 2 µg of the plasmid WT-N(-) was allowed to react with a restriction enzyme *BamHI* (manufactured by Takara Shuzo) at 37°C for 16 hours, DNA was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and allowed to react with a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 1 kbp was recovered by using Gene Clean II Kit (manufactured by BIO 101) in accordance with the manufacture's instructions. The plasmid pBluescript SK(-) (3 µg; manufactured by Stratagene) was allowed to react with a restriction enzyme *BamHI* (manufactured by Takara Shuzo) at 37°C for 16 hours, DNA was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and allowed to react with a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 3 kbp was recovered. The respective recovered DNA fragments were ligated by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α was transformed using the obtained recombinant plasmid DNA to obtain a plasmid WT-N(-) in pBS (cf. Fig. 23).

[0598] Next, 2 µg of the plasmid WT-N(-) in pBS was allowed to react with a restriction enzyme *HindIII* (manufactured by Takara Shuzo) at 37°C for 16 hours, DNA was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and allowed to react with a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 4 kbp was

recovered. After 2 µg of the plasmid ATG was allowed to react with a restriction enzyme *HindIII* (manufactured by Takara Shuzo) at 37°C for 16 hours, DNA was recovered by carrying out phenol/chloroform extraction and ethanol precipitation and allowed to react with a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 150 bp was recovered. The respective recovered DNA fragments were ligated by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α was transformed using the obtained recombinant plasmid DNA to obtain a plasmid WT in pBS (Fig. 24).

[0599] Next, 2 µg of the plasmid pAGE249 was allowed to react with restriction enzymes *HindIII* and *BamHI* (both manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 6.5 kbp was recovered. The plasmid WT (2 µg) in pBS was allowed to react with restriction enzymes *HindIII* and *BamHI* (both manufactured by Takara Shuzo) at 37°C for 16 hours. A digested DNA was fractionated by agarose electrophoresis and then a DNA fragment of about 1.2 kbp was recovered. The respective recovered DNA fragments were ligated by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α was transformed using the obtained recombinant plasmid DNA to obtain a plasmid pAGE249GMD (Fig. 25).

(2) Stable expression of GMD gene in clone CHO/CCR4-LCA

[0600] The CHO cell-derived GMD gene expression vector pAGE249GMD (5 µg) which was made into linear form by digesting it with a restriction enzyme *FspI* (manufactured by NEW ENGLAND BIOLABS), was introduced into 1.6×10⁶ cells of the clone CHO/CCR4-LCA by electroporation [*Cytotechnology*, 3, 133 (1990)]. Then, the cells were suspended in 30 ml of IMDM-dFBS(10) medium [IMDM medium (manufactured by GIBCO BRL) supplemented with 10% of dFBS] comprising 200 nmol/L MTX (manufactured by SIGMA), and cultured in a 182 cm² flask (manufactured by Greiner) at 37°C for 24 hours in a 5% CO₂ incubator. After culturing, the medium was changed to IMDM-dFBS(10) medium containing 0.5 mg/ml hygromycin and 200 nmol/L MTX (manufactured by SIGMA), followed by culturing for 19 days to obtain colonies of hygromycin-resistant transformants.

[0601] In the same manner, the pAGE249 vector was introduced into the clone CHO/CCR4-LCA by the same method as above to obtain colonies of hygromycin-resistant transformants.

(3) Culturing of GMD gene-expressed clone CHO/CCR4-LCA and purification of antibody

[0602] Using IMDM-dFBS(10) medium comprising 200 nmol/L MTX (manufactured by SIGMA) and 0.5 mg/ml hygromycin, the GMD-expressing transformant cells obtained in the item 2(2) of Example 8 were cultured in a 182 cm² flask (manufactured by Greiner) at 37°C in a 5% CO₂ incubator. Several days thereafter, when the cell density reached confluent, the culture supernatant was discarded and the cells were washed with 25 ml of PBS buffer (manufactured by GIBCO BRL) and mixed with 35 ml of EXCELL301 medium (manufactured by JRH). After culturing at 37°C in a 5% CO₂ incubator for 7 days, the culture supernatant was recovered. An anti-CCR4 chimeric antibody was purified from the culture supernatant by using Prosep-A column (manufactured by Millipore) in accordance with the manufacture's instructions.

[0603] In the same manner, the pAGE249 vector-introduced transformant cells were cultured by the same method as above and then anti-CCR4 chimeric antibody was recovered and purified from the culture supernatant.

(4) Measurement of lectin resistance in transformed cells

[0604] The GMD gene-expressing transformant cells obtained in the item 2(2) of Example 8 were suspended in IMDM-dFBS(10) medium comprising 200 nmol/L MTX (manufactured by SIGMA) and 0.5 mg/ml hygromycin to give a density of 6×10⁴ cells/ml, and the suspension was dispensed at 50 µl/well into a 96 well culture plate (manufactured by Iwaki Glass). Next, a medium prepared by suspending LCA (*Lens culinaris* agglutinin: manufactured by Vector Laboratories) at concentrations of 0 mg/ml, 0.4 mg/ml, 1.6 mg/ml or 4 mg/ml in IMDM-dFBS(10) medium containing 200 nmol/L MTX (manufactured by SIGMA) and 0.5 mg/ml hygromycin was added to the plate at 50 µl/well, followed by culturing at 37°C for 96 hours in a 5% CO₂ incubator. After culturing, WST-1 (manufactured by Boehringer) was added at 10 µl/well and incubated at 37°C for 30 minutes in a 5% CO₂ incubator for color development, and then the absorbance at 450 nm and 595 nm (hereinafter referred to as "OD450" and "OD595", respectively) was measured by using Microplate Reader (manufactured by BIO-RAD). In the same manner, the pAGE249 vector-introduced transformant cells were measured by the same method as above. The above-mentioned test was carried out twice independently.

[0605] Fig. 26 shows the number of survived cells in each well by percentage when a value calculated by subtracting OD595 from OD450 measured in the above is used as the survived number of each cell group and the number of survived cells in each of the LCA-free wells is defined as 100%. As shown in Fig. 26, decrease in the LCA-resistance

was observed in the GMD-expressed clone CHO/CCR4-LCA, and the survival ratio was about 40% in the presence of 0.2 mg/ml LCA and the survival ratio was about 20% in the presence of 0.8 mg/ml LCA. On the other hand, in the pAGE249 vector-introduced strain CHO/CCR4-LCA, the survival ratio was 100% in the presence of 0.2 mg/ml LCA and the survival ratio was about 80% even in the presence of 0.8 mg/ml LCA. Based on the above results, it was suggested that expression amount of GMD gene in the clone CHO/CCR4-LCA was decreased and, as a result, the resistance against LCA was obtained.

(5) ADCC activity of anti-CCR4 chimeric antibody obtained from GMD-expressed clone CHO/CCR4-LCA

[0606] ADCC activity of the purified anti-CCR4 chimeric antibody obtained in the item 2(3) of Example 8 was measured in accordance with the following method.

i) Preparation of target cell suspension

[0607] CCR4/EL4 cell described in W001/64754 was prepared at 1×10^6 cells and 3.7 MBq equivalent of a radioactive substance $\text{Na}_2^{51}\text{CrO}_4$ was added to thereto, followed by reaction at 37°C for 90 minutes to thereby label the cells with a radioisotope. After the reaction, the cells were washed three times by suspension in the RPMI1640-FBS(10) medium and subsequent centrifugation, resuspended in the medium and then incubated at 4°C for 30 minutes on ice for spontaneous dissociation of the radioactive substance. After centrifugation, the cells were adjusted to 2.0×10^5 cells/ml by adding 5 ml of the RPMI1640-FBS(10) medium and used as a target cell suspension.

ii) Preparation of effector cell suspension

[0608] From a healthy donor, 50 ml of venous blood was collected and gently mixed with 0.5 ml of heparin sodium (manufactured by Takeda Pharmaceutical). Using Lymphoprep (manufactured by Nycomed Pharma AS), the mixture was centrifuged in accordance with the manufacture's instructions to separate a mononuclear cell layer. The cells were washed three times by centrifuging with the RPMI1640-FBS(10) medium and then resuspended in the medium to give a density of 2.5×10^6 cells/ml and used as a effector cell suspension.

iii) Measurement of ADCC activity

[0609] The target cell suspension prepared in the above i) was dispensed at 50 μl (1×10^4 cells/well) into each well of a 96 well U-bottom plate (manufactured by Falcon). Next, 100 μl of the effector cell suspension prepared in the above ii) was added thereto (2×10^5 cells/well, ratio of the effector cells to the target cells was 25 : 1). Each of various anti-CCR4 chimeric antibodies was further added thereto to give a final concentration of 0.0025 to 2.5 $\mu\text{g/ml}$, followed by reaction at 37°C for 4 hours. After the reaction, the plate was centrifuged and the amount of ^{51}Cr in the supernatant was measured with a γ -counter. The amount of the spontaneously dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the effector cell suspension and antibody solution, and measuring the amount of ^{51}Cr in the supernatant. The amount of the total dissociated ^{51}Cr was calculated by carrying out the same procedure using the medium alone instead of the antibody solution and adding 1 mol/L hydrochloric acid instead of the effector cell suspension and measuring the amount of ^{51}Cr in the supernatant. The ADCC activity was calculated based on the above equation (1).

[0610] Results of the measurement of ADCC activity are shown in Fig. 27. As shown in Fig. 27, ADCC activity of the purified anti-CCR4 chimeric antibody obtained from the GMD-expressed clone CHO/CCR4-LCA was decreased to a similar degree to that of the KM3060 produced by the normal CHO cell-derived antibody-producing clone obtained in Example 4. On the other hand, ADCC activity of the purified anti-CCR4 chimeric antibody obtained from the pAGE249 vector-introduced clone CHO/CCR4-LCA showed a similar degree of ADCC activity to that of the purified anti-CCR4 chimeric antibody obtained from the clone CHO/CCR4-LCA. Based on the above results, it was suggested that expression amount of GMD gene in the clone CHO/CCR4-LCA is decreased and, as a result, an antibody having high ADCC activity can be produced.

(6) Sugar chain analysis of anti-CCR4 chimeric antibody derived from GMD-expressed clone CHO/CCR4-LCA

[0611] Sugar chains binding to the purified anti-CCR4 chimeric antibody obtained in the item 2(3) of Example 8 were analyzed in accordance with the method shown in the item 1 or Example 3, and the analyzed results are shown in Fig. 28. In comparison with the purified anti-CCR4 chimeric antibody prepared from the clone CHO/CCR4-LCA in Example 7, the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the purified anti-CCR4 chimeric antibody derived from the GMD gene-expressed clone

CHO/CCR4-LCA was decreased to 9% when calculated from the peak area. Thus, it was shown that the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the antibody produced by the cell is decreased to similar level of the antibody produced by the clone 5-03.

Example 9

Preparation of anti-fibroblast growth factor-8 human chimeric antibody

1. Preparation of cells stably producing anti-fibroblast growth factor-8 human chimeric antibody

[0612] Using a tandem type expression vector pKANTEX134 of an anti-fibroblast growth factor-8 (hereinafter referred to as "FGF-8") human chimeric antibody described in Reference Example 3, cells stably producing the anti-FGF-8 human chimeric antibody (hereinafter referred to as "anti-FGF-8 chimeric antibody") was prepared as follows.

(1) Preparation of producing cell using rat myeloma YB2/0 cell

[0613] After introducing 10 μ g of the anti-FGF-8 chimeric antibody expression vector pKANTEX1334 into 4×10^6 cells of rat myeloma YB2/0 cell (ATCC CRL 1662) by electroporation [*Cytotechnology*, 3, 133 (1990)], the cells were suspended in 40 ml of Hybridoma-SFM-FBS(5) [Hybridoma-SFM medium (manufactured by Invitrogen) containing 5% FBS (manufactured by PAA Laboratories)] and dispensed at 200 μ l/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 0.5 mg/ml, followed by culturing for 1 to 2 weeks. Culture supernatants were recovered from wells in which colonies of transformants showing G418 resistance were formed and their growth was confirmed, and antigen-binding activity of the anti-FGF-8 chimeric antibody in the supernatants was measured by the ELISA described in the item 2 of Example 9.

[0614] Regarding the transformants in wells in which production of the anti-FGF-8 chimeric antibody was found in the culture supernatants, in order to increase the antibody production amount by using a *dhfr* gene amplification system, each of them was suspended to give a density of 1 to 2×10^5 cells/ml in the Hybridoma-SFM-FBS(5) medium containing 0.5 mg/ml G418 and 50 nmol/l DHFR inhibitor MTX (manufactured by SIGMA) and dispensed at 1 ml into each well of a 24 well plate (manufactured by Greiner). After culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nmol/l MTX resistance were induced. Antigen-binding activity of the anti-FGF-8 chimeric antibody in culture supernatants in wells where growth of transformants was observed was measured by the ELISA described in the item 2 of Example 9.

[0615] Regarding the transformants in wells in which production of the anti-FGF-8 chimeric antibody was found in culture supernatants, the MTX concentration was increased by a method similar to the above to thereby finally obtain a transformant 5-D capable of growing in the Hybridoma-SFM-FBS(5) medium containing 0.5 mg/ml G418 and 200 nmol/l MTX and also highly producing the anti-FGF-8 chimeric antibody. The resulting transformant was subjected to cloning by limiting dilution, and the resulting transformant cell clone was named 5-D-10.

(2) Preparation of producing cell using CHO/DG44 cell

[0616] In accordance with the method described in Example 4, the anti-FGF-8 chimeric antibody expression plasmid pKANTEX1334 was introduced into CHO/DG44 cell and gene amplification was carried out by using the drug MTX to obtain a transformant highly producing the anti-FGF-8 chimeric antibody. The antibody expression amount was measured using the ELISA described in the item 2 of Example 9. The resulting transformant was cloned twice by limiting dilution, and the resulting transformant cell clone was named 7-D-1-5.

2. Binding activity of antibody to FGF-8 partial peptide (ELISA)

[0617] Compound 2 (SEQ ID NO:21) was selected as a human FGF-8 peptide with which the anti-FGF-8 chimeric antibody can react. For the activity measurement by the ELISA, a conjugate with BSA (manufactured by Nacalai Tesque) was prepared by the following method and used as the antigen. That is, 100 ml of a 25 mg/ml SMCC [4-(*N*-maleimidomethyl)cyclohexane-1-carboxylic acid *N*-hydroxysuccinimide ester] (manufactured by SIGMA)-DMSO solution was added dropwise to 900 ml of a PBS solution containing 10 mg of BSA under stirring, followed by slowly stirred for 30 minutes. To a gel filtration column such as NAP-10 column or the like which had been equilibrated with 25 ml of PBS, 1 ml of the reaction solution was applied, and the eluate eluted with 1.5 ml of PBS was used as a BSA-SMCC solution (BSA concentration was calculated from A₂₈₀ measurement). Next, 250 ml of PBS was added to 0.5 mg of Compound

2, 250 ml of DMF was added thereto and completely dissolved, and then the above BSA-SMCC solution (1.25 mg as BSA) was added thereto under stirring, followed by slow stirring for 3 hours. The reaction solution was dialyzed against PBS at 4°C overnight, sodium azide was added thereto to give a final concentration of 0.05% and then filtered through a 0.22 µm filter and used as a BSA-compound 2 solution.

[0618] The conjugate prepared in the above was dispensed at 1 µg/ml and 50 µl/well into a 96 well plate for ELISA (manufactured by Greiner) and adhered thereto by allowing it to stand at 4°C overnight. After washing with PBS, 1% BSA-PBS was added at 100 µl/well and allowed to react at room temperature for 1 hour to block the remaining active groups. After washing each well with Tween-PBS, culture supernatant of a transformant or a purified antibody was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing of each well with Tween-PBS, a peroxidase-labeled goat anti-human IgG (γ) antibody solution (manufactured by American Qualex) diluted 3,000-fold with 1% BSA-PBS was added as a secondary antibody solution at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was added at 50 µl/well to develop color, and the reaction was stopped 10 minutes thereafter by adding 5% SDS solution at 50 µl/well. Thereafter, OD415 was measured.

3. Purification of anti-FGF-8 chimeric antibody

(1) Culturing of YB2/0 cell-derived producing cell and purification of antibody

[0619] The anti-FGF-8 chimeric antibody-expressing transformant 5-D obtained in the item 1(1) of Example 9 was cultured in Hybridoma-SFM (manufactured by Invitrogen) medium containing 200 nmol/l of MTX and 5% Daigo's GF21 (manufactured by Wako Pure Chemical Industries) in a 182 cm² flask (manufactured by Greiner) at 37°C in a 5% CO₂ incubator. After culturing for 8 to 10 days, the anti-FGF-8 chimeric antibody was purified from the culture supernatant recovered by using Prosep-A (manufactured by Millipore) column and in accordance with the attached manufacture's instructions. The purified anti-FGF-8 chimeric antibody was named YB2/0-FGF8 chimeric antibody.

(2) Culturing of CHO-DG44 cell-derived antibody-producing cells and purification of antibody

[0620] The anti-FGF-8 chimeric antibody-producing transformant cell clone 7-D-1-5 obtained in the item 1(2) of Example 9 was cultured in the IMDM-dFBS(10) medium in a 182 cm² flask (manufactured by Greiner) at 37°C in a 5% CO₂ incubator. At the stage where the cell density reached confluent several days thereafter, the culture supernatant was discarded, the cells were washed with 25 ml of PBS buffer and then 35 ml of EXCELL301 medium (manufactured by JRH) was added thereto. After the culturing for 7 days at 37°C in a 5% CO₂ incubator, the culture supernatant was recovered. The anti-FGF-8 chimeric antibody was purified from the culture supernatant by using Prosep-A (manufactured by Millipore) column and in accordance with the manufacture's instructions. The purified anti-FGF-8 chimeric antibody was named CHO-FGF8 chimeric antibody.

[0621] When the binding activity of the YB2/0-FGF8 chimeric antibody and CHO-FGF8 chimeric antibody to FGF-8 was measured by the ELISA described in the item 2 of Example 9, they showed similar binding activity.

4. Analysis of purified anti-FGF-8 chimeric antibody

[0622] Each 4 µg of the two anti-FGF-8 chimeric antibodies produced by respective animal cells and purified in the item 3 of Example 9 was subjected to SDS-PAGE according to a known method [*Nature*, 227, 680 (1970)] and the molecular weight and purity were analyzed. In each of the purified anti-FGF-8 chimeric antibodies, a single band of about 150 Kd in molecular weight was found under non-reducing conditions and two bands of about 50 Kd and about 25 Kd were found under reducing conditions. These molecular weights almost coincided with the molecular weights deduced from the cDNA nucleotide sequences of the antibody H chain and L chain (H chain: about 49 Kd, L chain: about 23 Kd, whole molecule: about 144 Kd), and also coincided with the reports showing that the IgG type antibody shows a molecular weight of about 150 Kd under non-reducing conditions and is degraded into H chain having a molecular weight of about 50 Kd and L chain having a molecular weight of about 25 Kd under reducing conditions due to cleavage of the intramolecular S-S bond (*Antibodies*, Chapter 14 (1988); *Monoclonal Antibodies*). Thus it was confirmed that the anti-FGF-8 chimeric antibodies were expressed and purified as antibody molecules having correct structures.

5. Sugar chain analysis of purified anti-FGF-8 chimeric antibodies

[0623] Sugar chain analysis of the YB2/0-FGF8 chimeric antibody which is YB2/0 cell-derived anti-FGF-8 chimeric antibody and anti-FGF-8 chimeric antibody CHO-FGF8 chimeric antibody which is CHO/DG44 cell-derived prepared

in the item 4 of Example 9 was carried out in accordance with the method described in the item 1 of Example 3. As a result, the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond were 58% in the YB2/0-FGF8 chimeric antibody and 13% in the CHO-FGF8 chimeric antibody. Hereinafter, these samples are called anti-FGF-8 chimeric antibody (58%) and anti-FGF-8 chimeric antibody (13%).

Example 10

Preparation of soluble human Fc γ RIIIa protein

1. Construction of a soluble human Fc γ RIIIa protein expression vector

(1) Preparation of human peripheral blood monocyte cDNA

[0624] From a healthy donor, 30 ml of vein blood was collected, gently mixed with 0.5 ml of heparin sodium (manufactured by Shimizu Pharmaceutical) and then mixed with 30 ml of physiological saline (manufactured by Otsuka Pharmaceutical). After the mixing, 10 ml of each mixture was gently overlaid on 4 ml of Lymphoprep (manufactured by NYCOMED PHARMA AS) and centrifuged at 2,000 rpm for 30 minutes at room temperature. The separated monocyte fractions in respective centrifugation tubes were combined and suspended in 30 ml of RPMI1640-FBS(10). After centrifugation at room temperature and at 1,200 rpm for 15 minutes, the supernatant was discarded and the cell were suspended in 20 ml of RPMI1640-FBS(10). This washing operation was repeated twice and then 2×10^6 cells/ml of peripheral blood monocyte suspension was prepared using RPMI1640-FBS(10).

[0625] After 5 ml of the resulting peripheral blood monocyte suspension was centrifuged at room temperature and at 800 rpm for 5 minutes in 5 ml of PBS, the supernatant was discarded and the residue was suspended in 5 mL of PBS. After centrifugation at room temperature and at 800 rpm for 5 minutes, the supernatant was discarded and total RNA was extracted by QIAamp RNA Blood Mini Kit (manufactured by QIAGEN) and in accordance with the manufacture's instructions.

[0626] A single-stranded cDNA was synthesized by reverse transcription reaction to 2 μ g of the resulting total RNA, in a series of 40 μ l containing oligo(dT) as primers using SUPERSCRIP™ Preamplification System for First Strand cDNA Synthesis (manufactured by Life Technologies) according to the manufacture's instructions.

(2) Preparation method of cDNA encoding human Fc γ RIIIa protein

[0627] A cDNA encoding a human Fc γ RIIIa protein (hereinafter referred to as "hFc γ RIIIa") was prepared as follows.

[0628] First, a specific forward primer containing a translation initiation codon (represented by SEQ ID NO:22) and a specific reverse primer containing a translation termination codon (represented by SEQ ID NO:26) were designed from the nucleotide sequence of hFc γ RIIIa cDNA [*J. Exp. Med.*, 170, 481 (1989)].

[0629] Next, using a DNA polymerase ExTaq (manufactured by Takara Shuzo), 50 μ l of a reaction solution [1 \times concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 1 μ mol/l of the above gene-specific primers (SEQ ID NOs:22 and 26)] containing 5 μ l of 20-fold diluted solution of the human peripheral blood monocyte-derived cDNA solution prepared in the item 1(1) of Example 10 was prepared, and PCR was carried out. The PCR was carried out by 35 cycles of a reaction at 94°C for 30 seconds, at 56°C for 30 seconds and at 72°C for 60 seconds as one cycle.

[0630] After the PCR, the reaction solution was purified by using QIAquick PCR Purification Kit (manufactured by QIAGEN) and dissolved in 20 μ l of sterile water. The products were digested with restriction enzymes *Eco*RI (manufactured by Takara Shuzo) and *Bam*HI (manufactured by Takara Shuzo) and subjected to 0.8% agarose gel electrophoresis to recover about 800 bp of a specific amplification fragment.

[0631] On the other hand, 2.5 μ g of a plasmid pBluescript II SK(-) (manufactured by Stratagene) was digested with restriction enzymes *Eco*RI (manufactured by Takara Shuzo) and *Bam*HI (manufactured by Takara Shuzo), and digested products were subjected to 0.8% agarose gel electrophoresis to recover a fragment of about 2.9 kbp.

[0632] The human peripheral blood monocyte cDNA-derived amplification fragment and plasmid pBluescript II SK(-)-derived fragment obtained in the above were ligated by using DNA Ligation Kit Ver. 2.0 (manufactured by Takara Shuzo). The strain *Escherichia coli* DH5 α (manufactured by TOYOBO) was transformed by using the reaction solution, and a plasmid DNA was isolated from each of the resulting ampicillin-resistant colonies according to a known method.

[0633] A nucleotide sequence of the cDNA inserted into each plasmid was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction Kit (manufactured by Parkin Elmer) according to the manufacture's instructions. It was confirmed that all of the inserted cDNAs whose sequences were determined by this method encodes a complete ORF sequence of the cDNA encoding hFc γ RIIIa. A

plasmid DNA containing absolutely no reading error of bases in the sequence accompanied by PCR was selected from them. Hereinafter, the plasmid is called pBSFcγRIIIa5-3.

[0634] The thus determined full length cDNA sequence for hFcγRIIIa is represented by SEQ ID NO:27, and its corresponding amino acid sequence is shown in by SEQ ID NO:28.

(3) Preparation of a cDNA encoding soluble hFcγRIIIa

[0635] A cDNA encoding soluble hFcγRIIIa (hereinafter referred to as "shFcγRIIIa") having the extracellular region of hFcγRIIIa (positions 1 to 193 in SEQ ID NO:28) and a His-tag sequence at the C-terminal was constructed as follows.

[0636] First, a primer FcγR3-1 (represented by SEQ ID NO:29) specific for the extracellular region was designed from the nucleotide sequence of hFcγRIIIa cDNA (represented by SEQ ID NO:27).

[0637] Next, using a DNA polymerase ExTaq (manufactured by Takara Shuzo), 50 μl of a reaction solution [1× concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs, 1 μmol/l of the primer FcγR3-1, 1 μmol/l of the primer M13M4 (manufactured by Takara Shuzo)] containing 5 ng of the plasmid pBSFcγRIIIa5-3 prepared in the item 1(2) of Example 10 was prepared, and PCR was carried out. The PCR was carried out by 35 cycles of a reaction at 94°C for 30 seconds, at 56°C for 30 seconds and at 72°C for 60 seconds as one cycle.

[0638] After the PCR, the reaction solution was purified by using QIAquick PCR Purification Kit (manufactured by QIAGEN) and dissolved in 20 μl of sterile water. The products were digested with restriction enzymes *Pst*I (manufactured by Takara Shuzo) and *Bam*HI (manufactured by Takara Shuzo) and subjected to 0.8% agarose gel electrophoresis to recover about 110 bp of a specific amplification fragment.

[0639] On the other hand, 2.5 μg of the plasmid pBSFcγRIIIa5-3 was digested with restriction enzymes *Pst*I (manufactured by Takara Shuzo) and *Bam*HI (manufactured by Takara Shuzo), and the digested products were subjected to 0.8% agarose gel electrophoresis to recover a fragment of about 3.5 kbp.

[0640] The hFcγRIIIa cDNA-derived amplification fragment and plasmid pBSFcγRIIIa5-3-derived fragment obtained in the above were ligated by using DNA Ligation Kit Ver. 2.0 (manufactured by Takara Shuzo). The strain *Escherichia coli* DH5α (manufactured by TOYOBO) was transformed by using the reaction solution, and a plasmid DNA was isolated from each of the resulting ampicillin-resistant colonies according to a known method.

[0641] A nucleotide sequence of the cDNA inserted into each plasmid was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction Kit (manufactured by Parkin Elmer) according to the manufacture's instructions. It was confirmed that all of the inserted cDNAs whose sequences were determined by this method encodes a complete ORF sequence of the cDNA encoding shFcγRIIIa of interest. A plasmid DNA containing absolutely no reading error of bases in the sequence accompanied by PCR was selected from them. Hereinafter, this plasmid is named pBSFcγRIIIa+His3. The thus determined full length cDNA sequence for shFcγRIIIa is represented by SEQ ID NO:30, and its corresponding amino acid sequence is represented by SEQ ID NO:31.

(4) Construction of shFcγRIIIa expression vector

[0642] The shFcγRIIIa expression vector was constructed as follows.

[0643] After 3.0 μg of the plasmid pBSFcγRIIIa+His3 obtained in the item 1(3) of Example 10 was digested with restriction enzymes *Eco*RI (manufactured by Takara Shuzo) and *Bam*HI (manufactured by Takara Shuzo), the digested products were subjected to 0.8% agarose gel electrophoresis to recover a fragment of about 620 bp.

[0644] Separately, 2.0 μg of the plasmid pKANTEX93 described in WO97/10354 was digested with restriction enzymes *Eco*RI (manufactured by Takara Shuzo) and *Bam*HI (manufactured by Takara Shuzo), and the digested products were subjected to 0.8% agarose gel electrophoresis to recover a fragment of about 10.7 kbp.

[0645] The DNA fragment containing shFcγRIIIa cDNA and the plasmid pKANTEX93-derived fragment obtained in the above were ligated by using DNA Ligation Kit Ver. 2.0 (manufactured by Takara Shuzo). The strain *Escherichia coli* DH5α (manufactured by TOYOBO) was transformed by using the reaction solution, and a plasmid DNA was isolated from each of the resulting ampicillin-resistant colonies according to a known method.

[0646] A nucleotide sequence of the cDNA inserted into each plasmid was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction Kit (manufactured by Parkin Elmer) in accordance with the manual attached thereto. It was confirmed that all of the plasmids whose sequences were determined by this method encodes the cDNA of interest encoding shFcγRIIIa. Hereinafter, the obtained expression vector was named pKANTEXFcγRIIIa-His.

2. Preparation of cell stably producing shFcγRIIIa

[0647] Cells stably producing shFcγRIIIa were prepared by introducing the shFcγRIIIa expression vector pKAN-

TEXFcyRIIIa-His constructed in the item 1 of Example 10 into rat myeloma YB2/0 cell [ATCC CRL-1662, *J. Cell. Biol.*, 93, 576 (1982)],

[0648] pKANTEXFcyRIIIa-His was digested with a restriction enzyme *AatII* to obtain a linear fragment, 10 µg thereof was introduced into 4×10^6 cells by electroporation [*Cytotechnology*, 3, 133 (1990)], and the resulting cells were suspended in 40 ml of Hybridoma-SFM-FBS(10) [Hybridoma-SFM medium (manufactured by Life Technologie) containing 10% FBS] and dispensed at 200 µl/well into a 96 well culture plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added to give a concentration of 1.0 mg/ml, followed by culturing for 1 to 2 weeks. Culture supernatants were recovered from wells in which colonies of transformants showing G418 resistance were formed and their growth was confirmed, and expression amount of shFcyRIIIa in the supernatants was measured by the ELISA described in the item 3 of Example 10.

[0649] Regarding the transformants in wells in which expression of the shFcyRIIIa was confirmed in the culture supernatants, in order to increase the antibody production by using a *dhfr* gene amplification system, each of them was suspended to give a density of 1 to 2×10^5 cells/ml in the Hybridoma-SFM-FBS(10) medium containing 1.0 mg/ml G418 and 50 nmol/l DHFR inhibitor MTX (manufactured by SIGMA) and dispensed at 2 ml into each well of a 24 well plate (manufactured by Greiner). After culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator, transformants showing 50 nmol/l MTX resistance were induced. Expression level of shFcyRIIIa in culture supernatants in wells where growth of transformants was observed was measured by the ELISA described in The item 3 of Example 10. Regarding the transformants in wells in which expression of the shFcyRIIIa was found in culture supernatants, the MTX concentration was increased to 100 nmol/l and then to 200 nmol/l sequentially by a method similar to the above to thereby finally obtain a transformant capable of growing in the Hybridoma-SFM-FBS(10) medium containing 1.0 mg/ml G418 and 200 nmol/l MTX and also of highly producing shFcyRIIIa. The resulting transformant was cloned twice by limiting dilution. The obtained transformant was named clone KC1107.

3. Detection of shFcyRIIIa (ELISA)

[0650] shFcyRIIIa in culture supernatant or purified shFcyRIIIa was detected or determined by the ELISA shown below.

[0651] A solution of a mouse antibody against His-tag, Tetra-His Antibody (manufactured by QIAGEN), adjusted to 5 µg/ml with PBS was dispensed at 50 µl/well into each well of a 96 well plate for ELISA (manufactured by Greiner) and allowed to react at 4°C for 12 hours or more. After the reaction, 1% BSA-PBS was added at 100 µl/well and allowed to react at room temperature for 1 hour to block the remaining active groups. After 1% BSA-PBS was discarded, culture supernatant of the transformant or each of various dilution solutions of purified shFcyRIIIa was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing of each well with Tween-PBS, a biotin-labeled mouse anti-human CD16 antibody solution (manufactured by PharMingen) diluted 50-fold with 1% BSA-PBS was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, a peroxidase-labeled Avidin D solution (manufactured by Vector) diluted 4,000-fold with 1% BSA-PBS was added at 50 µl/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was added at 50 µl/well to develop color, and OD415 was measured.

4. Purification of shFcyRIIIa

[0652] The shFcyRIIIa-producing transformant cell clone KC1107 obtained in the item 2 of Example 10 was suspended in Hybridoma-SFM-GF(5) [Hybridoma-SFM medium (manufactured by Life Technologie) containing 5% Daigo's GF21 (manufactured by Wako Pure Chemical Industries)] to give a density of 3×10^5 cells/ml and dispensed at 50 ml into 182 cm² flasks (manufactured by Greiner). After culturing at 37°C for 4 days in a 5% CO₂ incubator, the culture supernatants were recovered. shFcyRIIIa was purified from the culture supernatants by using Ni-NTA agarose (manufactured by QIAGEN) column according to the manufacture's instructions.

5. Analysis of purified shFcyRIIIa

[0653] A concentration of purified shFcyRIIIa obtained in the item 4 of Example 10 was calculated by amino acid composition analysis as follows. A part of purified shFcyRIIIa was suspended in 6 mol/l hydrochloric acid-1% phenol solution, and hydrolysed in a gas phase at 110°C for 20 hours. Work Station manufactured by Waters was used for the hydrolysis. Amino acids after the hydrolysis were converted into PTC-amino acid derivatives in accordance with the method of Bidlingmeyer *et al.* [*J. Chromatogr.*, 336, 93 (1984)] and analyzed by using PicoTag Amino Acid Analyzer (manufactured by Waters).

[0654] Next, about 0.5 µg of purified shFcyRIIIa was subjected to SDS-PAGE under reducing conditions according

to a known method [*Nature*, 227, 680 (1970)] to analyze its molecular weight and purity. The results are shown in Fig. 6. As shown in Fig. 29, a broad band of 36 to 38 Kd in molecular weight was detected in purified shFcγRIIIa. Since it is known that five sites to which *N*-glycoside-linked sugar chains can be bound are present in the extracellular region of hFcγRIIIa [*J. Exp. Med.*, 170, 481 (1989)], it was considered that the broad molecular weight distribution of purified shFcγRIIIa is based on the irregularity of sugar chain addition. On the other hand, when the N-terminal amino acid sequence of purified shFcγRIIIa was analyzed by automatic Edman degradation using a protein sequencer PPSQ-10 (manufactured by Shimadzu), a sequence expected from the cDNA of shFcγRIIIa was obtained, so that it was confirmed that shFcγRIIIa of interest was purified.

Example 11

Evaluation of binding activity of various chimeric antibodies to shFcγRIIIa

1. Evaluation of shFcγRIIIa-binding activity of anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond

[0655] The shFcγRIIIa-binding activity of the anti-GD3 chimeric antibody (45%) and anti-GD3 chimeric antibody (7%) described in the item 1 of Example 3 which are two anti-GD3 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was measured by ELISA as follows.

[0656] According to the method described in the item 3 of Example 1, GD3 was immobilized at 100 pmol/well on a 96 well plate for ELISA (manufactured by Greiner). The 1% BSA-PBS was added at 100 μ l/well and allowed to react at room temperature for 1 hour to block the remaining active groups. After washing each well with Tween-PBS, a solution of each anti-GD3 chimeric antibody diluted with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing of each well with Tween-PBS, an shFcγRIIIa solution prepared by diluting it to 2.3 μ g/ml with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, a solution of a mouse antibody against His-tag, Tetra-His Antibody (manufactured by QIAGEN), adjusted to 1 μ g/ml with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, a peroxidase-labeled goat anti-mouse IgG1 antibody solution (manufactured by ZYMED) diluted 200-fold with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was added at 50 μ l/well to develop color, and OD415 was measured. In addition, it was confirmed that there is no difference in the amount of the anti-GD3 chimeric antibodies bound to the plate by adding each of the anti-GD3 chimeric antibodies to another plate and carrying out the ELISA described in item 3 of Example 1. The results of the measurement of the binding activity of the various anti-GD3 chimeric antibodies for shFcγRIIIa are shown in Fig. 30. As shown in Fig. 30, regarding the binding activity of the anti-GD3 chimeric antibodies to shFcγRIIIa, the anti-GD3 chimeric antibody (45%) having a high ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond had 10 times or more higher activity.

2. Evaluation of shFcγRIIIa-binding activity of anti-FGF-8 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond

[0657] The shFcγRIIIa-binding activity of the anti-FGF-8 chimeric antibody (58%) and anti-FGF-8 chimeric antibody (13%) described in the item 5 of Example 9 which were as two anti-FGF-8 chimeric antibodies having different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond was measured by ELISA as follows.

[0658] The human FGF-8 peptide conjugate prepared in the item 2 of Example 9 at a concentration of 1.0 μ g/ml was dispensed at 50 μ l/well into a 96-well plate for ELISA (manufactured by Greiner) and adhered thereto by allowing it to stand at 4°C overnight. After washing with PBS, 1% BSA-PBS was added at 100 μ l/well and allowed to react at room temperature for 1 hour to block the remaining active groups. After washing each well with Tween-PBS, a solution of each anti-FGF-8 chimeric antibody diluted with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing of each well with Tween-PBS, an shFcγRIIIa solution prepared by diluting it to 3.0 μ g/ml with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, a solution of a mouse antibody against His-tag, Tetra-His Antibody (manufactured by QIAGEN), adjusted to 1 μ g/ml with 1% BSA-PBS, was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, a peroxidase-labeled goat anti-mouse IgG1 antibody solution (manufactured by ZYMED) diluted 200-fold with 1%

BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was added at 50 μ l/well to develop color, and OD415 was measured. In addition, by adding each of the anti-FGF-8 chimeric antibodies to another plate and carrying out the ELISA described in the item 2 of Example 9, it was confirmed that there is no difference in the amount of the various anti-FGF-8 chimeric antibodies bound to the plate. The results of the measurement of the binding activity of the various anti-FGF-8 chimeric antibodies for shFc γ R1IIa are shown in Fig. 31. As shown in Fig. 31, regarding the binding activity of the anti-FGF8 chimeric antibodies to shFc γ R1IIa, the anti-FGF-8 chimeric antibody (58%) having a high ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond had 100 times or more higher activity.

3. Evaluation of shFc γ R1IIa-binding activity of anti-CCR4 chimeric antibodies having a different ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond

[0659] The shFc γ R1IIa-binding activity of the seven anti-CCR4 chimeric antibodies described in the item 5 of Example 4, having a different ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond, was measured by ELISA as follows.

[0660] The human CCR4 extracellular peptide conjugate prepared in the item 2 of Example 4 at a concentration of 1.0 μ g/ml was dispensed at 50 μ l/well into a 96 well plate for ELISA (manufactured by Greiner) and adhered thereto by allowing it to stand at 4°C overnight. After washing with PBS, 1% BSA-PBS was added at 100 μ l/well and allowed to react at room temperature for 1 hour to block the remaining active groups. After washing each well with Tween-PBS, a solution of each anti-CCR4 chimeric antibody diluted with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing of each well with Tween-PBS, an shFc γ R1IIa solution prepared by diluting it to 3.0 μ g/ml with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, a solution of a mouse antibody against His-tag, Tetra-His Antibody (manufactured by QIAGEN), adjusted to 1 μ g/ml with 1% BSA-PBS, was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, a peroxidase-labeled goat anti-mouse IgG1 antibody solution (manufactured by ZYMED) diluted 200-fold with 1% BSA-PBS was added at 50 μ l/well and allowed to react at room temperature for 1 hour. After the reaction and subsequent washing with Tween-PBS, the ABTS substrate solution was added at 50 μ l/well to develop color, and OD415 was measured. In addition, it was confirmed that there is no difference in the amount of the anti-CCR4 chimeric antibodies bound to the plate by adding each of the anti-CCR4 chimeric antibodies to another plate and carrying out the ELISA described in the item 2 of Example 4. The results of the measurement of the binding activity of the various anti-CCR4 chimeric antibodies to shFc γ R1IIa are shown in Fig. 32A. As shown in Fig. 32A, the binding activity of the anti-CCR4 chimeric antibodies to shFc γ R1IIa increased in proportion to the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond. Fig. 32B shows a plotted graph on a relationship between the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond, at antibody concentrations of 4 μ g/ml and 40 μ g/ml (the abscissa) and the shFc γ R1IIa-binding activity (the ordinate). As shown in Fig. 32B, the shFc γ R1IIa-binding activity was hardly detected in the anti-CCR4 chimeric antibody (8%), anti-CCR4 chimeric antibody (9%) and anti-CCR4 chimeric antibody (18%), as antibodies having 20% or less of the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond.

[0661] The above results clearly show that antibodies having a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond have higher shFc γ R1IIa-binding activity than antibodies having a α 1,6-fucose sugar chain. In addition, since antibodies having a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond has higher ADCC activity than antibodies having a α 1,6-fucose sugar chain as shown in Examples 3 and 4, it was strongly suggested that the high ADCC activity of antibodies a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is based on the high shFc γ R1IIa-binding activity. As shown in Fig. 32B, the proportion between Fc γ R1IIa binding activity and the ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is found. Therefore, Fc γ R1IIa-binding activity can be measured and the ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond can be determined by constructing such a calibration curve in advance. By using the above method, the cytotoxic activity can be prospected easily without determining the cytotoxic activity of the antibody composition.

Example 12

Evaluation of antibody produced by lectin-resistant CHO/DG44 cells to shFcγRIIIa

[0662] Binding activities of the anti-CCR4 chimeric antibody produced by the lectin-resistant clone CHO/CCR4-LCA purified in the item 3 of Example 7 [hereinafter referred to as "anti-CCR4 chimeric antibody (48%)"], the anti-CCR4 chimeric antibody KM2760-1 produced by a YB2/0 cell-derived antibody-producing clone purified in the item 3 of Example 4 [anti-CCR4 chimeric antibody (87%)] and the anti-CCR4 chimeric antibody KM3060 produced by a CHO/DG44 cell-derived antibody-producing clone 5-03 [anti-CCR4 chimeric antibody (8%)] to shFcγRIIIa were measured according to the method described in the item 3 of Example 11. As a result, as shown in Fig. 33, the anti-CCR4 chimeric antibody (48%) produced by the lectin-resistant clone CHO/CCR4-LCA showed 100 times or more higher binding activity to shFcγRIIIa than the anti-CCR4 chimeric antibody (8%) produced by the clone 5-03. Also, this activity was about 1/3 of the anti-CCR4 chimeric antibody (87%) produced by the YB2/0 cell-derived antibody-producing clone.

[0663] The above results clearly show that an antibody having 100 times or more higher binding activity to shFcγRIIIa can be prepared by using lectin-resistant CHO/DG44 cell than by using CHO/DG44 cell which is its parent cell.

Example 13

Method for screening antibody composition having high ADCC activity based on binding activity to shFcγRIIIa

[0664] An anti-GD3 chimeric antibody expression plasmid pChi641LHGM4 was introduced into the LCA-resistant clone CHO-LCA obtained in the item 1 of Example 7 according to the method described in the item 2(2) of Example 1, and gene amplification was carried out by using MTX to produce a transformant producing an anti-GD3 chimeric antibody. Cloning was carried out by limiting dilution method using the obtained transformant to obtain several clones. After each clone was cultured, the culture medium was recovered when it became confluent. A concentration of the anti-GD3 chimeric antibody in the culture supernatant was diluted to 1 μg/ml, and the shFcγRIIIa-binding activity was measured using the diluted antibody solution by the ELISA described in the item 1 of Example 11. At the same time, solutions of the anti-GD3 chimeric antibody produced by a YB2/0 cell-derived antibody-producing clone and the anti-GD3 chimeric antibody produced by a CHO/DG44 cell-derived antibody-producing clone, purified in the item 4 of Example 1, were prepared by diluting them to 1 μg/ml, and their shFcγRIIIa-binding activities were also measured.

[0665] Based on the measured results, a transformant cell clone capable of producing an antibody showing the activity equal to or higher than the binding activity of the anti-GD3 chimeric antibody produced by the CHO/DG44 cell-derived antibody-producing clone and also equal to or lower than the binding activity of the anti-GD3 chimeric antibody produced by the YB2/0 cell-derived antibody-producing clone was selected.

[0666] A purified antibody was obtained from the culture supernatant by culturing the selected transformant cell clone according to the method described in the item 4(2) of Example 1. When monosaccharide composition analysis of the purified antibody was carried out according to the method described in the item 1 of Example 3, the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond was 42%. Hereinafter, this sample is called anti-GD3 chimeric antibody (42%). When antigen-binding activity of the purified anti-GD3 chimeric antibody (42%) was evaluated by the ELISA described in the item 3 of Example 1, it was equivalent to those of the anti-GD3 chimeric antibody produced by a YB2/0 cell-derived antibody-producing clone and the anti-GD3 chimeric antibody produced by a CHO/DG44 cell-derived antibody-producing clone purified in the item 4 of Example 1. In addition, the ADCC activity of each anti-GD3 chimeric antibody was evaluated according to the method described in the item 2 of Example 2. For comparison, the ADCC activity of a sample of the anti-GD3 chimeric antibody produced by a CHO/DG44 cell-derived antibody-producing clone, wherein the ratio of a sugar chain in which 1-position of fucose was not bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond was 12% as a result of its monosaccharide composition analysis [hereinafter named "anti-GD3 chimeric antibody (12%)"] was measured. The results are shown in Fig. 34. In comparison with the anti-GD3 chimeric antibody (12%) produced by a CHO/DG44 cell-derived antibody-producing clone, about 30 times increase in the ADCC activity was observed in the anti-GD3 chimeric antibody (42%) produced by an antibody-producing clone selected based on the high binding activity to shFcγRIIIa.

[0667] Based on the above results, it was found that an antibody composition having high ADCC activity can be screened by screening an antibody composition having high binding activity to shFcγRIIIa.

Example 14

Preparation of FGF-8b/Fc fusion protein:

1. Construction of the expression vector of FGF-8b/Fc fusion protein

[0668] A humanized antibody expression vector pKANTEX93 [*Mol. Immunol.*, 37, 1035 (2000)] was digested with restriction enzymes *Apal* and *Bam*HI and a fragment containing about 1.0 kbp of human IgG1 subclass CH (hC γ 1) was obtained by using QIAquick Gel Extraction Kit (manufactured by QIAGEN). A plasmid pBluescript II SK(-) (manufactured by STRATAGENE) was also digested with similar restriction enzymes to obtain a fragment of about 2.9 kbp. The fragments were ligated using Solution I of TAKARA DNA Ligation Kit Ver. 2 (manufactured by Takara Shuzo) and *E. coli* DH5 α (manufactured by TOYOBO) was transformed to construct a plasmid pH γ 1/SK (-).

[0669] In order to ligate hC γ 1 with cDNA of FGF8, a synthetic DNA having the nucleotide sequence represented by SEQ ID NO:86 was designed. The synthetic DNA contains plural restriction enzyme recognition sequences at its 5'-terminal for cloning to pBluescript II SK(-), and the synthesis of the DNA was consigned to Proligo. To 50 μ l of a solution containing EX Taq Buffer (Mg $^{2+}$ plus) in TaKaRa Ex Taq (manufactured by Takara Shuzo) at 1 \times concentration, 1 ng of plasmid pH γ 1/SK (-), 0.25 mmol/L dNTPs, 0.5 μ mol/l of the synthetic DNA having the nucleotide sequence represented by SEQ ID NO:86, 0.5 μ mol/L M13 primer RV and 1.25 unit of TaKaRa Ex Taq were added, and the resulting solution was subjected to 30 cycles of heating at 94°C for 30 minutes, 56°C for 30 minutes and 72°C for 1 minute as one cycle by using DNA thermal cycler GeneAmp PCR System 9700 (manufactured by PERKIN ELMER). A PCR amplified fragment was purified from the total reaction solution by using QIAquick PCR purification Kit (manufactured by QIAGEN). Then, the purified fragment was digested with restriction enzymes *Kpn*I and *Bam*HI to obtain a fragment of about 0.75 kbp. In addition, the pBluescript II SK (-) (manufactured by STRATAGENE) was digested with similar restriction enzymes to obtain a fragment of about 2.9 kbp. These fragments were ligated by using Ligation high (manufactured by TOYOBO) and *E. coli* DH5 α (manufactured by TOYOBO) was transformed to construct a plasmid p Δ hC γ 1/SK(-).

[0670] Using the FGF-8b gene cloned plasmid pSC17 [*Proc. Natl. Acad. Sci.*, 89, 8928 (1992)] as the template, PCR was carried out as follows to obtain a structure gene region fragment of FGF-8b. To 50 μ l of a solution containing EX Taq Buffer (Mg $^{2+}$ plus) in TaKaRa Ex Taq (manufactured by Takara Shuzo) at 1x concentration, 1 ng of plasmid pSC17, 0.25 mmol/L dNTPs, 10 μ mol/l of synthetic DNAs having the nucleotide sequence represented by SEQ ID NOs:87 and 88 and 2.5 unit of TaKaRa Ex Taq were added, and the resulting solution was subjected to 35 cycles of heating at 94°C for 1 minute, 55°C for 1 minute and 72°C for 2 minutes as one cycle, followed by heating at 72°C for 10 minutes, by using DNA thermal cycler GeneAmp PCR System 9700 (manufactured by PERKIN ELMER). A PCR amplified fragment was purified from the total reaction solution, and the purified fragment was digested with restriction enzymes *Eco*RI and *Bam*HI to obtain a fragment of about 0.66 kbp. Also, plasmid pBluescript II SK(-) (manufactured by STRATAGENE), was digested with similar restriction enzymes to obtain a fragment of about 2.9 kbp. Then, the fragments were ligated by using T4 DNA Ligase (manufactured by Takara Shuzo) and *E. coli* DH5 α (manufactured by TOYOBO) was transformed to construct a plasmid pFGF-8b/SK(-).

[0671] Next, the plasmid p Δ hC γ 1/SK(-) was digested with restriction enzymes *Apal* and *Eco*RI to obtain a fragment of about 3.7 kbp. In addition, the plasmid pFGF-8b/SK(-) was digested with similar restriction enzymes to obtain a fragment of about 0.6 kbp. Then, the fragments were ligated by using Ligation high (manufactured by TOYOBO) and *E. coli* DH5 α (manufactured by TOYOBO) was transformed to construct a plasmid pFGF8b+hlgG/SK(-).

[0672] Next, the thus constructed plasmid pFGF8b+hlgG/SK (-) was digested with restriction enzymes *Eco*RI and *Bam*HI, a fragment of about 1.34 kbp was obtained. Also, pKANTEX93 was digested with similar restriction enzymes to obtain a fragment of about 8.8 kbp. Then, the fragments were ligated by using Ligation high (manufactured by TOYOBO) and *E. coli* DH5 α (manufactured by TOYOBO) was transformed to construct an expression vector pKANTEX/FGF8Fc for animal cell containing cDNA of FGF8b-Fc fusion protein represented by SEQ ID NO:89.

2. Stable expression using animal cell of FGF-8b/Fc fusion protein

[0673] A stable expression clone of FGF-8Fc fusion protein was prepared by introducing the FGF-8 fusion protein expression vector pKANTEX/FGF8Fc for animal cell which was constructed in the item 1 of Example 14 into various cells, and selecting a suitable clone.

(1) Preparation of producing cell using rat myeloma YB2/0 cells

[0674] After 10 μ g of the FGF8b-Fc fusion protein expression vector pKANTEX/FGF8Fc was introduced into 4 \times 10 6 cells of rat myeloma YB2/0 cell by electroporation, the resulting cells were suspended in 20 to 40 ml of RPMI1640-FBS

(10) and the solution was dispensed at 200 μ l/well into a 96-well culture plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, G418 was added thereto to give a concentration of 0.5 ml/ml, followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of G418 resistance transformants showing were formed and growth of colonies was observed, and the binding activity of the FGF8-Fc fusion protein in the supernatant to an anti-FGF-8 antibody was measured by the ELISA described in the item 4 of Example 14. As the anti-FGF-8 antibody, KM1334 (USP 5952472) was used.

[0675] Regarding the transformants in wells in which production of the FGF-8/Fc protein was observed in culture supernatants, in order to increase the production amount of the fusion protein by using a *dhfr* gene amplification system, each of them was suspended in the Hybridoma-SFM-FBS(5) medium comprising 0.5 mg/ml G418 and 50 nmol/L DHFR inhibitor, MTX (manufactured by SIGMA), and dispensed into each well of a 24 well plate for expansion culturing. After culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator to induce a transformant showing 50 nmol/l MTX resistance. Binding activity of FGF8b-Fc fusion protein in the supernatant in wells where growth of the transformant was observed, to KM1334 was measured by the ELISA described in the item 4 of Example 14.

[0676] Regarding the transformant in wells where production of FGF-8/Fc fusion protein was observed in the culture supernatant, according to the method similar to the above, a transformant KC1178 was obtained, which could grow in Hybridoma-SFM-FBS(5) medium comprising 0.5 mg/ml G418 and 200 nmol/l MTX as final concentrations by increase of the MTX concentration and highly produce FGF8-Fc fusion protein. It was found that KC 1178 was a lectin-resistant clone having a relatively low transcript amount obtained by the measuring method of FUT8 gene transcript described in Example 8 of WO00/61739. Also, KC1178 has been deposited on April 1, 2003, as FERM BP-8350 in International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology (AIST Tsukuba Central 6, 1-1, Higashi 1-Chome Tsukuba-shi, Ibaraki-ken, Japan).

(2) Preparation of producing cell using CHO/DG44 cells

[0677] After 10 μ g of the FGF8b-Fc fusion protein expression vector pKANTEX/FGF8Fc was introduced into 1.6×10^6 cells of CHO/DG44 cell by electroporation, the resulting cells were suspended in 30 ml of IMDM-dFBS(10)-HT(1) and the solution was dispensed at 100 μ l/well into a 96-well culture plate (manufactured by Sumitomo Bakelite). After culturing at 37°C for 24 hours in a 5% CO₂ incubator, the medium was changed to M7M-dFBE(10), followed by culturing for 1 to 2 weeks. The culture supernatant was recovered from wells in which colonies of transformants showing HT-independent growth were formed and growth of colonies was observed, and the binding activity of the FGF8-Fc fusion protein in the supernatant to KM1334 was measured by the ELISA shown in the item 4 of Example 14.

[0678] Regarding the transformants in wells in which production of the FGF-8/Fc protein was observed in culture supernatants, in order to increase the production amount of the fusion protein by using a *dhfr* gene amplification system, each of them was suspended in the IMDM-dFBS(10) medium comprising 50 nmol/L of MTX (manufactured by SIGMA), and dispensed into each well of a 24 well plate for expansion culturing. After culturing at 37°C for 1 to 2 weeks in a 5% CO₂ incubator to induce a transformant showing 50 nmol/l MTX resistance. Binding activity of FGF8b-Fc fusion protein in the supernatant in wells where growth was observed, to KM1334 was measured by the ELISA described in the item 4 of Example 14.

[0679] Regarding the transformant in wells where production of FGF-8/Fc fusion protein was observed in the culture supernatant, according to the method similar to the above, a transformant KC1179 was obtained, which could grow in IMDM-dFBS(10) medium comprising 500 nmol/l MTX by increase of the MTX concentration and highly produce FGF8-Fc fusion protein. Also, KC1179 has been deposited on April 1, 2003, as FERM BP-8351 in International Patent Organism Depository, National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1, Higashi 1-Chome Tsukuba-shi, Ibaraki-ken, Japan).

3. Purification of FGF8-Fc fusion protein

[0680] The producing cell of FGF8-Fc fusion protein prepared in the item 2 of Example 14 was cultured in an appropriate culture medium (e.g., H-SFM comprising 5% GF21 (manufactured by Wako Pure Chemical Industries), 0.5 mg/ml of G418 and 200 nmol/l of MTX for YB2/0 cell-derived cells; EXCELL301 (manufactured by JRH) comprising 500 nmol/l MTX for CHO/DG44 cell-derived cells) at a scale of 100 to 200 ml. FGF8-Fc fusion protein was purified from the culture supernatant by using Prosep G (manufactured by Millipore) column according to the manufacture's instruction. The deduced amino acid sequence of the purified protein is shown by SEQ ID NO:90.

4. Binding activity to anti-FGF-8 antibody

[0681] Binding activities of the FGF-8/Fc fusion protein produced by YB2/0 and the FGF-8/Fc fusion protein produced by CHO described in the item 2 of Example 14, to KM1334 were measured by the ELISA as follows. KM1334 of 1 μ g/

ml was dispensed at 50 μ l/well into a 96-well plate for ELISA (manufactured by Greiner) and allowed to stand at 4°C overnight for adsorption. After washing with PBS, 1% BSA-PBS was added at 100 μ l/well and allowed to react at room temperature for 1 hour to block the remaining active group. After washing each well with Tween-PBS, the culture supernatant of the transformant or a purified protein was added at 50 μ l/well, followed reaction at room temperature. After the reaction, each well was washed with Tween-PBS, and peroxidase labeled goat-anti-human IgG(γ) antibody solution (manufactured by American Qualex), which was diluted 3000-fold with 1% BSA-PBS, was added at 50 μ l/well as the secondary antibody solution, followed by reaction at room temperature for 1 hour. After the reaction, each well was washed with Tween-PBS, and ABTS substrate solution was added at 50 μ l/well, followed by reaction. After the color developed sufficiently, 5% SDS solution was added at 50 μ l/well to stop the reaction. Then, the absorbance was measured at a wavelength of 415 nm and a reference wavelength of 490. As shown in Fig. 35, the FGF-8/Fc fusion protein obtained in the item 2 of Example 14 showed binding activity to KM1334.

5. Binding activity to Fc γ RIIIa

[0682] Binding activities of FGF-8/Fc fusion protein produced by YB2/0 and FGF-8/Fc fusion protein derived from CHO described in the item 2 of Example 14, to Fc γ RIIIa was measured by ELISA as follows. Tetra-His Antibody, mouse antibody against His-tag (manufactured by QIAGEN), of 5 μ g/ml was dispensed at 50 μ l/well into a 96-well plate for ELISA (manufactured by GREINER) and allowed to stand at 4°C overnight for adsorption. After washing with PBS, 1% BSA-PBS was added at 100 μ l/well, followed by reaction at room temperature for 1 hour to block the remaining active group. After washing each well with Tween-PBS, shFc γ RIIIa (V) solution diluted to 5 μ g/ml with 1% BSA-PBS was added at 50 μ l/well, followed by reaction at room temperature for 2 hours. After the reaction, each well was washed with Tween-PBS, and a solution prepared by diluting purified FGF-8/Fc fusion protein to different concentrations with 1% BSA-PBS was added at 50 μ l/well, followed by reaction at room temperature for 2 hours. After the reaction, each well was washed with Tween-PBS, and biotinized KM1334 diluted to 1 μ g/ml with 1% BSA-PBS was added at 50 μ l/well, followed by reaction at room temperature for 1 hour. After the reaction, each well was washed with Tween-PBS, and a solution of peroxidase labeled Avidin-D (manufactured by VECTOR) diluted 4000-fold with 1% BSA-PBS was added at 50 μ l/well, followed by reaction at room temperature for 1 hour. After the reaction, each well was washed with Tween-PBS, and ABTS substrate solution was added at 50 μ l/well for color development, and after 15 minutes, 5% SDS solution was added at 50 μ l/well to stop the reaction. The absorbance of the resulting solution was measured at a wavelength of 415 nm and a reference wavelength of 490 nm.

[0683] Fig. 36 shows the measurement results of the binding activity of various FGF-8/Fc fusion proteins to shFc γ RIIIa (V). As apparent in Fig. 36, the FGF-8/Fc fusion protein produced by YB2/0 showed higher binding activity to shFc γ RIIIa (V) than the FGF-8/Fc fusion protein produced by CHO/DG44 cell.

Reference Example 1

[0684] Preparation of genes encoding various enzymes relating to sugar chain synthesis derived from CHO cell:

1. Determination of FX cDNA sequence in CHO cell

(1) Extraction of total RNA derived from CHO/DG44 cell

[0685] CHO/DG44 cells were suspended in IMDM medium containing 10% fetal bovine serum (manufactured by Life Technologies) and 1 \times concentration HT supplement (manufactured by Life Technologies), and 15 ml of the suspension was inoculated into a T75 flask for adhesion cell culture use (manufactured by Greiner) to give a density of 2 \times 10⁵ cells/ml. On the second day after culturing at 37°C in a 5% CO₂ incubator, 1 \times 10⁷ of the cells were recovered and a total RNA was extracted therefrom by using RNAeasy (manufactured by QIAGEN) in accordance with the manufacture's instructions.

(2) Preparation of total single-stranded cDNA from CHO/DG44 cell

[0686] The total RNA prepared in the item 1(1) of Reference Example 1 was dissolved in 45 μ l of sterile water, and 1 μ l of RQ1 RNase-Free DNase (manufactured by Promega), 5 μ l of the attached 10 \times DNase buffer and 0.5 μ l of RNasin Ribonuclease Inhibitor (manufactured by Promega) were added thereto, followed by reaction at 37°C for 30 minutes to degrade genomic DNA contaminated in the sample. After the reaction, the total RNA was purified again using RNAeasy (manufactured by QIAGEN) and dissolved in 50 μ l of sterile water.

[0687] In a 20 μ l of reaction mixture using oligo(dT) as a primer, single-stranded cDNA was synthesized from 3 μ g of the obtained total RNA samples by carrying out reverse transcription reaction using SUPERSRIPT™ Preamplic-

cation System for First Strand cDNA Synthesis (manufactured by Life Technologies) in accordance with the manufacture's instructions. A 50 fold-diluted aqueous solution of the reaction solution was used in the cloning of GFPP and FX. This was stored at -80°C until use.

(3) Preparation method of cDNA partial fragment of Chinese hamster-derived FX

[0688] An FX cDNA partial fragment derived from Chinese hamster was prepared by the following procedure.

[0689] First, primers (represented by SEQ ID NOs:42 and 43) specific for common nucleotide sequences registered at a public data base, namely a human FX cDNA (Genebank Accession No. U58766) and a mouse cDNA (Genebank Accession No. M30127), were designed.

[0690] Next, 25 µl of a reaction solution [1× concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/l dNTPs and 0.5 µmol/l of the above gene-specific primers (SEQ ID NOs:42 and 43)] containing 1 µl of the CHO/DG44-derived single-stranded cDNA prepared in the item 1(2) of Reference Example 1 was prepared by using a DNA polymerase ExTaq (manufactured by Takara Shuzo), and PCR was carried out. The PCR was carried out by heating at 94°C for 5 minutes, subsequent 30 cycles of heating at 94°C for 1 minute, 58°C for 2 minutes and 72°C for 3 minutes as one cycle, and further heating at 72°C for 10 minutes.

[0691] After the PCR, the reaction solution was subjected to 2% agarose gel electrophoresis, and a specific amplified fragment of 301 bp was purified using QuiaexII Gel Extraction Kit (manufactured by QIAGEN) and eluted with 20 µl of sterile water (hereinafter, the method was used for the purification of DNA fragments from agarose gel). Into a plasmid pCR2.1, 4 µl of the above amplified fragment was employed to insert in accordance with the instructions attached to TOPO TA Cloning Kit (manufactured by Invitrogen), and *E. coli* DH5α was transformed with the reaction solution. Each plasmid DNA was isolated in accordance with a known method from the obtained several kanamycin-resistant colonies to obtain 2 clones into which FX cDNA partial fragments were respectively inserted. They were named pCRFX clone 8 and pCRFX clone 12.

[0692] The nucleotide sequence of the cDNA inserted into each of the FX clone 8 and FX clone 12 was determined using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method of the manufacture's instructions. It was confirmed that each of the inserted cDNA whose sequence was determined encodes an ORF partial sequence of the Chinese hamster FX.

(4) Synthesis of single-stranded cDNA for RACE

[0693] Single-stranded cDNA samples for 5' and 3' RACE were prepared from the CHO/DG44 total RNA extracted in the item 1(1) of Reference Example 1 using SMART™ RACE cDNA Amplification Kit (manufactured by CLONTECH) in accordance with the manufacture's instructions. In the case, PowerScript™ Reverse Transcriptase (manufactured by CLONTECH) was used as the reverse transcriptase. Each single-stranded cDNA after the preparation was diluted 10-fold with the Tricin-EDTA buffer attached to the kit and used as the template of PCR.

(5) Determination of Chinese hamster-derived FX full length cDNA by RACE method

[0694] Based on the FX partial sequence derived from Chinese hamster determined in the item 1(3) of Reference Example 1, primers FXGSP1-1 (SEQ ID NO:44) and FXGSP1-2 (SEQ ID NO:45) for the Chinese hamster FX-specific 5' RACE and primers FXGSP2-1 (SEQ ID NO:46) and FXGSP2-2 (SEQ ID NO:47) for the Chinese hamster FX-specific 3' RACE were designed.

[0695] Next, 50 µl of a reaction solution [1× concentration Advantage2 PCR buffer (manufactured by CLONTECH), 0.2 mmol/L dNTPs, 0.2 µmol/l Chinese hamster FX-specific primers for RACE and 1 x concentration of common primers (manufactured by CLONTECH)] containing 1 µl of the CHO/DG44-derived single-stranded cDNA for RACE prepared in the item 1(4) of Reference Example 1 was prepared by using Advantage2 PCR Kit (manufactured by CLONTECH) and PCR was carried out. The PCR was carried out by 20 cycles of heating at 94°C for 5 seconds, 68°C for 10 seconds and 72°C for 2 minutes as one cycle.

[0696] After completion of the reaction, 1 µl of the reaction solution was diluted 50-fold with the Tricin-EDTA buffer, and 1 µl of the diluted solution was used as a template. The reaction solution was again prepared and the PCR was carried out under the same conditions. The templates, the combination of primers used in the first and second PCRs and the length of amplified DNA fragments by the PCRs are shown in Table 6.

Table 6

| Combination of primers used in Chinese hamster FX cDNA RACE PCR and the size of PCR products | | | |
|--|---------------------|-------------------------------|----------------------------|
| 5' RACE | FX-specific primers | Common primers | PCR-amplified product size |
| First | FXGSP1-1 | UPM (Universal primer mix) | 300 bp |
| Second | FXGSP1-2 | NUP (Nested Universal primer) | |
| 3' RACE | FX-specific primers | Common primers | PCR-amplified product size |
| First | FXGSP2-1 | UPM (Universal primer mix) | 1,100 bp |
| Second | FXGSP2-2 | NUP (Nested Universal primer) | |

[0697] After the PCR, the reaction solution was subjected to 1% agarose gel electrophoresis, and the specific amplified fragment of interest was recovered and eluted with 20 μ l of sterile water. Into a plasmid pCR2.1, 4 μ l of the amplified fragment was inserted, and *E. coli* DH5 α was transformed by using the reaction solution in accordance with the instructions attached to TOPO TA Cloning Kit (manufactured by Invitrogen). Plasmid DNAs were isolated from the obtained kanamycin-resistant colonies to obtain 5 cDNA clones containing Chinese hamster FX 5' region. They were named FX5' clone 25, FX5' clone 26, FX5' clone 27, FX5' clone 28, FX5' clone 31 and FX5' clone 32.

[0698] In the same manner, 5 cDNA clones containing Chinese hamster FX 3' region were obtained. These FX3' clones were named FX3' clone 1, FX3' clone 3, FX3' clone 6, FX3' clone 8 and FX3' clone 9.

[0699] The nucleotide sequence of the cDNA moiety of each of the clones obtained by the 5' and 3' RACE was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) in accordance with the method described in the manufacture's instructions. By comparing the cDNA nucleotide sequences determined by the method, reading errors of nucleotide bases due to PCR were excluded and the full length nucleotide sequence of Chinese hamster FX cDNA was determined. The determined sequence is represented by SEQ ID NO:48.

2. Determination of CHO cell-derived GFPP cDNA sequence

(1) Preparation of GFPP cDNA partial fragment derived from Chinese hamster

[0700] GFPP cDNA partial fragment derived from Chinese hamster was prepared by the following procedure.

[0701] First, nucleotide sequences of a human GFPP cDNA (Genebank Accession No. AF017445), mouse EST sequences having high homology with the sequence (Genebank Accession Nos. AI467195, AA422658, BE304325 and AI466474) and rat EST sequences (Genebank Accession Nos. BF546372, AI058400 and AW144783), registered at public data bases, were compared, and primers GFPP FW9 and GFPP RV9 (SEQ ID NOs:49 and 50) specific for rat GFPP were designed on a highly preserved region among these three species.

[0702] Next, 25 μ l of a reaction solution [1 \times concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs and 0.5 μ mol/l of the above GFPP-specific primers GFPP FW9 and GFPP RV9 (SEQ ID NOs:49 and 50)] containing 1 μ l of the CHO/DG44-derived single-stranded cDNA prepared in the item 1(2) of Reference Example 1 was prepared by using a DNA polymerase ExTaq (manufactured by Takara Shuzo), and PCR was carried out. The PCR was carried out by heating at 94°C for 5 minutes, subsequent 30 cycles of heating at 94°C for 1 minute, 58°C for 2 minutes and 72°C for 3 minutes as one cycle, and further heating at 72°C for 10 minutes.

[0703] After the PCR, the reaction solution was subjected to 2% agarose gel electrophoresis, and a specific amplified fragment of 1.4 Kbp was recovered and eluted with 20 μ l of sterile water. Into a plasmid pCR2.1, 4 μ l of the above amplified fragment was inserted in accordance with the instructions attached to TOPO TA Cloning Kit (manufactured by Invitrogen), and *E. coli* DH5 α was transformed by using the reaction solution. Plasmid DNAs were isolated from the obtained kanamycin-resistant clones to obtain 3 clones into which GFPP cDNA partial fragments were respectively integrated. They were named GFPP clone 8, GFPP clone 11 and GFPP clone 12.

[0704] The nucleotide sequence of the cDNA inserted into each of the GFPP clone 8, GFPP clone 11 and GFPP clone 12 was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) and BigDye Terminator Cycle Sequencing FS Ready Reaction kit (manufactured by Parkin Elmer) in accordance with the method described in the manufacture's instructions. It was confirmed that the inserted cDNA whose sequence was determined according to the present invention encodes an ORF partial sequence of the Chinese hamster GFPP.

(2) Determination of Chinese hamster GFPP full length cDNA by RACE method

[0705] Based on the Chinese hamster FX partial sequence determined in the item 2(1) of Reference Example 1, primers GFPP GSP1-1 (SEQ ID NO:52) and GFPP GSP1-2 (SEQ ID NO:53) for the Chinese hamster FX-specific 5' RACE and primers GFPP GSP2-1 (SEQ ID NO:54) and GFPP GSP2-2 (SEQ ID NO:55) for the Chinese hamster GFPP-specific 3' RACE were designed.

[0706] Next, 50 μ l of a reaction solution [$1\times$ concentration Advantage2 PCR buffer (manufactured by CLONTECH), 0.2 mmol/L dNTPs, 0.2 μ mol/l Chinese hamster GFPP-specific primers for RACE and $1\times$ concentration of common primers (manufactured by CLONTECH)] containing 1 μ l of the CHO/DG44 cell-derived single-stranded cDNA for RACE prepared in the item 1(4) of Reference Example 1 was prepared by using Advantage2 PCR Kit (manufactured by CLONTECH), and PCR was carried out. The PCR was carried out by 20 cycles of heating at 94°C for 5 seconds, 68°C for 10 seconds and 72°C for 2 minutes as one cycle.

[0707] After completion of the reaction, 1 μ l of the reaction solution was diluted 50-fold with the Tricin-EDTA buffer, and 1 μ l of the diluted solution was used as a template. The reaction solution was again prepared and the PCR was carried out under the same conditions. The templates, the combination of primers used in the first and second PCRs and the size of amplified DNA fragments by the PCRs are shown in Table 7.

Table 7

| Combination of primers used in Chinese hamster GFPP cDNA RACE PCR and the size of PCR products | | | |
|--|-----------------------|-------------------------------|----------------------------|
| 5' RACE | GFPP-specific primers | Common primers | PCR-amplified product size |
| First | GFPPGSP1-1 | UPM (Universal primer mix) | 1,100 bp |
| Second | GFPPGSP1-2 | NUP (Nested Universal primer) | |
| 3' RACE | GFPP-specific primers | Common primers | PCR-amplified product size |
| First | GFPPGSP2-1 | UPM (Universal primer mix) | 1,400 bp |
| Second | GFPPGSP2-2 | NUP (Nested Universal primer) | |

[0708] After the PCR, the reaction solution was subjected to 1% agarose gel electrophoresis, and the specific amplified fragment of interest was recovered and eluted with 20 μ l of sterile water. Into a plasmid pCR2.1, 4 μ l of the above amplified fragment was inserted and *E. coli* DH5 α was transformed with the reaction solution in accordance with the instructions attached to TOPO TA Cloning Kit (manufactured by Invitrogen). Plasmid DNAs were isolated from the obtained several kanamycin-resistant clones to obtain 4 cDNA clones containing Chinese hamster GFPP 5' region. They were named GFPP5' clone 1, GFPP5' clone 2, GFPP5' clone 3 and GFPP5' clone 4.

[0709] In the same manner, 5 cDNA clones containing Chinese hamster GFPP 3' region were obtained. They were named GFPP3' clone 10, GFPP3' clone 16 and GFPP3' clone 20.

[0710] The nucleotide sequence of the cDNA protein of each of the clones obtained in the above 5' and 3' RACE was determined by using DNA Sequencer 377 (manufactured by Parkin Elmer) in accordance with the method described in the manufacture's instructions. By comparing the cDNA nucleotide sequences after the nucleotide sequence determination, reading errors of bases due to PCR were excluded and the full length nucleotide sequence of Chinese hamster GFPP cDNA was determined. The determined sequence is represented by SEQ ID NO:51.

Reference Example 2

Preparation of CHO cell-derived GMD gene:

1. Determination of CHO cell-derived GMD cDNA sequence

(1) Preparation of CHO cell-derived GMD gene cDNA (preparation of partial cDNA excluding 5'- and 3'-terminal sequences)

[0711] Rodents-derived GMD cDNA was searched in a public data base (BLAST) using a human-derived GMD cDNA sequence (GenBank Accession No. AF042377) registered at GenBank as a query, and three kinds of mouse EST sequences were obtained (GenBank Accession Nos. BE986856, BF158988 and BE284785). By ligating these EST sequences, a deduced mouse GMD cDNA sequence was determined.

[0712] On the basis of the mouse-derived GMD cDNA sequence, a 28 mer primer having the nucleotide sequence

represented by SEQ ID NO:56, a 27 mer primer having the nucleotide sequence represented by SEQ ID NO:57, a 25 mer primer having the nucleotide sequence represented by SEQ ID NO:58, a 24 mer primer having the nucleotide sequence represented by SEQ ID NO:59 and a 25 mer primer having the sequence represented by SEQ ID NO:60 were prepared.

[0713] Next, in order to amplify the CHO cell-derived GMD cDNA, PCR was carried out by the following method. A reaction solution at 20 μ l [$1\times$ concentration Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 μ mol/L of two synthetic DNA primers] containing 0.5 μ l of the CHO cell-derived single-stranded cDNA prepared in the item 1(1) of Example 8 as the template was prepared. In this case, combinations of SEQ ID NO:56 with SEQ ID NO:57, SEQ ID NO:58 with SEQ ID NO:57, SEQ ID NO:56 with SEQ ID NO:59 and SEQ ID NO:56 with SEQ ID NO:60 were used as the synthetic DNA primers. The reaction was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle.

[0714] The PCR reaction solution was fractionated by agarose electrophoresis to find that a DNA fragment of about 1.2 kbp was amplified in the PCR product when synthetic DNA primers of SEQ ID NOs:56 and 57 were used, a fragment of about 1.1 kbp was amplified in the PCR product when synthetic DNA primers of SEQ ID NOs:57 and 59 were used, a fragment of about 350 bp was amplified in the PCR product when synthetic DNA primers of SEQ ID NOs:56 and 59 were used and a fragment of about 1 kbp was amplified in the PCR product when synthetic DNA primers of SEQ ID NOs:56 and 60 were used. The DNA fragments were recovered. The recovered DNA fragments were ligated to a pT7Blue(R) vector (manufactured by Novagen) by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5 strain (manufactured by Toyobo) was transformed by using the obtained recombinant plasmid DNA samples to thereby obtain plasmids 22-8 (having a DNA fragment of about 1.2 kbp amplified from synthetic DNA primers of SEQ ID NO:56 and SEQ ID NO:57), 23-3 (having a DNA fragment of about 1.1 kbp amplified from synthetic DNA primers of SEQ ID NO:58 and SEQ ID NO:57), 31-5 (a DNA fragment of about 350 bp amplified from synthetic DNA primers of SEQ ID NO:56 and SEQ ID NO:59) and 34-2 (having a DNA fragment of about 1 kbp amplified from synthetic DNA primers of SEQ ID NO:56 and SEQ ID NO:60). The CHO cell-derived GMD cDNA sequence contained in these plasmids was determined in the usual way by using a DNA sequencer ABI PRISM 377 (manufactured by Parkin Elmer) (since a sequence of 28 bases in downstream of the initiation codon methionine in the 5'-terminal side and a sequence of 27 bases in upstream of the termination codon in the 3'-terminal side are originated from synthetic oligo DNA sequences, they are mouse GMD cDNA sequences).

[0715] In addition, the following steps were carried out in order to prepare a plasmid in which the CHO cell-derived GMD cDNAs contained in the plasmids 22-8 and 34-2 are combined. The plasmid 22-8 (1 μ g) was allowed to react with a restriction enzyme *Eco*RI (manufactured by Takara Shuzo) at 37°C for 16 hours, the digest was subjected to agarose electrophoresis and then a DNA fragment of about 4 kbp was recovered. The plasmid 34-2 (2 μ g) was allowed to react with a restriction enzyme *Eco*RI at 37°C for 16 hours, the digest was subjected to agarose electrophoresis and then a DNA fragment of about 150 bp was recovered. The recovered DNA fragments were respectively subjected to terminal dephosphorylation using Calf Intestine Alkaline Phosphatase (manufactured by Takara Shuzo) and then ligated by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5 α strain (manufactured by Toyobo) was transformed by using the obtained recombinant plasmid DNA to obtain a plasmid CHO-GMD (cf. Fig. 37).

(2) Determination of 5'-terminal sequence of CHO cell-derived GMD cDNA

[0716] A 24 mer primer having the nucleotide sequence represented by SEQ ID NO:61 was prepared from 5'-terminal side non-coding region nucleotide sequences of CHO cell-derived human and mouse GMD cDNA, and a 32 mer primer having the nucleotide sequence represented by SEQ ID NO:62 from CHO cell-derived GMD cDNA sequence were prepared, and PCR was carried out by the following method to amplify cDNA. Then, 20 μ l of a reaction solution [$1\times$ concentration Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of Ex Taq polymerase (manufactured by Takara Shuzo) and 0.5 μ mol/L of the synthetic DNA primers of SEQ ID NO:61 and SEQ ID NO:62] containing 0.5 μ l of the CHO cell-derived single-stranded cDNA obtained in the item 1(1) of Example 8 was prepared as the template, and the reaction was carried out therein by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes, subsequent 20 cycles of heating at 94°C for 1 minute, 55°C for 1 minute and 72°C for 2 minutes as one cycle and further 18 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After fractionation of the PCR reaction solution by agarose electrophoresis, a DNA fragment of about 300 bp was recovered. The recovered DNA fragment was ligated to a pT7Blue(R) vector (manufactured by Novagen) by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5 α strain (manufactured by Toyobo) was transformed by using the obtained recombinant plasmid DNA samples to thereby obtain a plasmid 5'GMD. Using DNA Sequencer 377 (manufactured by Parkin Elmer), the nucleotide sequence of 28 bases in downstream of the initiation methionine of CHO -derived GMD cDNA contained in the plasmid was determined.

(3) Determination of 3'-terminal sequence of CHO cell-derived GMD cDNA

[0717] In order to obtain 3'-terminal cDNA sequence of CHO cell-derived GMD, RACE method was carried out by the following method. A single-stranded cDNA for 3' RACE was prepared from the CHO cell-derived RNA obtained in the item 1(1) of Example 8 by using SMART™ RACE cDNA Amplification Kit (manufactured by CLONTECH) in accordance with the manufacture's instructions. In the case, PowerScript™ Reverse Transcriptase (manufactured by CLONTECH) was used as the reverse transcriptase. The single-stranded cDNA after the preparation was diluted 10-fold with the Tricin-EDTA buffer attached to the kit and used as the template of PCR.

[0718] Next, 20 µl of a reaction solution [1× concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of EX Taq polymerase (manufactured by Takara Shuzo), 0.5 µmol/L of the 24 mer synthetic DNA primer represented by SEQ ID NO:63 [prepared on the basis of the CHO cell-derived GMD cDNA sequence determined in the item 1(1) of Reference Example 2] and 1x concentration of Universal Primer Mix (attached to SMART™ RACE cDNA Amplification Kit; manufactured by CLONTECH) containing 1 µl of the above single-stranded cDNA for 3' RACE as the template was prepared, and the reaction was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle.

[0719] After completion of the reaction, 1 µl of the PCR reaction solution was diluted 20-fold with Tricin-EDTA buffer (manufactured by CLONTECH). Then, 20 µl of a reaction solution [1× concentration ExTaq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of EX Taq polymerase (manufactured by Takara Shuzo), 0.5 µmol/L of the 25 mer synthetic DNA primer represented by SEQ ID NO:64 [prepared on the basis of the CHO cell-derived GMD cDNA sequence determined in the item 1(1) of Reference Example 2] and 0.5 µmol/L of Nested Universal Primer (attached to SMART™ RACE cDNA Amplification Kit; manufactured by CLONTECH)] containing 1 µl of the 20 fold-diluted aqueous solution as the template was prepared, and the reaction was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle.

[0720] After completion of the reaction, the PCR reaction solution was fractionated by agarose electrophoresis and then a DNA fragment of about 700 bp was recovered. The recovered DNA fragment was ligated to a pT7Blue(R) vector (manufactured by Novagen) by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α strain (manufactured by Toyobo) was transformed by using the obtained recombinant plasmid DNA, thereby obtaining a plasmid 3'GMD. Using DNA Sequencer 377 (manufactured by Parkin Elmer), the nucleotide sequence of 27 bases in upstream of the termination codon of CHO-derived GMD cDNA and 415 bp of 3'-terminal side non-coding region contained in the plasmid was determined.

[0721] The full length cDNA sequence of the CHO-derived GMD gene determined by the items 1(1), 1(2) and 1(3) of Reference Example 2 is represented by SEQ ID NO:65, and the corresponding amino acid sequence is represented by SEQ ID NO:71.

2. Determination of genomic sequence containing CHO/DG44-derived cell GMD gene

[0722] A 25 mer primer having the nucleotide sequence represented by SEQ ID NO:66 was prepared from the mouse GMD cDNA sequence determined in the item 1 of Reference Example 2. Next, a CHO cell-derived genomic DNA was obtained by the following method. A CHO/DG44 cell-derived KC861 cell was suspended in IMDM-dFBS(10)-HT(1) medium [IMDM-dFBS(10) medium comprising 1 x concentration of HT supplement (manufactured by Invitrogen)] to give a density of 3×10^5 cells/ml, and the suspension was dispensed at 2 ml/well into a 6 well flat bottom plate for adhesion cell use (manufactured by Greiner). After culturing at 37°C in a 5% CO₂ incubator until the cells became confluent on the plate, genomic DNA was prepared from the cells on the plate by a known method [*Nucleic Acids Research*, 3, 2303 (1976)] and dissolved overnight in 150 µl of TE-RNase buffer (pH 8.0) (10 mmol/l Tris-HCl, 1 mmol/l EDTA, 200 µg/ml RNase A).

[0723] A reaction solution (20 µl) [1× concentration Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of EX Taq polymerase (manufactured by Takara Shuzo) and 0.5 µmol/L of synthetic DNA primers of SEQ ID NO:59 and SEQ ID NO:66] containing 100 ng of the obtained CHO/DG44 cell-derived genomic DNA was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle. After completion of the reaction, the PCR reaction solution was fractionated by agarose electrophoresis and then a DNA fragment of about 100 bp was recovered. The recovered DNA fragment was ligated to a pT7Blue(R) vector (manufactured by Novagen) by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α strain (manufactured by Toyobo) was transformed by using the obtained recombinant plasmid DNA, thereby obtaining a plasmid ex3. Using DNA Sequencer 377 (manufactured by Parkin Elmer), the nucleotide sequence of CHO cell-derived genomic DNA contained in the plasmid was determined. The nucleotide sequence was represented by SEQ ID NO:67.

[0724] Next, a 25 mer primer having the nucleotide sequence represented by SEQ ID NO:68 and a 25 mer primer having the nucleotide sequence represented by SEQ ID NO:69 were prepared on the basis of the CHO cell-derived GMD cDNA sequence determined in the item 1 of Reference Example 2. Next, 20 µl of a reaction solution [1× concentration Ex Taq buffer (manufactured by Takara Shuzo), 0.2 mmol/L dNTPs, 0.5 unit of EX Taq polymerase (manufactured by Takara Shuzo) and 0.5 µmol/L of the synthetic DNA primers of SEQ ID NO:68 and SEQ ID NO:69] containing 100 ng of the CHO/DG44-derived genomic DNA was prepared, and PCR was carried out by using DNA Thermal Cycler 480 (manufactured by Perkin Elmer) by heating at 94°C for 5 minutes and subsequent 30 cycles of heating at 94°C for 1 minute and 68°C for 2 minutes as one cycle.

[0725] After completion of the reaction, the PCR reaction solution was fractionated by agarose electrophoresis and then a DNA fragment of about 200 bp was recovered. The recovered DNA fragment was ligated to a pT7Blue(R) vector (manufactured by Novagen) by using DNA Ligation Kit (manufactured by Takara Shuzo), and *E. coli* DH5α strain (manufactured by Toyobo) was transformed by using the obtained recombinant plasmid DNA, thereby obtaining a plasmid ex4. Using DNA Sequencer 377 (manufactured by Parkin Elmer), the nucleotide sequence of CHO cell-derived genomic DNA contained in the plasmid was determined. The result nucleotide sequence was represented by SEQ ID NO:70.

Reference Example 3

Preparation of anti-FGF-8 chimeric antibody

1. Isolation and analysis of a cDNA encoding the V region of a mouse antibody against FGF-8

(1) Preparation of mRNA from hybridoma cells which produces a mouse antibody against FGF-8

[0726] About 8 µg of mRNA was prepared from 1×10^7 cells of a hybridoma KM1334 (FERM BP-5451) which produces a mouse antibody against FGF-8 (anti-FGF-8 mouse antibody), using a mRNA preparation kit Fast Track mRNA Isolation Kit (manufactured by Invitrogen) according to the attached manufacture's instructions.

(2) Production of cDNA libraries of anti-FGF-8 mouse antibody H chain and L chain

[0727] A cDNA having *EcoRI*-*NotI* adapters on both termini was synthesized from 5 µg of the KM1334 mRNA obtained in the item 1(1) of Reference Example 3 by using Time Saver cDNA Synthesis Kit (manufactured by Amersham Pharmacia Biotech) according to the attached manufacture's instructions. A full amount of the prepared cDNA was dissolved in 20 µl of sterile water and then fractionated by agarose gel electrophoresis, and about 1.5 kb of a cDNA fragment corresponding to the H chain of an IgG class antibody and about 1.0 kb of a cDNA fragment corresponding to the L chain of a κ class were recovered each at about 0.1 µg. Next, 0.1 µg of the cDNA fragment of about 1.5 kb and 0.1 µg of the cDNA fragment of about 1.0 kb were respectively digested with restriction enzyme *EcoRI* and then ligated with 1 µg of λZAPII vector whose termini had been dephosphorylated with calf intestine alkaline phosphatase, using λZAPII Cloning Kit (manufactured by Stratagene) according to the attached manufacture's instructions.

[0728] Using Gigapack II Packaging Extracts Gold (manufactured by Stratagene), 4 µl of each reaction solution after ligation was packaged in λ phage according to the manufacture's instructions, and *Escherichia coli* XL1-Blue [*Biotechniques*, 5, 376 (1987)] was infected with an adequate amount of the package to obtain about 8.1×10^4 and 5.5×10^4 phage clones as H chain cDNA library and L chain cDNA library, respectively, of KM1334. Next, respective phages were immobilized on a nylon membrane according to a known method (*Molecular Cloning*, Second Edition).

(3) Cloning of cDNAs of anti-FGF-8 mouse antibody H chain and L chain

[0729] Nylon membranes of the H chain cDNA library and L chain cDNA library of KM1334 prepared in the item 1 (2) in Reference Example 3 were detected using a cDNA of the C region of a mouse antibody [H chain is a DNA fragment containing mouse Cγ1 cDNA (*J. Immunol.*, 146, 2010 (1991)), L chain is a DNA fragment containing mouse Cκ cDNA (*Cell*, 22, 197 (1980))] as a probe, using ECL Direct Nucleic Acid Labeling and Detection Systems (manufactured by Amersham Pharmacia Biotech) according to the attached manufacture's instructions, and phage clones strongly linked to the probe, 10 clones for each of H chain and L chain, were obtained. Next, each phage clone was converted into a plasmid by the *in vivo* excision method according to the manufacture's instructions attached to λZAPII Cloning Kit (manufactured by Stratagene). A nucleotide sequence of a cDNA contained in each of the obtained plasmids was determined by the dideoxy method (*Molecular Cloning*, Second Edition) by using Big Dye Terminator Kit ver. 2 (manufactured by Applied Biosystems). As a result, a plasmid pKM1334H7-1 containing a full length and functional H chain cDNA and a plasmid pKM1334L7-1 containing L chain cDNA, having an ATG sequence considered to be the initiation

codon in the 5'-terminal of the cDNA were obtained.

(4) Analysis of amino acid sequence of V region of anti-FGF-8 mouse antibody

[0730] A full length nucleotide sequence of VH contained in the plasmid pKM1334H7-1 and a deduced complete length amino acid sequence are represented by SEQ ID NO:72 and SEQ ID NO:73, respectively, and a full length nucleotide sequence of VL contained in the plasmid pKM1334L7-1 and a deduced complete length amino acid sequence are represented by SEQ ID NO:74 and SEQ ID NO:75, respectively. As a result of comparing these sequences to both known sequence data of mouse antibodies (*Sequences of Proteins of Immunological Interest*, U.S. Dept. Health and Human Services, 1991) and the comparison with the results of analysis of N-terminal amino acid sequences of H chain and L chain of the purified anti-FGF-8 mouse antibody KM1334, carried out by their automatic Edman degradation using a protein sequencer PPSQ-10 (manufactured by Shimadzu), it was found that each of the isolated cDNA is a full length cDNA encoding the anti-FGF-8 mouse antibody KM1334 containing a secretory signal sequence, and positions 1 to 19 in the amino acid sequence represented by SEQ ID NO:73 and positions 1 to 19 in the amino acid sequence described in SEQ ID NO:75 are secretory signal sequences of H chain and L chain, respectively.

[0731] Next, novelty of the amino acid sequences (sequences excluding secretory signal sequence) of VH and VL of the anti-FGF-8 mouse antibody KM1334 was examined. Using GCG Package (version 9.1, manufactured by Genetics Computer Group) as a sequence analyzing system, an amino acid sequence data base of known proteins (PIR-Protein (Release 56.0)) was searched by the BLAST method [*J. Mol. Biol.*, 215, 403 (1990)]. As a result, completely coincided sequences were not found for both of the H chain and L chain, so that it was confirmed that the VH and VL of the anti-FGF-8 mouse antibody KM1334 are novel amino acid sequences.

[0732] Also, the CDR of VH and VL of the anti-FGF-8 mouse antibody KM1334 was identified by comparing with amino acid sequences of known antibodies. Amino acid sequences of CDR 1, 2 and 3 of VH of the anti-FGF-8 mouse antibody KM1334 are represented by SEQ ID NOs:76, 77 and 78, respectively, and amino acid sequences of CDR 1, 2 and 3 of VL in SEQ ID NOs:79, 80 and 81, respectively.

2. Stable expression of anti-FGF-8 chimeric antibody using animal cell

(1) Construction of anti-FGF-8 chimeric antibody expression vector pKANTEX1334

[0733] An anti-FGF-8 chimeric antibody expression vector pKANTEX1334 was constructed as follows using the vector pKANTEX93 for humanized antibody expression described in W097/10354 and the plasmids pKM1334H7-1 and pKM1334L7-1 obtained in the item 1(3) of Reference Example 3.

[0734] Using 50 ng of the plasmid pKM1334H7-1 obtained in the item 1(3) of Reference Example 3 as the template and by adding synthetic DNAs having the nucleotide sequences described in SEQ ID NOs:24 and 25 (manufactured by GENSET) as primers to give a final concentration of 0.3 μ M, PCR were carried out in a system of 50 μ l by first heating at 94°C for 2 minutes and subsequent 30 cycles of heating at 94°C for 15 seconds, at 55°C for 30 seconds and at 68°C for 1 minute according to the manufacture's instructions attached to KOD plus polymerase (manufactured by TOYOBO). The reaction solution was precipitated with ethanol, dissolved in sterile water and then allowed to react at 37°C for 1 hour by using 10 units of a restriction enzyme *Apal* (manufactured by Takara Shuzo) and 10 units of a restriction enzyme *NotI* (manufactured by New England Biolabs). About 0.3 μ g of an *Apal*-*NotI* fragment of about 0.47 kb was recovered. By fractionating the reaction solution by agarose gel electrophoresis.

[0735] Next, 3 μ g of the vector pKANTEX93 for humanized antibody expression was allowed to react at 37°C for 1 hour by using 10 units of restriction enzyme *Apal* (manufactured by Takara Shuzo) and 10 units of restriction enzyme *NotI* (manufactured by New England Biolabs). About 2 μ g of an *Apal*-*NotI* fragment of about 12.75 kb was recovered, by fractionating the reaction solution by an agarose gel electrophoresis.

[0736] Next, 0.1 μ g of the *NotI*-*Apal* fragment derived from the PCR product and 0.1 μ g of the *NotI*-*Apal* fragment derived from the plasmid pKANTEX93, obtained in the above, were added to 10 μ l of sterile water in total amount and ligated by using Ligation High (manufactured by TOYOBO). The plasmid pKANTEX1334H shown in Fig. 38 was obtained by transforming *Escherichia coli* JM109 by using the recombinant plasmid DNA solution obtained in this manner.

[0737] Next, using 50 ng of the plasmid pKM1334L7-1 obtained in the item 1(3) of Reference Example 1 as the template and by adding synthetic DNAs having the nucleotide sequences described in SEQ ID NOs:82 and 83 (manufactured by GENSET) as primers to give a final concentration of 0.3 μ M, PCR was carried out in a system of 50 μ l by first heating at 94°C for 2 minutes and subsequent 30 cycles of heating at 94°C for 15 seconds, at 55°C for 30 seconds and 68°C for 1 minute according to the manufacture's instructions attached to KOD plus polymerase (manufactured by TOYOBO). The reaction solution was precipitated with ethanol, dissolved in sterile water and then allowed to react at 37°C for 1 hour by using 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and 10 units of a restriction enzyme *BsWI* (manufactured by New England Biolabs). About 0.3 μ g of an *EcoRI*-*BsWI* fragment of about

0.44 kb was recovered by fractionating the reaction solution by agarose gel electrophoresis.

[0738] Next, 3 µg of the plasmid pKANTEX1134H obtained in the above was allowed to react at 37°C for 1 hour by using 10 units of a restriction enzyme *EcoRI* (manufactured by Takara Shuzo) and a restriction enzyme *BsWI* (manufactured by New England Biolabs). About 2 µg of an *EcoRI*-*BsWI* fragment of about 13.20 kb was recovered by fractionating said reaction solution by an agarose gel electrophoresis.

[0739] Next, 0.1 µg of the *EcoRI*-*BsWI* fragment derived from the PCR product and 0.1 µg of the *EcoRI*-*BsWI* fragment derived from the plasmid pKANTEX1334H, obtained in the above, were added to 10 µl of sterile water in total amount and ligated by using Ligation High (manufactured by TOYOBO). The plasmid pKANTEX1334 shown in Fig. 38 was obtained by transforming *Escherichia coli* JM109 using the recombinant plasmid DNA solution obtained in this manner.

[0740] As a result of carrying out analysis of a nucleotide sequence using 400 ng of the obtained plasmid by the dideoxy method (*Molecular Cloning*, Second Edition) using Big Dye Terminator Kit ver. 2 (manufactured by Applied Biosystems), it was confirmed that a plasmid comprising a cloned DNA of interest was obtained.

INDUSTRIAL APPLICABILITY

[0741] The present invention relates to a method for enhancing a binding activity of an antibody composition to Fcγ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule; a method for enhancing an antibody-dependent cell-mediated cytotoxic activity of an antibody composition; a process for producing an antibody composition having an enhanced binding activity to Fcγ receptor IIIa; a method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among total complex *N*-glycoside-linked sugar chains bound to the Fc region in an antibody composition; an Fc fusion protein composition produced by using a cell resistant to a lectin which recognizes a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α-bond in a complex *N*-glycoside-linked sugar chain; and a process for producing the same.

Free Text in Sequence Listing

[0742]

SEQ ID NO:4 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:5 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:8 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:9 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:10 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO: 11 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:12 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:13 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:14 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:15 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:16 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:17 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:18 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:22 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:26 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:29 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:32 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:33 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:34 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:35 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:36 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:37 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:38 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:39 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:40 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:41 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:42 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:43 - Explanation of synthetic sequence: synthetic DNA

SEQ ID NO:44 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:45 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:46 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:47 - Explanation of synthetic sequence: synthetic DNA
 5 SEQ ID NO:49 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:50 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:52 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:53 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:54 - Explanation of synthetic sequence: synthetic DNA
 10 SEQ ID NO:55 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:56 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:57 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:58 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:59 - Explanation of synthetic sequence: synthetic DNA
 15 SEQ ID NO:60 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:61 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:62 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:63 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:64 - Explanation of synthetic sequence: synthetic DNA
 20 SEQ ID NO:66 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:68 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:69 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:82 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:83 - Explanation of synthetic sequence: synthetic DNA
 25 SEQ ID NO:84 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:85 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:86 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:87 - Explanation of synthetic sequence: synthetic DNA
 SEQ ID NO:88 - Explanation of synthetic sequence: synthetic DNA
 30 SEQ ID NO:89 - Explanation of synthetic sequence: synthetic DNA

35

40

45

50

55

SEQUENCE LISTING

<110> KYOWA HAKKO KOGYO CO., LTD.

<120> Method of enhancing of binding activity of antibody composition to Fcγ receptor IIIa

<130> P044077

<150> P2002-106950

<151> 2002-04-09

<160> 100

<170> PatentIn Ver. 2.1

<210> 1

<211> 2008

<212> DNA

<213> Cricetulus griseus

<400> 1

```

aacagaaact tattttcctg tgtggctaac tagaaccaga gtacaatgtt tccaattctt 60
tgagctccga gaagacagaa gggagttgaa actctgaaaa tgcgggcatg gactgggtcc 120
tggcggttga ttatgctcat tctttttgcc tgggggacct tattgtttta tatagggtgt 180
catttggttc gagataatga ccaccctgac cattctagca gagaactctc caagattctt 240
gcaaagctgg agcgcttaaa acaacaaaat gaagacttga ggagaatggc tgagtctctc 300
cgaataccag aaggccctat tgatcagggg acagctacag gaagagtccg tgtttttagaa 360
gaacagcttg ttaaggccaa agaacagatt gaaaattaca agaaacaagc taggaatgat 420
ctgggaaagg atcatgaaat cttaaggagg aggattgaaa atggagctaa agagctctgg 480
ttttttctac aaagtgaatt gaagaaatta aagaaattag aaggaaacga actccaaga 540
catgcagatg aaattctttt ggatttagga catcatgaaa ggtctatcat gacagatcta 600
tactacctca gtcaaacaga tggagcaggt gagtggcggg aaaaagaagc caaagatctg 660
acagagctgg tccagcggag aataacatat ctgcagaatc ccaaggactg cagcaaagcc 720
agaaagctgg tatgtaatat caacaaaggc tgtggtatg gatgtcaact ccatcatgtg 780
gtttactgct tcatgattgc ttatggcacc cagcgaacac tcatcttgga atctcagaat 840
tggcgctatg ctactggagg atgggagact gtgttttagac ctgtaagtga gacatgcaca 900
gacaggctctg gcctctccac tggacactgg tcagggtgaag tgaaggacaa aaatgttcaa 960

```

gtggtcgagc tccccattgt agacagcctc catcctcgtc ctccctactt acccttggct 1020
 gtaccagaag accttgcaga tcgactcctg agagtccatg gtgatcctgc agtgtgggtg 1080
 gtatcccagt ttgtcaaata cttgatccgt ccacaacctt ggctggaaag ggaaatagaa 1140
 gaaaccacca agaagcttgg cttcaaacat ccagttattg gagtccatgt cagacgcact 1200
 gacaaagtgg gaacagaagc agccttccat cccattgagg aatacatggt acacgttgaa 1260
 gaacattttc agcttctcga acgcagaatg aaagtggata aaaaaagagt gtatctggcc 1320
 actgatgacc ctcttttgtt aaaggaggca aagacaaagt actccaatta tgaatttatt 1380
 agtgataact ctatttcttg gtcagctgga ctacacaacc gatacacaga aaattcactt 1440
 cggggcggtga tcttgatat acactttctc tcccaggctg acttccctgt gtgtactttt 1500
 tcatcccagg tctgtagggt tgcttatgaa atcatgcaaa cactgcatcc tgatgcctct 1560
 gcaaaacttc attctttaga tgacatctac tattttggag gccaaaatgc ccacaaccag 1620
 attgcagttt atcctcacca acctcgaact aaagaggaaa tcccatgga acctggagat 1680
 atcattgggtg tggctggaaa ccattggaat ggttactcta aagggtgcaa cagaaaacta 1740
 ggaaaaacag gcctgtaccc ttcttcaaaa gtccgagaga agatagaaac agtcaaatac 1800
 cctacatctc ctgaagctga aaaatagaga tggagtgtaa gagattaaca acagaattta 1860
 gttcagacca tctcagccaa gcagaagacc cagactaaca tatggttcat tgacagacat 1920
 gctccgcacc aagagcaagt gggaaccctc agatgctgca ctggtggaac gcctctttgt 1980
 gaagggtgc tgtgccctca agcccatg 2008

<210> 2

<211> 1728

<212> DNA

<213> *Mus musculus*

<400> 2

atgcgggcat ggactggttc ctggcggttg attatgctca ttctttttgc ctgggggacc 60
 ttgttatttt atataggttg tcatttgggt cgagataatg accaccctga tcactccagc 120
 agagaactct ccaagattct tgcaaagctt gaacgcttaa aacagcaaaa tgaagacttg 180
 aggcgaatgg ctgagtctct ccgaatacca gaaggcccca ttgaccaggg gacagctaca 240
 ggaagagtcc gtgtttttaga agaacagctt gttaaggcca aagaacagat tgaaaattac 300
 aagaaacaag ctagaaatgg tctggggaag gatcatgaaa tcttaagaag gaggattgaa 360
 aatggagcta aagagctctg gtttttcta caaagcgaac tgaagaaatt aaagcattta 420
 gaaggaaatg aactccaaag acatgcagat gaaattcttt tggatttagg acaccatgaa 480
 aggtctatca tgacagatct atactacctc agtcaaacag atggagcagg ggattggcgt 540
 gaaaaagagg ccaagatct gacagagctg gtccagcgga gaataacata tctccagaat 600
 cctaaggact gcagcaaagc caggaagctg gtgigtaca tcaataaagg ctgtggctat 660
 ggttgtcaac tccatcacgt ggtctactgt ttcatgattg cttatggcac ccagcgaaca 720

ctcatcttgg aatctcagaa ttggcgctat gctactgggtg gatgggagac tgtgtttaga 780
 cctgtaagtg agacatgtac agacagatct ggccctctcca ctggacactg gtcaggtgaa 840
 gtaaatgaca aaaacattca agtggctcag ctccccattg tagacagcct ccattcctcg 900
 cctccttact taccactggc tgttccagaa gaccttgacg accgactcct aagagtccat 960
 ggtgaccctg cagtgtgggtg ggtgtcccag tttgtcaaat acttgattcg tccacaacct 1020
 tggctggaaa aggaaataga agaagccacc aagaagcttg gcttcaaaca tccagttatt 1080
 ggagtccatg tcagacgcac agacaaagtg ggaacagaag cagccttcca ccccatcgag 1140
 gagtacatgg tacacgttga agaacatttt cagcttctcg cagcgagaat gcaagtggat 1200
 aaaaaagag tatatctggc tactgatgat cctactttgt taaaggaggc aaagacaaag 1260
 tactccaatt atgaatttat tagtgataac tctatttctt ggtagctgg actacacaat 1320
 cggtagacag aaaattcact tcgggggtgtg atcctggata tacactttct ctcacaggct 1380
 gactttctag tgtgtacttt ttcattccag gtcgtcggg ttgcttatga aatcatgcaa 1440
 accctgcac ctagatccctc tgcgaacttc cattcttttg atgacatcta ctattttgga 1500
 ggccaaaatg cccacaatca gattgctgtt tatcctcaca aacctcgaa tgaagaggaa 1560
 attccaatgg aacctggaga tatcattggt gtggctggaa accattggga tggttattct 1620
 aaaggatatc acagaaaact tggaaaaaca ggcttatatc cctcctacaa agtccgagag 1680
 aagatagaaa cagtcaagta tcccacatat cctgaagctg aaaaatag 1728

<210> 3

<211> 9196

<212> DNA

<213> Cricetulus griseus

<400> 3

tctagaccag gctggctctg aactcacaga gaaccacctg cctctgccac ctgagtgtg 60
 ggattaaagg tgtgcaccac caccgccgg cgtaaaatca tatttttgaa tattgtgata 120
 atttacatta taattgtaag taaaaatttt cagcctattt tgttatacat ttttgcgtaa 180
 attattcttt ttgaaagt ttgtttgcc taatagtcta gggaaacata aagttataat 240
 tttgtctat gtatttgc atatatctat ttaatctcct aatgtccagg aaataaatag 300
 ggtatgtaat agcttcaaca tgtggatatga tagaattttt cagtgtctata taagttgtta 360
 cagcaaagtg ttattaattc atatgtccat atttcaattt tttatgaatt attaaattga 420
 atccttaagc tgccagaact agaattttat tttaatcagg aagccccaaa tctgttcatt 480
 ctttctatat atgtggaaag gtaggcctca ctaactgatt cttcacctgt tttagaacat 540
 ggtccaagaa tggagttag taaggggaat tacaagtgtg agaaaactcc tagaaaacaa 600
 gatgagtctt gtgaccttag ttcttttaaa aacacaaaat tcttggaaat tgttttcatt 660
 ttctctccag gtggatagga gtgagtttat ttcagattat ttattacaac tggctgttgt 720
 tacttgtttc tatgtcttta tagaaaaaca tatttttttt gccacatgca gcttgcctt 780

atgattttat acttgtgtga ctcttaactc tcagagtata aattgtctga tgctatgaat 840
 aaagttggct attgtatgag acttcagccc acttcaatta ttggcttcat tcctcagat 900
 5 cccaccacct ccagagtggg aaacaacttg aaccattaaa cagacttttag tctttatttg 960
 aatgatagat ggggatatca gatttatagg cacagggttt tgagaaaggg agaaggtaaa 1020
 cagtagagtt taacaacaac aaaaagtata cttigttaaac gtaaaactat ttattaaagt 1080
 10 agtagacaag acattaaata ttcttggga ttagtgttt ttgaattttg ctttcaaata 1140
 atagtcagtg agtatacccc tccccattc tatatttttag cagaaatcag aataaatggt 1200
 gtttcigtga cattcttttg tagagaattt atttctttg ggtttttgtg catttaaagt 1260
 15 caataaaaat taagggtcag taatagaaaa aaactctga tttttggaat cccctttctt 1320
 cagcttttct atttaactc ttaatgataa ttaattttgt ggccatgtgg tcaaagtata 1380
 tagccttgta tatgtaaatg ttttaaccaa cctgccttta cagtaactat ataattttat 1440
 tctataatat atgacttttc ttccatagct tttagagttgc ccagtcactt taagttacat 1500
 20 tttcatatat gttctttgtg ggaggagata attttatttc taagagaatc ctaagcatac 1560
 tgattgagaa atggcaaaca aaacacataa ttaaagctga taaagaacga acatttggag 1620
 tttaaaatac atagccaccc taagggttta actgttgta gccttctttt ggaattttta 1680
 ttagttcata tagaaaaatg gattttatcg tgacatttcc atatatgtat ataataat 1740
 25 tacatcatat ccacctgtaa ttattagtgt ttttaaatat atttgaanaa ataatggtct 1800
 ggtttgatcc atttgaacct ttgatgttt ggtgtggttg ccaattgggt gatggttatg 1860
 ataacctttg cttctctaag gttcaagta gtttgagaat atgtcctcta aaaatgacag 1920
 30 gttgcaagtt aagtagtgag atgacagcga gatggagtga tgagaatttg tagaatgaa 1980
 ttcacttata ctgagaactt gttttgcttt tagataatga acatattagc ctgaagtaca 2040
 tagccgaatt gattaattat tcaaagatat aatcttttaa tccctataaa agaggattta 2100
 cacaacaatt caagaaagat agaattagac ttccagtatt ggagtgaacc atttgttacc 2160
 35 aggtagaacc ctaacgtgtg tggttgactt aaagtgttta ctttttacct gatactgggt 2220
 agctaattgt ctttcagcct cctggccaaa gataccatga aagtcaactt acgttgtatt 2280
 ctatatctca aacaactcag ggtgtttctt actctttcca cagcatgtag agcccaggaa 2340
 40 gcacaggaca agaaagctgc ctcttgtat caccaggaag atctttttgt aagagtcac 2400
 acagtatacc agagagacta attttgtctg aagcatcatg tgttgaaaca acagaaactt 2460
 attttcctgt gtggctaact agaaccagag tacaatgttt ccaattcttt gagctccgag 2520
 aagacagaag ggagttgaaa ctctgaaaa gcgggcatgg actggttcct ggcgttggat 2580
 45 tatgtcatt ctttttgcct gggggacctt attgttttat atagggtgtc atttggttcg 2640
 agataatgac caccctgacc attctagcag agaactctcc aagattcttg caaagctgga 2700
 gcgccttaaaa caacaaaatg aagacttgag gagaatggct gagtctctcc ggtaggtttg 2760
 50 aaatactcaa ggatttgatg aaatactgtg cttgacctt aggtataggg tctcagtcg 2820
 ctgttgaaaa atataatttc tacaaccgt ctttgtaaaa ttttaagtat ttagcagac 2880
 tttttaaaag tcagtatac atctatatag tcaatatagg ttacatagt tgcaatctta 2940
 ttttgcatat gaatcagtat atagaagcag tggcatttat atgcttatgt tgcatttaca 3000
 55 attatgttta gacgaacaca aactttatgt gatttggatt agtgctcatt aaattttttt 3060

attctatgga ctacaacaga gacataaatt ttgaaaggct tagttactct taaattctta 3120
 5 tgatgaaaag caaaaattca ttgttaaata gaacagtgca tccggaatgt gggttaattat 3180
 tgccatattt ctagtctact aaaaattigt gcataactgt tcaaagtcac cagttgtttg 3240
 gaaagccaaa gtctgattta aatggaaaac ataaacaatg atatctattt ctagatacct 3300
 10 ttaacttgca gttactgagt ttacaagttg tctgacaact ttggattctc ttacttcata 3360
 tctaagaatg atcatgtgta cagtgcctac tgtcacttta aaaaactgca gggctagaca 3420
 tgcagatatg aagactttga cattagatgt ggtaattggc actaccagca agtgggtatta 3480
 agatacagct gaatatatta ctttttgagg aacataattc atgaatggaa agtggagcat 3540
 15 tagagaggat gccttctggc tctccacac cactgtttgc atccattgca tttcacactg 3600
 cttttagaac tcagatgttt catatgggat attgtgtaac tcaccatcag ttttatcttt 3660
 aaatgtctat ggatgataat gttgtatgtt aacactttta caaaaacaaa tgaagccata 3720
 tcttcgggtg gagtgtgat ggtggtaatt gtcacaatag gattattcag caaggaaacta 3780
 20 agtcagggac aagaagtggt cgatactttg ttggattaaa tcattttact ggaagttcat 3840
 cagggagggt tatgaaagt gtggtccttg aactgaaatt atatgtgatt cattattctt 3900
 gatttaggcc ttgctaatag taactatcat ttattgggaa ttgtcatat gtgccaattt 3960
 25 gtcatgggcc agacagcgtg ttttactgaa tttctagata tctttatgag attctagtag 4020
 tgttttcagc cttttacag atgaagaatc ttaaaaaatg ttaaataatt tagtttgccc 4080
 aagattatac gttacaacaaat ggtagaacct tctttgaatt ctggcagtat ggctacacag 4140
 tccgaactct tatcttctta agctgaaaac agaaaaagca atgaccagaa aaattttatt 4200
 30 taaaagtcct aggagagact tcccatcctg agaagatctc ttttcccttt tataatttag 4260
 gctcctgaat aatcaactgaa ttttctccat gttccatcta tagtactgtt atttctgttt 4320
 tcttttttct ttaccacaaa gtatcttgtt ttigtctgtat gaaagaaaat gtgttattgt 4380
 35 aatgtgaaat tctctgtccc tgcaggggtc cactccgcc tcaatcccaa ataaacacac 4440
 agaggctgta ttaattatga aactgtttgt cagttggcta gggcttctta ttggctagct 4500
 ctgtcttaat tattaaccaa taactactat tgtaagtatt tccatgtggt cttatcttac 4560
 caaggaaagg gtccaggac ctttactcc tctggcgtgt tggcagtgaa gaggagagag 4620
 40 cgatttctta tttgtctctg cttattttct gattctgctc agctatgtca cttcctgctt 4680
 ggccaatcag ccaatcagtg ttttattcat tagccaataa aagaacatt tacacagaag 4740
 gacttcccc atcatgttat ttgtatgagt tcttcagaaa atcatagtat cttttaatac 4800
 45 taatttttat aaaaaattaa ttgtattgaa aattatgtgt atatgtgtct gtgtgtcgat 4860
 ttgtgtcat aagtagcatg gagtgcagaa gagggaaatca gatctttttt taagggacaa 4920
 agagtttatt cagattacat ttaagggtga taatgtatga ttgcaagggt atcaacatgg 4980
 cagaaatgtg aagaagctgg tcacattaca tccagagtca agagtagaga gcaatgaatt 5040
 50 gatgcatgca ttctgtgct cagctcactt ttctggagc tgagctgatt gtaagccatc 5100
 tgatgtcttt gctgggaact aactcaaagg caagttcaaa acctgttctt aagtataagc 5160
 catctctcca gtccctcata tggctctta agacacittc tttatattct tgtacataga 5220
 55 aattgaattc ctaacaactg cattcaaatt acaaaatagt ttttaaaagc tgatataata 5280
 aatgtaata caatctagaa catttttata aataagcata ttaactcagt aaaaataaat 5340

gcatggttat tttccttcat tagggaagta tgtctcccca ggctgttctc tagattctac 5400
 tagtaatgct gtttgtacac catccacagg ggttttatatt taaagctaag acatgaatga 5460
 5 tggacatgct tgtagcatt tagacttttt tcttactat aattgagcta gtatttttgt 5520
 gctcagtttg atactgtta attcagataa atgtaatagt aggtaatttc ttigtgataa 5580
 aggcataata attgaagtig gaaaacaaaa gcctgaaatg acagttttta agattcagaa 5640
 10 caataatttt caaaagcagt taccacactt tccaaataca atctgcagtt ttcttgatat 5700
 gtgataaatt tagacaaaga aatagcacat tttaaaatag ctatttactc ttgatttttt 5760
 tttcaaatat aggctagtgc actagtgttg tgtaaggta tggtgcgaaa catctttgac 5820
 15 tcttggttag ggaatccagg atgatttacg tgtttggcca aaatcttggt ccattctggg 5880
 tttcttctct atctaggttag ctagcacaag ttaaagggtg ggtagtattg gaaggctctc 5940
 aggtatataat ttctatattc tgtatttttt tctctgtca tatatttgct ttctgtttta 6000
 ttgatttcta ctgttagttt gatacttact ttcttacct ttctttggga tttattttgc 6060
 20 tgttctaaga tttcttagca agttcatatc actgatttta acagtgtgct cttttgtaat 6120
 atagactgaa tgcccttat ttgaaatgct tgggatcaga aacacagatt tgaacttttc 6180
 ttttttaata ttccatcaa gtttaccagc tgaatgtcct gatccaagaa tatgaaatct 6240
 25 gaaatgcttt gaaatctgaa acttttagag tgataaagct tccctttaaa ttaattgttg 6300
 ttctatattt ttgacaaag tcaacctttc attgttatcc aatgagtga catattttca 6360
 atttttttgt ttgatctgtt atattttgat ctgaccatat ttataaaatt ttatttaatt 6420
 tgaatgttgt gctgttactt atctttatta ttatttttgc ttattttcta gccaaatgaa 6480
 30 attatactct gtattatttt agtttgaatt ttactttgtg gcttagtaac tgccttttgt 6540
 tggatgaatgc ttaagaaaaa cgtgttgtct actgatattg gttctaactt tatatagcat 6600
 gttgtttgtt aggtagtga ttatgctggt cagattgtct tgagtttatg caaatgtaaa 6660
 atatttagat gcttgttttg ttgtctaaga acaaagtatg ctgtctgtct cctatcggtt 6720
 35 ctggtttttc cattcatctc ttcaagctgt ttgtgtgtt gaataactaac tccgtactat 6780
 cttgttttct gtgaattaac cctttttcaa aggtttcttt tctttttttt ttttaaggac 6840
 aacaagttaa ttacagattac attttaagct gataatgtat gattgcaagg ttatcaacat 6900
 40 ggcagaaatg tgaagaagct aggcacatta catccacatg gagtcaagag cagagagcag 6960
 tgaattaatg catgcattcc tgtggtcagc tcacttttcc tattcttaga tagtctagga 7020
 tcataaacct ggggaatagt gctaccacaa tgggcataac cacttacttc agttcatgca 7080
 atcaaccaag gcacatccac aggaaaaact gatthagaca acctctcatt gagactcttc 7140
 45 ccagatgatt agactgtgtc aagttgacaa ttaaaactat cacacctgaa gccatcacta 7200
 gtaaatataa tgaaaatgtt gattatcacc ataattcacc tgtatccctt tgttattgta 7260
 gattttgtga agttcctatt caagtcctg ttcccttctt aaaaacctgt ttttagtta 7320
 50 aataggtttt ttagtgttcc tgtctgtaaa tactttttta aagttagata ttattttcaa 7380
 gtatgttctc ccagcttttg gcttgtattt tcatcccttc aatacatata tttttgtaat 7440
 ttattttttt tatttaaaatt agaaacaaag ctgcttttac atgtcagttc cagtccctc 7500
 55 tccctccctt cctccctgc tccccaccta agcccaatt ccaactcctt tcttctccc 7560
 aggaagggtg aggcctcca tgggggaaat cttcaatgic tgcataatca ttggagcag 7620

5 ggccatagacc ctccccagtg tgtctaggct gagagagtat ccctctatgt ggagagggct 7680
 cccaaagtgc atttgtgtac taggggtaaa tactgatcca ctatcagtgg ccccatagat 7740
 tgtccggacc tccaaactga ctctctcctt cagggagtct ggaacagttc tatgttggtt 7800
 tcccagatat cagtctgggg tccatgagca accccttggt caggtcagtt gtttctgtag 7860
 10 gtttccccag cccggctctt acccctttgc tcatcacttc tccctctctg caactggatt 7920
 ccagagttca gctcagtggt tagctgtggg tgtctgcacg tgcctccatc agctactgga 7980
 tgagggtctt aggatggcat ataaggtagt catcagtcct attatcagag aagggtcttt 8040
 aaggtagcct cttagattatt gcttagattg ttagttgggg tcaaccttgt aggtctctgg 8100
 15 acagtgcacg aattctcttt aaacctataa tggctccctc tgtggtggta tcccttttct 8160
 tgcctctatc cgttctctcc ctgactagat ctctctgctc cctcatgtcc tctctctccc 8220
 tccccctctc ccttctcttt tcttctaact cctctctccc tccaccacg atccccatta 8280
 gcttatgaga tcttgtcctt attttagcaa aacctttttg gctataaaat taattaattt 8340
 20 aatatgctta tatcagggtt attttgcta gtatttgtat gtgtttggtt agtgttttta 8400
 accttaattg acatgtatcc ttatatattg acacagattt aaatatattg agtttttttt 8460
 tttttttttt ttaaagattt atttattttt tatgtcttct gcctgcatgc cagaagaggg 8520
 25 caccagatct cattcaaggt ggttgtgagc caccatgtgg ttgctgggaa ttgaactcag 8580
 gacctctgga agaacagtcg gtgctcttaa ccgctgagcc atctctccag cccctgaagt 8640
 gtttctttta aagaggatag cagtgcacga tttttccctt tgaccaatga ctctacatt 8700
 actgaattgt tttagccatt tataatgtaat gctgttacca ggtttacatt ttcttttctc 8760
 30 ttgctaaatt tcttccctgt ttgtctcatc tcttattttt gtctgttgga ttatataggc 8820
 ttttattttt ctgtttttac agtaagttat atcaaattaa aattatttta tggaatgggt 8880
 gtgttgacta catgtatgtc tgtgcaccat gtgctgacct ggtcttggcc agaagaaggt 8940
 35 gtcataattct ctgaaactgg tattgtggat gttacgaact gccatagggt gctaggaatc 9000
 aaacccacgc tctcttgga aagcagccac tgctctgagc cactgagtc tctcttcaag 9060
 cagggtgatgc caacttttaa tggttaccag tggataagag tgcttgatc tctagcacc 9120
 atgaaaattt atgcattgct atatgggctt gtcacttcag catttgttga cagagacagg 9180
 40 aggatcccaa gagctc 9196

45 <210> 4

<211> 25

<212> DNA

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence: Synthetic DNA

55

<400> 4

actcatcttg gaatctcaga attgg

25

5

<210> 5

<211> 24

10

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 5

20

cttgaccgtt tctatcttct ctcg

24

25

<210> 6

<211> 979

<212> DNA

<213> *Cricetulus griseus*

30

<400> 6

actcatcttg gaatctcaga attggcgcta tgctactgga ggatgggaga ctgtgttttag 60

35

acctgtaagt gagacatgca cagacaggtc tggcctctcc actggacact ggtcagggtga 120

agtgaaggac aaaaatgttc aagtggtcga gctccccatt gtagacagcc tccatcctcg 180

tctccttac ttacccttgg ctgtaccaga agaccttgca gatcgactcc tgagagtcca 240

tgggtgatcct gcagtgtggt gggatatcca gtttgtcaaa tacttgatcc gtccacaacc 300

40

ttggctggaa agggaaatag aagaaaccac caagaagctt ggcttcaaac atccagttat 360

tggagtccat gtcagacgca ctgacaaagt gggaacagaa gcagccttcc atcccattga 420

ggaatacatg gtacacgttg aagaacattt tcagcttctc gaacgcagaa tgaaagtgga 480

45

taaaaaaaga gtgtatctgg ccactgatga cccttctttg ttaaaggagg caaagacaaa 540

gtactccaat tatgaattta ttagtgataa ctctatitct tggtcagctg gactacacaa 600

ccgatacaca gaaaattcac ttcggggcgt gatcctggat atacacttcc tctcccaggc 660

tgacttcctt gtgtgtactt tttcatccca ggtctgtagg gttgcttatg aaatcatgca 720

50

aacactgcat cctgatgcct ctgcaaactt ccattcttta gatgacatct actattttgg 780

aggccaaaat gcccaaac agattgcagt ttatcctcac caacctcgaa cttaaaggga 840

aatcccatg gaacctggag atatcattgg tgtggctgga aaccattgga atggttactc 900

55

taaagggtgc aacagaaaac taggaaaaac aggcctgtac ccttcctaca aagtcaggga 960

gaagatagaa acggtcaag

979

5

<210> 7

<211> 979

<212> DNA

10

<213> Rattus norvegicus

<400> 7

15

actcatcttg gaatctcaga attggcgcta tgctactggt ggaatgggaga ctgtgttttag 60

acctgtaagt gagacatgca cagacagatc tggcctctcc actggacact ggtcagggtga 120

agtgaatgac aaaaatattc aagtgggtgga gctcccccatt gtagacagcc ttcatectcg 180

gcctccttac ttaccactgg ctgttcgaga agaccttgca gatcgactcg taagagtcca 240

20

tggtgatcct gcagtgtggt ggggtgtccca gttcgtcaaa tatttgattc gtccacaacc 300

tiggctagaa aaggaaatag aagaagccac caagaagctt ggcttcaaac atccagtcac 360

tggagtccat gtcagacgca cagacaaagt gggaacagag gcagccttcc atcccatcga 420

25

agagtacatg gtacatgttg aagaacattt tcagcttctc gcacgcagaa tgcaagtggga 480

taaaaaaaga gtatatctgg ctaccgatga cctgctttg ttaaaggagg caaagacaaa 540

gtactccaat tatgaattta ttagtgataa ctctatttct tggtcagctg gactacacaa 600

tcggtacaca gaaaattcac ttcggggcgt gatcctggat atacacttcc tctctcagge 660

30

tgacttccia gigtgtactt tttcatccca ggtctgtcgg gttgcttatg aaatcatgca 720

aaccctgcat cctgatgcct ctgcaaaactt ccactcttta gatgacatct actatttttg 780

aggccaaaat gccacacacc agattgccgt ttatcctcac aaacctcgaa ctgatgagga 840

35

aattccaatg gaacctggag atatcattgg tgtggctgga aaccattggg atggttattc 900

taaagggtgc aacagaaaac ttggaaaaac aggcttatat ccctctaca aagtcgaga 960

gaagatagaa acggtcaag 979

40

<210> 8

<211> 40

<212> DNA

45

<213> Artificial Sequence

<220>

50

<223> Description of Artificial Sequence: Synthetic DNA

<400> 8

55

aagtataagc ttacatggat gacgatatcg ctgcgctcgt

40

5 <210> 9
 <211> 40
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA
 15
 <400> 9
 atttaactgc aggaagcatt tgcggtggac gatggagggg 40
 20
 <210> 10
 <211> 40
 <212> DNA
 25 <213> Artificial Sequence
 <220>
 30 <223> Description of Artificial Sequence: Synthetic DNA
 <400> 10
 atttaaggta ccgaagcatt tgcggtgcac gatggagggg 40
 35
 <210> 11
 40 <211> 23
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA
 50
 <400> 11
 ctccaattat gaatttatta gtg 23
 55
 <210> 12

5
 <211> 25
 <212> DNA
 <213> Artificial Sequence

 10
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

 15
 <400> 12
 ggatgtttga agccaagctt ctgg 25

 20
 <210> 13
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 25
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

 30
 <400> 13
 gtccatgggtg atcctgcagt gtgg 24

 35
 <210> 14
 <211> 23
 <212> DNA
 40
 <213> Artificial Sequence

 45
 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

 50
 <400> 14
 caccaatgat atctccaggt tcc 23

 55
 <210> 15
 <211> 24
 <212> DNA

<213> Artificial Sequence

5

<220>

<223> Description of Artificial Sequence: Synthetic DNA

10

<400> 15

gatatcgctg cgctcgttgt cgac

24

15

<210> 16

<211> 24

<212> DNA

20

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

25

<400> 16

caggaaggaa ggctggaaaa gacg

24

30

<210> 17

<211> 24

35

<212> DNA

<213> Artificial Sequence

40

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 17

45

gatatcgctg cgctcgtcgt cgac

24

50

<210> 18

<211> 24

<212> DNA

<213> Artificial Sequence

55

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 18

caggaaggaa ggctggaaga gagc

24

<210> 19

<211> 321

<212> PRT

<213> Cricetulus griseus

<400> 19

Met Gly Glu Pro Gln Gly Ser Arg Arg Ile Leu Val Thr Gly Gly Ser
 1 5 10 15
 Gly Leu Val Gly Arg Ala Ile Gln Lys Val Val Ala Asp Gly Ala Gly
 20 25 30
 Leu Pro Gly Glu Glu Trp Val Phe Val Ser Ser Lys Asp Ala Asp Leu
 35 40 45
 Thr Asp Ala Ala Gln Thr Gln Ala Leu Phe Gln Lys Val Gln Pro Thr
 50 55 60
 His Val Ile His Leu Ala Ala Met Val Gly Gly Leu Phe Arg Asn Ile
 65 70 75 80
 Lys Tyr Asn Leu Asp Phe Trp Arg Lys Asn Val His Ile Asn Asp Asn
 85 90 95
 Val Leu His Ser Ala Phe Glu Val Gly Thr Arg Lys Val Val Ser Cys
 100 105 110
 Leu Ser Thr Cys Ile Phe Pro Asp Lys Thr Thr Tyr Pro Ile Asp Glu
 115 120 125
 Thr Met Ile His Asn Gly Pro Pro His Ser Ser Asn Phe Gly Tyr Ser
 130 135 140
 Tyr Ala Lys Arg Met Ile Asp Val Gln Asn Arg Ala Tyr Phe Gln Gln
 145 150 155 160
 His Gly Cys Thr Phe Thr Ala Val Ile Pro Thr Asn Val Phe Gly Pro
 165 170 175
 His Asp Asn Phe Asn Ile Glu Asp Gly His Val Leu Pro Gly Leu Ile
 180 185 190
 His Lys Val His Leu Ala Lys Ser Asn Gly Ser Ala Leu Thr Val Trp
 195 200 205

EP 1 498 491 A1

Gly Thr Gly Lys Pro Arg Arg Gln Phe Ile Tyr Ser Leu Asp Leu Ala
 210 215 220
 5 Arg Leu Phe Ile Trp Val Leu Arg Glu Tyr Asn Glu Val Glu Pro Ile
 225 230 235 240
 Ile Leu Ser Val Gly Glu Glu Asp Glu Val Ser Ile Lys Glu Ala Ala
 10 245 250 255
 Glu Ala Val Val Glu Ala Met Asp Phe Cys Gly Glu Val Thr Phe Asp
 260 265 270
 15 Ser Thr Lys Ser Asp Gly Gln Tyr Lys Lys Thr Ala Ser Asn Gly Lys
 275 280 285
 Leu Arg Ala Tyr Leu Pro Asp Phe Arg Phe Thr Pro Phe Lys Gln Ala
 290 295 300
 20 Val Lys Glu Thr Cys Ala Trp Phe Thr Asp Asn Tyr Glu Gln Ala Arg
 305 310 315 320
 Lys
 25
 <210> 20
 <211> 590
 30 <212> PRT
 <213> Cricetulus griseus
 <400> 20
 35 Met Ala Ser Leu Arg Glu Ala Ser Leu Arg Lys Leu Arg Arg Phe Ser
 1 5 10 15
 Glu Met Arg Gly Lys Pro Val Ala Thr Gly Lys Phe Trp Asp Val Val
 20 25 30
 40 Val Ile Thr Ala Ala Asp Glu Lys Gln Glu Leu Ala Tyr Lys Gln Gln
 35 40 45
 Leu Ser Glu Lys Leu Lys Arg Lys Glu Leu Pro Leu Gly Val Asn Tyr
 45 50 55 60
 His Val Phe Thr Asp Pro Pro Gly Thr Lys Ile Gly Asn Gly Gly Ser
 65 70 75 80
 50 Thr Leu Cys Ser Leu Gln Cys Leu Glu Ser Leu Tyr Gly Asp Lys Trp
 85 90 95
 Asn Ser Phe Thr Val Leu Leu Ile His Ser Gly Gly Tyr Ser Gln Arg
 100 105 110
 55 Leu Pro Asn Ala Ser Ala Leu Gly Lys Ile Phe Thr Ala Leu Pro Leu

EP 1 498 491 A1

| | | | |
|----|---|-----|-----|
| | 115 | 120 | 125 |
| 5 | Gly Glu Pro Ile Tyr Gln Met Leu Asp Leu Lys Leu Ala Met Tyr Met | | |
| | 130 | 135 | 140 |
| | Asp Phe Pro Ser Arg Met Lys Pro Gly Val Leu Val Thr Cys Ala Asp | | |
| | 145 | 150 | 155 |
| 10 | Asp Ile Glu Leu Tyr Ser Ile Gly Asp Ser Glu Ser Ile Ala Phe Glu | | |
| | 165 | 170 | 175 |
| | Gln Pro Gly Phe Thr Ala Leu Ala His Pro Ser Ser Leu Ala Val Gly | | |
| 15 | 180 | 185 | 190 |
| | Thr Thr His Gly Val Phe Val Leu Asp Ser Ala Gly Ser Leu Gln His | | |
| | 195 | 200 | 205 |
| | Gly Asp Leu Glu Tyr Arg Gln Cys His Arg Phe Leu His Lys Pro Ser | | |
| 20 | 210 | 215 | 220 |
| | Ile Glu Asn Met His His Phe Asn Ala Val His Arg Leu Gly Ser Phe | | |
| | 225 | 230 | 235 |
| 25 | Gly Gln Gln Asp Leu Ser Gly Gly Asp Thr Thr Cys His Pro Leu His | | |
| | 245 | 250 | 255 |
| | Ser Glu Tyr Val Tyr Thr Asp Ser Leu Phe Tyr Met Asp His Lys Ser | | |
| | 260 | 265 | 270 |
| 30 | Ala Lys Lys Leu Leu Asp Phe Tyr Glu Ser Val Gly Pro Leu Asn Cys | | |
| | 275 | 280 | 285 |
| | Glu Ile Asp Ala Tyr Gly Asp Phe Leu Gln Ala Leu Gly Pro Gly Ala | | |
| 35 | 290 | 295 | 300 |
| | Thr Ala Glu Tyr Thr Lys Asn Thr Ser His Val Thr Lys Glu Glu Ser | | |
| | 305 | 310 | 315 |
| | His Leu Leu Asp Met Arg Gln Lys Ile Phe His Leu Leu Lys Gly Thr | | |
| 40 | 325 | 330 | 335 |
| | Pro Leu Asn Val Val Val Leu Asn Asn Ser Arg Phe Tyr His Ile Gly | | |
| | 340 | 345 | 350 |
| | Thr Thr Glu Glu Tyr Leu Leu His Phe Thr Ser Asn Gly Ser Leu Gln | | |
| 45 | 355 | 360 | 365 |
| | Ala Glu Leu Gly Leu Gln Ser Ile Ala Phe Ser Val Phe Pro Asn Val | | |
| | 370 | 375 | 380 |
| 50 | Pro Glu Asp Ser His Glu Lys Pro Cys Val Ile His Ser Ile Leu Asn | | |
| | 385 | 390 | 395 |
| | Ser Gly Cys Cys Val Ala Pro Gly Ser Val Val Glu Tyr Ser Arg Leu | | |
| | 405 | 410 | 415 |
| 55 | Gly Pro Glu Val Ser Ile Ser Glu Asn Cys Ile Ile Ser Gly Ser Val | | |

EP 1 498 491 A1

420 425 430
 Ile Glu Lys Ala Val Leu Pro Pro Cys Ser Phe Val Cys Ser Leu Ser
 5 435 440 445
 Val Glu Ile Asn Gly His Leu Glu Tyr Ser Thr Met Val Phe Gly Met
 450 455 460
 10 Glu Asp Asn Leu Lys Asn Ser Val Lys Thr Ile Ser Asp Ile Lys Met
 465 470 475 480
 Leu Gln Phe Phe Gly Val Cys Phe Leu Thr Cys Leu Asp Ile Trp Asn
 485 490 495
 15 Leu Lys Ala Met Glu Glu Leu Phe Ser Gly Ser Lys Thr Gln Leu Ser
 500 505 510
 Leu Trp Thr Ala Arg Ile Phe Pro Val Cys Ser Ser Leu Ser Glu Ser
 20 515 520 525
 Val Ala Ala Ser Leu Gly Met Leu Asn Ala Ile Arg Asn His Ser Pro
 530 535 540
 Phe Ser Leu Ser Asn Phe Lys Leu Leu Ser Ile Gln Glu Met Leu Leu
 25 545 550 555 560
 Cys Lys Asp Val Gly Asp Met Leu Ala Tyr Arg Glu Gln Leu Phe Leu
 565 570 575
 30 Glu Ile Ser Ser Lys Arg Lys Gln Ser Asp Ser Glu Lys Ser
 580 585 590

35 <210> 21
 <211> 25
 <212> PRT
 40 <213> Homo sapiens

<400> 21
 Gln Val Thr Val Gln Ser Ser Pro Asn Phe Thr Gln His Val Arg Glu
 45 1 5 10 15
 Gln Ser Leu Val Thr Asp Gln Leu Cys
 20 25

50
 <210> 22
 <211> 32
 55 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 22

taaatagaat tcggcatcat gtggcagctg ct

32

<210> 23

<211> 575

<212> PRT

<213> Cricetulus griseus

<400> 23

Met Arg Ala Trp Thr Gly Ser Trp Arg Trp Ile Met Leu Ile Leu Phe

1 5 10 15

Ala Trp Gly Thr Leu Leu Phe Tyr Ile Gly Gly His Leu Val Arg Asp

20 25 30

Asn Asp His Pro Asp His Ser Ser Arg Glu Leu Ser Lys Ile Leu Ala

35 40 45

Lys Leu Glu Arg Leu Lys Gln Gln Asn Glu Asp Leu Arg Arg Met Ala

50 55 60

Glu Ser Leu Arg Ile Pro Glu Gly Pro Ile Asp Gln Gly Thr Ala Thr

65 70 75 80

Gly Arg Val Arg Val Leu Glu Glu Gln Leu Val Lys Ala Lys Glu Gln

85 90 95

Ile Glu Asn Tyr Lys Lys Gln Ala Arg Asn Asp Leu Gly Lys Asp His

100 105 110

Glu Ile Leu Arg Arg Arg Ile Glu Asn Gly Ala Lys Glu Leu Trp Phe

115 120 125

Phe Leu Gln Ser Glu Leu Lys Lys Leu Lys Lys Leu Glu Gly Asn Glu

130 135 140

Leu Gln Arg His Ala Asp Glu Ile Leu Leu Asp Leu Gly His His Glu

145 150 155 160

Arg Ser Ile Met Thr Asp Leu Tyr Tyr Leu Ser Gln Thr Asp Gly Ala

165 170 175

Gly Glu Trp Arg Glu Lys Glu Ala Lys Asp Leu Thr Glu Leu Val Gln

EP 1 498 491 A1

| | | | |
|----|---|-----|-----|
| | 180 | 185 | 190 |
| 5 | Arg Arg Ile Thr Tyr Leu Gln Asn Pro Lys Asp Cys Ser Lys Ala Arg | | |
| | 195 | 200 | 205 |
| | Lys Leu Val Cys Asn Ile Asn Lys Gly Cys Gly Tyr Gly Cys Gln Leu | | |
| | 210 | 215 | 220 |
| 10 | His His Val Val Tyr Cys Phe Met Ile Ala Tyr Gly Thr Gln Arg Thr | | |
| | 225 | 230 | 235 |
| | Leu Ile Leu Glu Ser Gln Asn Trp Arg Tyr Ala Thr Gly Gly Trp Glu | | |
| | 245 | 250 | 255 |
| 15 | Thr Val Phe Arg Pro Val Ser Glu Thr Cys Thr Asp Arg Ser Gly Leu | | |
| | 260 | 265 | 270 |
| | Ser Thr Gly His Trp Ser Gly Glu Val Lys Asp Lys Asn Val Gln Val | | |
| 20 | 275 | 280 | 285 |
| | Val Glu Leu Pro Ile Val Asp Ser Leu His Pro Arg Pro Pro Tyr Leu | | |
| | 290 | 295 | 300 |
| | Pro Leu Ala Val Pro Glu Asp Leu Ala Asp Arg Leu Leu Arg Val His | | |
| 25 | 305 | 310 | 315 |
| | Gly Asp Pro Ala Val Trp Trp Val Ser Gln Phe Val Lys Tyr Leu Ile | | |
| | 325 | 330 | 335 |
| 30 | Arg Pro Gln Pro Trp Leu Glu Arg Glu Ile Glu Glu Thr Thr Lys Lys | | |
| | 340 | 345 | 350 |
| | Leu Gly Phe Lys His Pro Val Ile Gly Val His Val Arg Arg Thr Asp | | |
| | 355 | 360 | 365 |
| 35 | Lys Val Gly Thr Glu Ala Ala Phe His Pro Ile Glu Glu Tyr Met Val | | |
| | 370 | 375 | 380 |
| | His Val Glu Glu His Phe Gln Leu Leu Glu Arg Arg Met Lys Val Asp | | |
| 40 | 385 | 390 | 395 |
| | Lys Lys Arg Val Tyr Leu Ala Thr Asp Asp Pro Ser Leu Leu Lys Glu | | |
| | 405 | 410 | 415 |
| | Ala Lys Thr Lys Tyr Ser Asn Tyr Glu Phe Ile Ser Asp Asn Ser Ile | | |
| 45 | 420 | 425 | 430 |
| | Ser Trp Ser Ala Gly Leu His Asn Arg Tyr Thr Glu Asn Ser Leu Arg | | |
| | 435 | 440 | 445 |
| 50 | Gly Val Ile Leu Asp Ile His Phe Leu Ser Gln Ala Asp Phe Leu Val | | |
| | 450 | 455 | 460 |
| | Cys Thr Phe Ser Ser Gln Val Cys Arg Val Ala Tyr Glu Ile Met Gln | | |
| | 465 | 470 | 475 |
| 55 | Thr Leu His Pro Asp Ala Ser Ala Asn Phe His Ser Leu Asp Asp Ile | | |

EP 1 498 491 A1

5 Tyr Tyr Phe Gly Gly Gln Asn Ala His Asn Gln Ile Ala Val Tyr Pro
 500 505 510
 His Gln Pro Arg Thr Lys Glu Glu Ile Pro Met Glu Pro Gly Asp Ile
 515 520 525
 10 Ile Gly Val Ala Gly Asn His Trp Asn Gly Tyr Ser Lys Gly Val Asn
 530 535 540
 Arg Lys Leu Gly Lys Thr Gly Leu Tyr Pro Ser Tyr Lys Val Arg Glu
 15 545 550 555 560
 Lys Ile Glu Thr Val Lys Tyr Pro Thr Tyr Pro Glu Ala Glu Lys
 565 570 575
 20
 <210> 24
 <211> 575
 25 <212> PRT
 <213> Mus musculus
 <400> 24
 30 Met Arg Ala Trp Thr Gly Ser Trp Arg Trp Ile Met Leu Ile Leu Phe
 1 5 10 15
 Ala Trp Gly Thr Leu Leu Phe Tyr Ile Gly Gly His Leu Val Arg Asp
 20 25 30
 35 Asn Asp His Pro Asp His Ser Ser Arg Glu Leu Ser Lys Ile Leu Ala
 35 40 45
 Lys Leu Glu Arg Leu Lys Gln Gln Asn Glu Asp Leu Arg Arg Met Ala
 40 50 55 60
 Glu Ser Leu Arg Ile Pro Glu Gly Pro Ile Asp Gln Gly Thr Ala Thr
 65 70 75 80
 Gly Arg Val Arg Val Leu Glu Glu Gln Leu Val Lys Ala Lys Glu Gln
 45 85 90 95
 Ile Glu Asn Tyr Lys Lys Gln Ala Arg Asn Gly Leu Gly Lys Asp His
 100 105 110
 50 Glu Ile Leu Arg Arg Arg Ile Glu Asn Gly Ala Lys Glu Leu Trp Phe
 115 120 125
 Phe Leu Gln Ser Glu Leu Lys Lys Leu Lys His Leu Glu Gly Asn Glu
 130 135 140
 55 Leu Gln Arg His Ala Asp Glu Ile Leu Leu Asp Leu Gly His His Glu

EP 1 498 491 A1

| | | | | |
|----|---|-----|-----|-----|
| | 145 | 150 | 155 | 160 |
| 5 | Arg Ser Ile Met Thr Asp Leu Tyr Tyr Leu Ser Gln Thr Asp Gly Ala | | | |
| | | 165 | 170 | 175 |
| | Gly Asp Trp Arg Glu Lys Glu Ala Lys Asp Leu Thr Glu Leu Val Gln | | | |
| | | 180 | 185 | 190 |
| 10 | Arg Arg Ile Thr Tyr Leu Gln Asn Pro Lys Asp Cys Ser Lys Ala Arg | | | |
| | | 195 | 200 | 205 |
| | Lys Leu Val Cys Asn Ile Asn Lys Gly Cys Gly Tyr Gly Cys Gln Leu | | | |
| 15 | | 210 | 215 | 220 |
| | His His Val Val Tyr Cys Phe Met Ile Ala Tyr Gly Thr Gln Arg Thr | | | |
| | 225 | | 230 | 235 |
| | Leu Ile Leu Glu Ser Gln Asn Trp Arg Tyr Ala Thr Gly Gly Trp Glu | | | |
| 20 | | 245 | 250 | 255 |
| | Thr Val Phe Arg Pro Val Ser Glu Thr Cys Thr Asp Arg Ser Gly Leu | | | |
| | | 260 | 265 | 270 |
| 25 | Ser Thr Gly His Trp Ser Gly Glu Val Asn Asp Lys Asn Ile Gln Val | | | |
| | | 275 | 280 | 285 |
| | Val Glu Leu Pro Ile Val Asp Ser Leu His Pro Arg Pro Pro Tyr Leu | | | |
| | | 290 | 295 | 300 |
| 30 | Pro Leu Ala Val Pro Glu Asp Leu Ala Asp Arg Leu Leu Arg Val His | | | |
| | 305 | | 310 | 315 |
| | Gly Asp Pro Ala Val Trp Trp Val Ser Gln Phe Val Lys Tyr Leu Ile | | | |
| | | 325 | 330 | 335 |
| 35 | Arg Pro Gln Pro Trp Leu Glu Lys Glu Ile Glu Glu Ala Thr Lys Lys | | | |
| | | 340 | 345 | 350 |
| | Leu Gly Phe Lys His Pro Val Ile Gly Val His Val Arg Arg Thr Asp | | | |
| 40 | | 355 | 360 | 365 |
| | Lys Val Gly Thr Glu Ala Ala Phe His Pro Ile Glu Glu Tyr Met Val | | | |
| | | 370 | 375 | 380 |
| | His Val Glu Glu His Phe Gln Leu Leu Ala Arg Arg Met Gln Val Asp | | | |
| 45 | | 385 | 390 | 395 |
| | Lys Lys Arg Val Tyr Leu Ala Thr Asp Asp Pro Thr Leu Leu Lys Glu | | | |
| | | 405 | 410 | 415 |
| 50 | Ala Lys Thr Lys Tyr Ser Asn Tyr Glu Phe Ile Ser Asp Asn Ser Ile | | | |
| | | 420 | 425 | 430 |
| | Ser Trp Ser Ala Gly Leu His Asn Arg Tyr Thr Glu Asn Ser Leu Arg | | | |
| | | 435 | 440 | 445 |
| 55 | Gly Val Ile Leu Asp Ile His Phe Leu Ser Gln Ala Asp Phe Leu Val | | | |

EP 1 498 491 A1

450 455 460
 Cys Thr Phe Ser Ser Gln Val Cys Arg Val Ala Tyr Glu Ile Met Gln
 5 465 470 475 480
 Thr Leu His Pro Asp Ala Ser Ala Asn Phe His Ser Leu Asp Asp Ile
 485 490 495
 10 Tyr Tyr Phe Gly Gly Gln Asn Ala His Asn Gln Ile Ala Val Tyr Pro
 500 505 510
 His Lys Pro Arg Thr Glu Glu Glu Ile Pro Met Glu Pro Gly Asp Ile
 515 520 525
 15 Ile Gly Val Ala Gly Asn His Trp Asp Gly Tyr Ser Lys Gly Ile Asn
 530 535 540
 Arg Lys Leu Gly Lys Thr Gly Leu Tyr Pro Ser Tyr Lys Val Arg Glu
 20 545 550 555 560
 Lys Ile Glu Thr Val Lys Tyr Pro Thr Tyr Pro Glu Ala Glu Lys
 565 570 575

25

<210> 25

<211> 18

30

<212> PRT

<213> Homo sapiens

<400> 25

35

Asp Glu Ser Ile Tyr Ser Asn Tyr Tyr Leu Tyr Glu Ser Ile Pro Lys

1

5

10

15

Pro Cys

40

<210> 26

<211> 34

45

<212> DNA

<213> Artificial Sequence

50

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 26

55

aataaaggat cctggggtca ttgtcttga gggt

34


```

5      <210> 27
      <211> 788
      <212> DNA
10     <213> Homo sapiens

      <220>
      <221> CDS
15     <222> (13)..(774)

      <400> 27
20     gaa ttc ggc atc atg tgg cag ctg ctc ctc cca act gct ctg cta ctt 48
          Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu
              1              5              10
25     cta gtt tca gct ggc atg cgg act gaa gat ctc cca aag gct gtg gtg 96
          Leu Val Ser Ala Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val
              15              20              25
30     ttc ctg gag cct caa tgg tac agg gtg ctc gag aag gac agt gtg act 144
          Phe Leu Glu Pro Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr
              30              35              40
35     ctg aag tgc cag gga gcc tac tcc cct gag gac aat tcc aca cag tgg 192
          Leu Lys Cys Gln Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp
              45              50              55              60
40     ttt cac aat gag agc ctc atc tca agc cag gcc tcg agc tac ttc att 240
          Phe His Asn Glu Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile
              65              70              75
45     gac gct gcc aca gtc gac gac agt gga gag tac agg tgc cag aca aac 288
          Asp Ala Ala Thr Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn
              80              85              90
50     ctc tcc acc ctc agt gac ccg gtg cag cta gaa gtc cat atc ggc tgg 336
          Leu Ser Thr Leu Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp
              95              100             105
55     ctg ttg ctc cag gcc cct cgg tgg gtg ttc aag gag gaa gac cct att 384
          Leu Leu Leu Gln Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile
              110             115             120
60     cac ctg agg tgt cac agc tgg aag aac act gct ctg cat aag gtc aca 432
          His Leu Arg Cys His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr

```

EP 1 498 491 A1

125 130 135 140
 5 tat tta cag aat ggc aaa ggc agg aag tat ttt cat cat aat tct gac 480
 Tyr Leu Gln Asn Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp
 145 150 155
 10 ttc tac att cca aaa gcc aca ctc aaa gac agc ggc tcc tac ttc tgc 528
 Phe Tyr Ile Pro Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys
 160 165 170
 15 agg ggg ctt ttt ggg agt aaa aat gtg tct tca gag act gtg aac atc 576
 Arg Gly Leu Phe Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile
 175 180 185
 acc atc act caa ggt ttg gca gtg tca acc atc tca tca ttc ttt cca 624
 Thr Ile Thr Gln Gly Leu Ala Val Ser Thr Ile Ser Ser Phe Phe Pro
 20 190 195 200
 cct ggg tac caa gtc tct ttc tgc ttg gtg atg gta ctc ctt ttt gca 672
 Pro Gly Tyr Gln Val Ser Phe Cys Leu Val Met Val Leu Leu Phe Ala
 25 205 210 215 220
 gtg gac aca gga cta tat ttc tct gtg aag aca aac att cga agc tca 720
 Val Asp Thr Gly Leu Tyr Phe Ser Val Lys Thr Asn Ile Arg Ser Ser
 225 230 235
 30 aca aga gac tgg aag gac cat aaa ttt aaa tgg aga aag gac cct caa 768
 Thr Arg Asp Trp Lys Asp His Lys Phe Lys Trp Arg Lys Asp Pro Gln
 240 245 250
 35 gac aaa tga ccc cag gat cc 788
 Asp Lys
 40 <210> 28
 <211> 254
 <212> PRT
 <213> Homo sapiens
 45
 <400> 28
 Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu Leu Val Ser Ala
 50 1 5 10 15
 Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro
 20 25 30
 55 Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr Leu Lys Cys Gln
 35 40 45

EP 1 498 491 A1

Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp Phe His Asn Glu
 50 55 60
 5 Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile Asp Ala Ala Thr
 65 70 75 80
 Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn Leu Ser Thr Leu
 85 90 95
 10 Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp Leu Leu Leu Gln
 100 105 110
 15 Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile His Leu Arg Cys
 115 120 125
 His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr Tyr Leu Gln Asn
 130 135 140
 20 Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp Phe Tyr Ile Pro
 145 150 155 160
 Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Phe
 165 170 175
 25 Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln
 180 185 190
 Gly Leu Ala Val Ser Thr Ile Ser Ser Phe Phe Pro Pro Gly Tyr Gln
 195 200 205
 30 Val Ser Phe Cys Leu Val Met Val Leu Leu Phe Ala Val Asp Thr Gly
 210 215 220
 Leu Tyr Phe Ser Val Lys Thr Asn Ile Arg Ser Ser Thr Arg Asp Trp
 225 230 235 240
 35 Lys Asp His Lys Phe Lys Trp Arg Lys Asp Pro Gln Asp Lys
 245 250
 40

<210> 29

<211> 51

45 <212> DNA

<213> Artificial Sequence

50 <220>

<223> Description of Artificial Sequence: Synthetic DNA

55 <400> 29

tggtggatcc tgtcaatgat gatgatgatg atgaccttga gtgatggtga t

51

5 <210> 30
<211> 620
<212> DNA
10 <213> Homo sapiens

<220>
15 <221> CDS
<222> (13).. (609)

<400> 30

20 gaa ttc ggc atc atg tgg cag ctg ctc ctc cca act gct ctg cta ctt 48
Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu
1 5 10
25 cta gtt tca gct ggc atg cgg act gaa gat ctc cca aag gct gtg gtg 96
Leu Val Ser Ala Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val
15 20 25
30 ttc ctg gag cct caa tgg tac agg gtg ctc gag aag gac agt gtg act 144
Phe Leu Glu Pro Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr
30 35 40
35 ctg aag tgc cag gga gcc tac tcc cct gag gac aat tcc aca cag tgg 192
Leu Lys Cys Gln Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp
45 50 55 60
40 ttt cac aat gag agc ctc atc tca agc cag gcc tcg agc tac ttc att 240
Phe His Asn Glu Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile
65 70 75
45 gac gct gcc aca gtc gac gac agt gga gag tac agg tgc cag aca aac 288
Asp Ala Ala Thr Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn
80 85 90
50 ctc tcc acc ctc agt gac ccg gtg cag cta gaa gtc cat atc ggc tgg 336
Leu Ser Thr Leu Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp
95 100 105
55 ctg ttg ctc cag gcc cct cgg tgg gtg ttc aag gag gaa gac cct att 384
Leu Leu Leu Gln Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile
110 115 120
cac ctg agg tgt cac agc tgg aag aac act gct ctg cat aag gtc aca 432
His Leu Arg Cys His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr

EP 1 498 491 A1

125 130 135 140
 tat tta cag aat ggc aaa ggc agg aag tat ttt cat cat aat tct gac 480
 5 Tyr Leu Gln Asn Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp
 145 150 155
 ttc tac att cca aaa gcc aca ctc aaa gac agc ggc tcc tac ttc tgc 528
 10 Phe Tyr Ile Pro Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys
 160 165 170
 agg ggg ctt ttt ggg agt aaa aat gtg tct tca gag act gtg aac atc 576
 15 Arg Gly Leu Phe Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile
 175 180 185
 acc atc act caa ggt cat cat cat cat cat cat tga cag gat cc 620
 20 Thr Ile Thr Gln Gly His His His His His His
 190 195

 25 <210> 31
 <211> 199
 <212> PRT
 <213> Homo sapiens
 30
 <400> 31
 Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu Leu Val Ser Ala
 1 5 10 15
 35 Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro
 20 25 30
 Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr Leu Lys Cys Gln
 40 35 40 45
 Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp Phe His Asn Glu
 50 55 60
 Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile Asp Ala Ala Thr
 45 65 70 75 80
 Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn Leu Ser Thr Leu
 85 90 95
 50 Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp Leu Leu Leu Gln
 100 105 110
 Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile His Leu Arg Cys
 115 120 125
 55 His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr Tyr Leu Gln Asn

EP 1 498 491 A1

130 135 140
 5 Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp Phe Tyr Ile Pro
 145 150 155 160
 Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Phe
 165 170 175
 10 Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln
 180 185 190
 Gly His His His His His His
 15 195

20 <210> 32
 <211> 24
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

30 <400> 32
 aggaagggtgg cgctcatcac gggc 24

35 <210> 33
 <211> 26
 <212> DNA
 40 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic DNA

45 <400> 33
 taaggccaca agtccttaatt gcatcc 26

50
 <210> 34
 <211> 27
 55

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 34

caggggtgtt cccttgagga ggtggaa

27

<210> 35

<211> 23

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 35

ccccacgc atgaagcctg gag

23

<210> 36

<211> 28

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 36

ggcaggagac caccttgcga gtgccac

28

<210> 37

<211> 28

<212> DNA

<213> Artificial Sequence

$\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 37

ggcgctggct taccggaga ggaatggg

28

<210> 38

<211> 28

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 38

aaaaggcctc agttagtga ctgtatgg

28

<210> 39

<211> 29

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 39

cgcggtatcct caagcgttgg ggttggtcc

29

<210> 40

<211> 45

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

5

<400> 40

cccaagcttg ccaccatggc tcacgctccc gctagctgcc cgagc

45

10

<210> 41

<211> 31

15

<212> DNA

<213> Artificial Sequence

<220>

20

<223> Description of Artificial Sequence: Synthetic DNA

<400> 41

ccggaattct gcccaagtatg agccatcctg g

25

31

<210> 42

30

<211> 17

<212> DNA

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 42

gccatccaga aggtggt

40

17

45

<210> 43

<211> 17

<212> DNA

50

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

55

<400> 43

gtcttgtcag ggaagat

17

<210> 44

<211> 28

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 44

ggcaggagac caccttgcga gtgcccac

28

<210> 45

<211> 28

⟨212⟩ DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 45

gggtgggctg taccttctgg aacagggc

28

<210> 46

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 46

ggcgctggct taccggaga ggaatggg

28

5

<210> 47

<211> 28

<212> DNA

10

<213> Artificial Sequence

<220>

15

<223> Description of Artificial Sequence: Synthetic DNA

<400> 47

20

ggaatgggtg tttgtctcctc caaagatgc

28

<210> 48

25

<211> 1316

<212> DNA

<213> *Cricetulus griseus*

30

<400> 48

35

gccccgcccc ctccacctgg accgagagta gctggagaat tgtgcaccgg aagtagctct 60
tggactgggtg gaacctgctg cagggtgcagc aacaatgggt gagccccagg gatccaggag 120
gatacctagt acagggggct ctggactgggt gggcagagct atccagaagg tggtcgcaga 180
tggcgctggc ttacccggag aggaatgggt gttgtctctc tccaaagatg cagatctgac 240
ggatgcagca caaaccctga cctgtttcca gaaggtacag cccacccatg tcatccatct 300
tgctgcaatg gtaggaggcc ttttcggaa tatcaaatac aacttggatt tctggaggaa 360
gaatgtgcac atcaatgaca acgtctgca ctacagcttc gaggtgggca ctgcgaaggt 420
ggtctcctgc ctgtccacct gtatcttccc tgacaagacc acctatccta ttgatgaaac 480
aatgatccac aatgggtccac cccacagcag caattttggg tactcgtatg ccaagaggat 540
gattgacgtg cagaacaggc ctaacttcca gcagcatggc tgcacettca ctgtgtcat 600
ccctaccaat gtctttggac ctcatgacaa cttcaacatt gaagatggcc atgtgtgcc 660
tggcctcatc cataagggtc atctggccaa gagtaatggt tcagccttga ctgtttgggg 720
tacagggaaa ccacggaggc agttcatcta ctactggac ctagcccggc tcttcatctg 780
ggtcctgcgg gagtacaatg aagttgagcc catcatcctc tcagtgggag aggaagatga 840
agtctccatt aaggaggcag ctgaggctgt agtggaggcc atggacttct gtggggaagt 900
cacttttgat tcaacaaagt cagatgggca gtataagaag acagccagca atggcaagct 960
tcgggcctac ttgcctgatt tccgtttcac acccttcaag caggctgtga aggagacctg 1020
tgccctggttc accgacaact atgagcaggc ccggaagtga agcatgggac aagcgggtgc 1080

55

tcagctggca atgccagtc agtaggtgc agtctcatca ttgcttgtc aagaactgag 1140
 5 gacagtatcc agcaacctga gccacatgct ggtctctctg ccagggggct tcatgcagcc 1200
 atccagtagg gcccatgttt gtccatcctc gggggaaggc cagaccaaca ccttgtttgt 1260
 ctgccttctg cccaacctca gtgcatccat gctggtcctg ctgtcccttg tctaga 1316

10

<210> 49

<211> 23

15

<212> DNA

<213> Artificial Sequence

<220>

20

<223> Description of Artificial Sequence: Synthetic DNA

<400> 49

25

gacacctgctg ggacaaaaat tgg

23

<210> 50

30

<211> 22

<212> DNA

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 50

40

cttaacatcc caagggatgc tg

22

45

<210> 51

<211> 1965

<212> DNA

50

<213> Cricetulus griseus

<400> 51

55

acggggggct cccggaagcg gggaccatgg cgtctctgcg cgaagcgagc ctgcggaagc 60

tgcggcgctt ttccgagatg agaggcaaac ctgtggcaac tgggaaattc tgggatgtag 120

5 ttgtaataac agcagctgac gaaaagcagg agcttgctta caagcaacag ttgtcggaga 180
 agctgaagag aaaggaattg ccccttggag ttaactacca tgttttcact gatcctcctg 240
 gaacccaaaat tggaaatgga ggatcaacac tttgttctct tcagtcctg gaaagcctct 300
 atggagacaa gtggaattcc ttcacagtcc tgtaattca ctctgggtggc tacagtcac 360
 10 gacttcccaa tgcaagcgt ttaggaaaaa tcttcacggc tttaccactt ggtgagccca 420
 tttatcagat gttggactta aaactagcca tgtacatgga tttccctca cgcatagaagc 480
 ctggagtttt ggtcacctgt gcagatgata ttgaactata cagcatiggg gactctgagt 540
 ccattgcatt tgagcagcct ggctttactg ccctagccca tccatctagt ctggctgtag 600
 15 gcaccacaca tggagtattt gtattggact ctgccggttc tttgcaacat ggtgacctag 660
 agtacaggca atgccaccgt ttctccata agcccagcat tgaaaacatg caccacttta 720
 atgccgtgca tagactagga agcttggtc aacaggactt gagtgggggt gacaccacct 780
 gtcattccatt gcactctgag tatgtctaca cagatagcct attttacatg gatcataaat 840
 20 cagccaaaaa gctacttgat ttctatgaaa gtgtaggccc actgaactgt gaaatagatg 900
 cctatggtga ctttctgcag gcactgggac ctggagcaac tgcagagtac accaagaaca 960
 cctcacacgt cactaaagag gaatcacact tgttggacat gaggcagaaa atattccacc 1020
 25 tcctcaaggg aacacccctg aatgttgttg tccttaataa ctccaggttt tatcacattg 1080
 gaacaacgga ggagtatctg ctacatttca ctccaatgg ttcgttacag gcagagctgg 1140
 gcttgcaatc catagcttcc agtgtcttc caaatgtgcc tgaagactcc catgagaaac 1200
 cctgtgtcat tcacagcatc ctgaattcag gatgtgtgt ggccccctggc tcagtggtag 1260
 30 aatattccag attaggacct gaggtgtcca tctcgaaaaa ctgcattatc agcggttctg 1320
 tcatagaaaa agctgttctg ccccatgtt ctttctgttg ctctttaagt gtggagataa 1380
 atggacactt agaatttca actatgtgt ttggcatgga agacaacttg aagaacagt 1440
 35 ttaaaacat atcagatata aagatgttc agttcttgg agtctgtttc ctgacttggt 1500
 tagatatttg gaaccttaaa gctatggaag aactattttc aggaagtaag acgcagctga 1560
 gcctgtggac tgctcgaatt ttccctgtct gttcttctct gagtgcagtc gttgcagcat 1620
 cccttgggat gttaaagcc attcgaaacc attcgccatt cagcctgagc aacttcaagc 1680
 40 tgctgtccat ccaggaaatg cttctctgca aagatgtagg agacatgctt gcttacaggg 1740
 agcaactctt tctagaaatc agttcaaaga gaaaacagtc tgattcggag aaatcttaaa 1800
 tacaatggat tttgcctgga aacaggattg caaatgcagg catattctat agatctctgg 1860
 gttcttcttt ctttctcccc tctctcttt ctttccctt tgatgtaatg acaaaggtaa 1920
 45 aaatggccac ttctgatgga aaaaaaaaaa aaaaaaaaaa aaaaaa 1965

50 <210> 52

<211> 27

<212> DNA

55 <213> Artificial Sequence

$\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 52

caggggtgtt cccttgagga ggtggaa

27

<210> 53

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 53

cactgagcca ggggccacac agcatcc

27

<210> 54

<211> 23

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

<400> 54

cccctcacgc atgaagcctg gag

23

<210> 55

<211> 27

<212> DNA

<213> Artificial Sequence

 $\langle 220 \rangle$

<223> Description of Artificial Sequence: Synthetic DNA

5 <400> 55
 tgccaccgtt tcctccataa gccccagc 27

10 <210> 56
 <211> 28
 <212> DNA
15 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

20 <400> 56
 atggctcaag ctcccgctaa gtgcccgga 28

25 <210> 57
 <211> 27
30 <212> DNA
 <213> Artificial Sequence

 <220>
35 <223> Description of Artificial Sequence: Synthetic DNA

 <400> 57
40 tcaagcgttt gggttggtcc tcatgag 27

45 <210> 58
 <211> 25
 <212> DNA
 <213> Artificial Sequence

50 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

55 <400> 58

| | | |
|----|---|----|
| | tccggggatg gcgagatggg caagc | 25 |
| 5 | | |
| | <210> 59 | |
| | <211> 24 | |
| 10 | <212> DNA | |
| | <213> Artificial Sequence | |
| 15 | <220> | |
| | <223> Description of Artificial Sequence: Synthetic DNA | |
| | <400> 59 | |
| 20 | cttgacatgg ctctgggctc caag | 24 |
| 25 | <210> 60 | |
| | <211> 25 | |
| | <212> DNA | |
| | <213> Artificial Sequence | |
| 30 | <220> | |
| | <223> Description of Artificial Sequence: Synthetic DNA | |
| 35 | <400> 60 | |
| | ccacttcagt cggtcggtag tattt | 25 |
| 40 | <210> 61 | |
| | <211> 24 | |
| | <212> DNA | |
| 45 | <213> Artificial Sequence | |
| | <220> | |
| 50 | <223> Description of Artificial Sequence: Synthetic DNA | |
| | <400> 61 | |
| 55 | cgctcaccgc cctgaggcga catg | 24 |

| | | |
|----|---|----|
| 5 | <210> 62 | |
| | <211> 32 | |
| | <212> DNA | |
| | <213> Artificial Sequence | |
| 10 | <220> | |
| | <223> Description of Artificial Sequence: Synthetic DNA | |
| 15 | <400> 62 | |
| | ggcagggtgct gtcggtgagg tcaccatagt gc | 32 |
| 20 | <210> 63 | |
| | <211> 24 | |
| | <212> DNA | |
| 25 | <213> Artificial Sequence | |
| | <220> | |
| 30 | <223> Description of Artificial Sequence: Synthetic DNA | |
| | <400> 63 | |
| 35 | ggggccatgc caaggactat gtgc | 24 |
| 40 | <210> 64 | |
| | <211> 25 | |
| | <212> DNA | |
| | <213> Artificial Sequence | |
| 45 | <220> | |
| | <223> Description of Artificial Sequence: Synthetic DNA | |
| 50 | <400> 64 | |
| | atgtggctga tgttacaaaa tgatg | 25 |
| 55 | <210> 65 | |

<211> 1504

<212> DNA

<213> *Cricetulus griseus*

<220>

<221> CDS

<222> (1).. (1119)

<400> 65

```

atg gct cac gct ccc gct agc tgc ccg agc tcc agg aac tct ggg gac 48
Met Ala His Ala Pro Ala Ser Cys Pro Ser Ser Arg Asn Ser Gly Asp
  1           5           10           15
ggc gat aag ggc aag ccc agg aag gtg gcg ctc atc acg ggc atc acc 96
Gly Asp Lys Gly Lys Pro Arg Lys Val Ala Leu Ile Thr Gly Ile Thr
      20           25           30
ggc cag gat ggc tca tac ttg gca gaa ttc ctg ctg gag aaa gga tac 144
Gly Gln Asp Gly Ser Tyr Leu Ala Glu Phe Leu Leu Glu Lys Gly Tyr
      35           40           45
gag gtt cat gga att gta cgg cga tcc agt tca ttt aat aca ggt cga 192
Glu Val His Gly Ile Val Arg Arg Ser Ser Ser Phe Asn Thr Gly Arg
      50           55           60
att gaa cat tta tat aag aat cca cag gct cat att gaa gga aac atg 240
Ile Glu His Leu Tyr Lys Asn Pro Gln Ala His Ile Glu Gly Asn Met
      65           70           75           80
aag ttg cac tat ggt gac ctc acc gac agc acc tgc cta gta aaa atc 288
Lys Leu His Tyr Gly Asp Leu Thr Asp Ser Thr Cys Leu Val Lys Ile
      85           90           95           100
atc aat gaa gtc aaa cct aca gag atc tac aat ctt ggt gcc cag agc 336
Ile Asn Glu Val Lys Pro Thr Glu Ile Tyr Asn Leu Gly Ala Gln Ser
      105           110           115
cat gtc aag att tcc ttt gac tta gca gag tac act gca gat gtt gat 384
His Val Lys Ile Ser Phe Asp Leu Ala Glu Tyr Thr Ala Asp Val Asp
      120           125           130
gga gtt ggc acc ttg cgg ctt ctg gat gca att aag act tgt ggc ctt 432
Gly Val Gly Thr Leu Arg Leu Leu Asp Ala Ile Lys Thr Cys Gly Leu
      135           140           145
ata aat tct gtg aag ttc tac cag gcc tca act agt gaa ctg tat gga 480
Ile Asn Ser Val Lys Phe Tyr Gln Ala Ser Thr Ser Glu Leu Tyr Gly

```

EP 1 498 491 A1

| | | | | |
|----|---|-----|-----|------|
| | 150 | 155 | 160 | |
| 5 | aaa gtg caa gaa ata ccc cag aaa gag acc acc cct ttc tat cca agg | | | 528 |
| | Lys Val Gln Glu Ile Pro Gln Lys Glu Thr Thr Pro Phe Tyr Pro Arg | | | |
| | 165 | 170 | 175 | 180 |
| 10 | tcg ccc tat gga gca gcc aaa ctt tat gcc tat tgg att gta gtg aac | | | 576 |
| | Ser Pro Tyr Gly Ala Ala Lys Leu Tyr Ala Tyr Trp Ile Val Val Asn | | | |
| | 185 | 190 | 195 | |
| 15 | ttt cga gag gct tat aat ctc ttt gcg gtg aac ggc att ctc ttc aat | | | 624 |
| | Phe Arg Glu Ala Tyr Asn Leu Phe Ala Val Asn Gly Ile Leu Phe Asn | | | |
| | 200 | 205 | 210 | |
| 20 | cat gag agt cct aga aga gga gct aat ttt gtt act cga aaa att agc | | | 672 |
| | His Glu Ser Pro Arg Arg Gly Ala Asn Phe Val Thr Arg Lys Ile Ser | | | |
| | 215 | 220 | 225 | |
| 25 | cgg tca gta gct aag att tac ctt gga caa ctg gaa tgt ttc agt ttg | | | 720 |
| | Arg Ser Val Ala Lys Ile Tyr Leu Gly Gln Leu Glu Cys Phe Ser Leu | | | |
| | 230 | 235 | 240 | |
| 30 | gga aat ctg gac gcc aaa cga gac tgg ggc cat gcc aag gac tat gtc | | | 768 |
| | Gly Asn Leu Asp Ala Lys Arg Asp Trp Gly His Ala Lys Asp Tyr Val | | | |
| | 245 | 250 | 255 | 260 |
| 35 | gag gct atg tgg ctg atg tta caa aat gat gaa cca gag gac ttt gtc | | | 816 |
| | Glu Ala Met Trp Leu Met Leu Gln Asn Asp Glu Pro Glu Asp Phe Val | | | |
| | 265 | 270 | 275 | |
| 40 | ata gct act ggg gaa gtt cat agt gtc cgt gaa ttt gtt gag aaa tca | | | 864 |
| | Ile Ala Thr Gly Glu Val His Ser Val Arg Glu Phe Val Glu Lys Ser | | | |
| | 280 | 285 | 290 | |
| 45 | ttc atg cac att gga aag acc att gtg tgg gaa gga aag aat gaa aat | | | 912 |
| | Phe Met His Ile Gly Lys Thr Ile Val Trp Glu Gly Lys Asn Glu Asn | | | |
| | 295 | 300 | 305 | |
| 50 | gaa gtg ggc aga tgt aaa gag acc ggc aaa att cat gtg act gtg gat | | | 960 |
| | Glu Val Gly Arg Cys Lys Glu Thr Gly Lys Ile His Val Thr Val Asp | | | |
| | 310 | 315 | 320 | |
| 55 | ctg aaa tac tac cga cca act gaa gtg gac ttc ctg cag gga gac tgc | | | 1008 |
| | Leu Lys Tyr Tyr Arg Pro Thr Glu Val Asp Phe Leu Gln Gly Asp Cys | | | |
| | 325 | 330 | 335 | 340 |
| 60 | tcc aag gcg cag cag aaa ctg aac tgg aag ccc cgc gtt gcc ttt gac | | | 1056 |
| | Ser Lys Ala Gln Gln Lys Leu Asn Trp Lys Pro Arg Val Ala Phe Asp | | | |
| | 345 | 350 | 355 | |
| 65 | gag ctg gtg agg gag atg gtg caa gcc gat gtg gag ctc atg aga acc | | | 1104 |

Glu Leu Val Arg Glu Met Val Gln Ala Asp Val Glu Leu Met Arg Thr

360 365 370

aac ccc aac gcc tga gcacctctac aaaaaaatc gcgagacatg gactatgggtg 1159

Asn Pro Asn Ala

375

cagagccagc caaccagagt ccagccactc ctgagaccat cgaccataaa ccctcgactg 1219

cctgtgtcgt cccacagct aagagctggg ccacagggtt gtgggcacca ggacggggac 1279

actccagagc taaggccact tcgcttttgt caaaggctcc tctcaatgat ttgaggaaat 1339

caagaagttt aaaatcacat actcatttta cttgaaatta tgtcactaga caacttaaat 1399

ttttgagtct tgagattgtt tttctctttt cttattaaat gatctttcta tgaccagca 1459

aaaaaaaaa aaaaaaggga tataaaaaa aaaaaaaaaa aaaaa 1504

<210> 66

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic DNA

<400> 66

atgaagttgc actatggtga cctca 25

<210> 67

<211> 59

<212> DNA

<213> Cricetulus griseus

<400> 67

ccgacagcac ctgcctagta aaaatcatca atgaagtcaa acctacagag atctacaat 59

<210> 68

<211> 25

<212> DNA

<213> Artificial Sequence

5 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

10 <400> 68
 gacttagcag agtacactgc agatg 25

15 <210> 69
 <211> 25
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> Description of Artificial Sequence: Synthetic DNA

25 <400> 69
 accttgata gaaaggggtg gtctc 25

30 <210> 70
 <211> 125
 <212> DNA
 35 <213> Cricetulus griseus

40 <400> 70
 ttgatggagt tggcaccitg cggttcttg atgcaattaa gacttgtggc cttataaatt 60
 ctgtgaagtt ctaccaggcc tcaactagtg aactgtatgg aaaagtgcaa gaaatacccc 120
 agaaa 125

45 <210> 71
 <211> 376
 50 <212> PRT
 <213> Cricetulus griseus

55 <400> 71
 Met Ala His Ala Pro Ala Ser Cys Pro Ser Ser Arg Asn Ser Gly Asp

EP 1 498 491 A1

| | | | | |
|----|---|-----|-----|-----|
| | 1 | 5 | 10 | 15 |
| 5 | Gly Asp Lys Gly Lys Pro Arg Lys Val Ala Leu Ile Thr Gly Ile Thr | | | |
| | 20 | 25 | 30 | |
| | Gly Gln Asp Gly Ser Tyr Leu Ala Glu Phe Leu Leu Glu Lys Gly Tyr | | | |
| | 35 | 40 | 45 | |
| 10 | Glu Val His Gly Ile Val Arg Arg Ser Ser Ser Phe Asn Thr Gly Arg | | | |
| | 50 | 55 | 60 | |
| | Ile Glu His Leu Tyr Lys Asn Pro Gln Ala His Ile Glu Gly Asn Met | | | |
| 15 | 65 | 70 | 75 | 80 |
| | Lys Leu His Tyr Gly Asp Leu Thr Asp Ser Thr Cys Leu Val Lys Ile | | | |
| | 85 | 90 | 95 | 100 |
| | Ile Asn Glu Val Lys Pro Thr Glu Ile Tyr Asn Leu Gly Ala Gln Ser | | | |
| 20 | 105 | 110 | 115 | |
| | His Val Lys Ile Ser Phe Asp Leu Ala Glu Tyr Thr Ala Asp Val Asp | | | |
| | 120 | 125 | 130 | |
| 25 | Gly Val Gly Thr Leu Arg Leu Leu Asp Ala Ile Lys Thr Cys Gly Leu | | | |
| | 135 | 140 | 145 | |
| | Ile Asn Ser Val Lys Phe Tyr Gln Ala Ser Thr Ser Glu Leu Tyr Gly | | | |
| | 150 | 155 | 160 | |
| 30 | Lys Val Gln Glu Ile Pro Gln Lys Glu Thr Thr Pro Phe Tyr Pro Arg | | | |
| | 165 | 170 | 175 | 180 |
| | Ser Pro Tyr Gly Ala Ala Lys Leu Tyr Ala Tyr Trp Ile Val Val Asn | | | |
| | 185 | 190 | 195 | |
| 35 | Phe Arg Glu Ala Tyr Asn Leu Phe Ala Val Asn Gly Ile Leu Phe Asn | | | |
| | 200 | 205 | 210 | |
| | His Glu Ser Pro Arg Arg Gly Ala Asn Phe Val Thr Arg Lys Ile Ser | | | |
| 40 | 215 | 220 | 225 | |
| | Arg Ser Val Ala Lys Ile Tyr Leu Gly Gln Leu Glu Cys Phe Ser Leu | | | |
| | 230 | 235 | 240 | |
| 45 | Gly Asn Leu Asp Ala Lys Arg Asp Trp Gly His Ala Lys Asp Tyr Val | | | |
| | 245 | 250 | 255 | 260 |
| | Glu Ala Met Trp Leu Met Leu Gln Asn Asp Glu Pro Glu Asp Phe Val | | | |
| | 265 | 270 | 275 | |
| 50 | Ile Ala Thr Gly Glu Val His Ser Val Arg Glu Phe Val Glu Lys Ser | | | |
| | 280 | 285 | 290 | |
| | Phe Met His Ile Gly Lys Thr Ile Val Trp Glu Gly Lys Asn Glu Asn | | | |
| | 295 | 300 | 305 | |
| 55 | Glu Val Gly Arg Cys Lys Glu Thr Gly Lys Ile His Val Thr Val Asp | | | |

EP 1 498 491 A1

310 315 320
 Leu Lys Tyr Tyr Arg Pro Thr Glu Val Asp Phe Leu Gln Gly Asp Cys
 5 325 330 335 340
 Ser Lys Ala Gln Gln Lys Leu Asn Trp Lys Pro Arg Val Ala Phe Asp
 345 350 355
 10 Glu Leu Val Arg Glu Met Val Gln Ala Asp Val Glu Leu Met Arg Thr
 360 365 370
 Asn Pro Asn Ala
 15 375

 <210> 72
 20 <211> 420
 <212> DNA
 <213> Mus musculus

 25 <220>
 <221> CDS
 <222> (1).. (420)

 30 <400> 72
 atg gaa tgg atc tgg atc ttt ctc ttc ttc ctc tca gga act aca ggt 48
 Met Glu Trp Ile Trp Ile Phe Leu Phe Phe Leu Ser Gly Thr Thr Gly
 35 1 5 10 15
 gtc tac tcc cag gtt cag ctg cag cag tct gga gct gag gtg gcg agg 96
 Val Tyr Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Val Ala Arg
 20 25 30
 40 ccc ggg gct tca gtg aaa ctg tcc tgc aag gct tct ggc tac acc ttc 144
 Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 35 40 45
 45 act gac tac tat cta aac tgg gtg aag cag agg tct gga cag ggc ctt 192
 Thr Asp Tyr Tyr Leu Asn Trp Val Lys Gln Arg Ser Gly Gln Gly Leu
 50 55 60
 50 gag tgg att gga gag att gat cct gga agt gat agt ata tat tat aat 240
 Glu Trp Ile Gly Glu Ile Asp Pro Gly Ser Asp Ser Ile Tyr Tyr Asn
 65 70 75 80
 gaa aac ttg gag ggc agg gcc aca ctg act gca gac aaa tcc tcc agc 288
 55 Glu Asn Leu Glu Gly Arg Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser

EP 1 498 491 A1

5 85 90 95
aca gcc tac atg cag ctc aac agc ctg aca tct gag gac tct gca gtc 336
Thr Ala Tyr Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val
 100 105 110
tat ttc tgt gca aga tat ggg tat tct aga tac gac gta agg ttt gtc 384
10 Tyr Phe Cys Ala Arg Tyr Gly Tyr Ser Arg Tyr Asp Val Arg Phe Val
 115 120 125
tac tgg ggc caa ggg act ctg gtc act gtc tct aca 420
15 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Thr
 130 135 140

20 <210> 73
 <211> 140
 <212> PRT
25 <213> Mus musculus

 <400> 73
Met Glu Trp Ile Trp Ile Phe Leu Phe Phe Leu Ser Gly Thr Thr Gly
30 1 5 10 15
Val Tyr Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Val Ala Arg
 20 25 30
Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe
35 35 40 45
Thr Asp Tyr Tyr Leu Asn Trp Val Lys Gln Arg Ser Gly Gln Gly Leu
 50 55 60
40 Glu Trp Ile Gly Glu Ile Asp Pro Gly Ser Asp Ser Ile Tyr Tyr Asn
 65 70 75 80
Glu Asn Leu Glu Gly Arg Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser
 85 90 95
45 Thr Ala Tyr Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val
 100 105 110
Tyr Phe Cys Ala Arg Tyr Gly Tyr Ser Arg Tyr Asp Val Arg Phe Val
50 115 120 125
Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Thr
 130 135 140

55

<210> 74

<211> 393

<212> DNA

<213> *Mus musculus*

<220>

<221> CDS

<222> (1)..(393)

<400> 74

```

atg aag ttg cct gtt agg ctg ttg gtg ctg atg ttc tgg att cct gct 48
Met Lys Leu Pro Val Arg Leu Leu Val Leu Met Phe Trp Ile Pro Ala
1           5           10          15
tcc agg agt gat gtt ttg atg acc caa act cca ctc tcc ctg cct gtc 96
Ser Arg Ser Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val
20          25          30
agt ctt gga gat caa gcc tcc atc tct tgc aga tct agt cag agt ctt 144
Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu
25          35          40          45
gta cat agt aat gga aga acc tat tta gaa tgg tac ctg cag aaa cct 192
Val His Ser Asn Gly Arg Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro
30          50          55          60
ggc cag tca cca aag gtc ctg atc tac aaa gtt tcc aac cga att tct 240
Gly Gln Ser Pro Lys Val Leu Ile Tyr Lys Val Ser Asn Arg Ile Ser
35          65          70          75          80
ggg gtc cca gac agg ttc agt ggc agt gga tca ggg aca gat ttc aca 288
Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
40          85          90          95
ctc aaa atc agc aga gtg gag gct gag gat ctg gga gtt tat ttc tgc 336
Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys
45          100         105         110
ttt cag ggt tca cat gtt ccg tac acg ttc gga ggg ggg acc aag ctg 384
Phe Gln Gly Ser His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu
50          115         120         125
gaa ata aaa 393
Glu Ile Lys
55          130

```

5 <210> 75
 <211> 131
 <212> PRT
 <213> Mus musculus
 10
 <400> 75
 Met Lys Leu Pro Val Arg Leu Leu Val Leu Met Phe Trp Ile Pro Ala
 15 1 5 10 15
 Ser Arg Ser Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val
 20 20 25 30
 Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu
 20 35 40 45
 Val His Ser Asn Gly Arg Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro
 50 55 60
 25 Gly Gln Ser Pro Lys Val Leu Ile Tyr Lys Val Ser Asn Arg Ile Ser
 65 70 75 80
 Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 85 90 95
 30 Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys
 100 105 110
 Phe Gln Gly Ser His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu
 115 120 125
 35 Glu Ile Lys
 130

40
 <210> 76
 <211> 5
 <212> PRT
 45 <213> Mus musculus

<400> 76
 50 Asp Tyr Tyr Leu Asn
 1 5

55 <210> 77

<211> 17

<212> PRT

<213> Mus musculus

<400> 77

Glu Ile Asp Pro Gly Ser Asp Ser Ile Tyr Tyr Asn Glu Asn Leu Glu

1

5

10

15

Gly

<210> 78

<211> 12

<212> PRT

<213> Mus musculus

<400> 78

Tyr Gly Tyr Ser Arg Tyr Asp Val Arg Phe Val Tyr

1

5

10

<210> 79

<211> 16

<212> PRT

<213> Mus musculus

<400> 79

Arg Ser Ser Gln Ser Leu Val His Ser Asn Gly Arg Thr Tyr Leu Glu

1

5

10

15

<210> 80

<211> 7

<212> PRT

<213> Mus musculus

<400> 80

Lys Val Ser Asn Arg Ile Ser

1

5

5 <210> 81
 <211> 9
 <212> PRT
 10 <213> Mus musculus

 <400> 81
 15 Phe Gln Gly Ser His Val Pro Tyr Thr
 1 5

 <210> 82
 20 <211> 22
 <212> DNA
 <213> Artificial Sequence

 25 <220>
 <223> Description of Artificial Sequence:synthetic DNA

 30 <400> 82
 ctgaattcgc ggccgctagt cc 22

 35 <210> 83
 <211> 39
 <212> DNA
 40 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence:synthetic DNA
 45 <400> 83
 atggggccctt ggtggaggct gtagagacag tgaccagag 39

 50

 <210> 84
 <211> 22
 55 <212> DNA

<213> Artificial Sequence
 5
 <220>
 <223> Description of Artificial Sequence:synthetic DNA
 10
 <400> 84
 ctgaattcgc ggccgctgct gt 22
 15
 <210> 85
 <211> 28
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 <223> Description of Artificial Sequence: synthetic DNA
 25
 <400> 85
 atcgtagctt ttatttcag cttggtcc 28
 30
 <210> 86
 <211> 57
 <212> DNA
 35
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence:synthetic DNA
 40
 <400> 86
 ttgttggtac cgaattcttt cagggccccc gagccccgag agcccaaata ttgtgac 57
 45
 <210> 87
 <211> 31
 <212> DNA
 <213> Artificial Sequence
 50
 <220>
 55

<223> Description of Artificial Sequence:synthetic DNA

5

<400> 87

gcgaattcca ccatgggcag cccccgctcc g

31

10

<210> 88

<211> 32

15

<212> DNA

<213> Artificial Sequence

<220>

20

<223> Description of Artificial Sequence:synthetic DNA

<400> 88

25

cgggatccct atcggggctc cggggcccaa gt

32

<210> 89

30

<211> 1341

<212> DNA

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence:synthetic DNA

40

<400> 89

atg ggc agc ccc cgc tcc gcg ctg agc tgc ctg ctg ttg cac ttg ctg 48

Met Gly Ser Pro Arg Ser Ala Leu Ser Cys Leu Leu Leu His Leu Leu

1

5

10

15

45

gtt ctc tgc ctc caa gcc cag gta act gtt cag tcc tca cct aat ttt 96

Val Leu Cys Leu Gln Ala Gln Val Thr Val Gln Ser Ser Pro Asn Phe

20

25

30

50

aca cag cat gtg agg gag cag agc ctg gtg acg gat cag ctc agc cgc 144

Thr Gln His Val Arg Glu Gln Ser Leu Val Thr Asp Gln Leu Ser Arg

35

40

45

55

cgc ctc atc cgg acc tac cag ctc tac agc cgc acc agc ggg aag cac 192

Arg Leu Ile Arg Thr Tyr Gln Leu Tyr Ser Arg Thr Ser Gly Lys His

EP 1 498 491 A1

| | | | | |
|----|---|-----|-----|-----|
| | 50 | 55 | 60 | |
| 5 | gtg cag gtc ctg gcc aac aag cgc atc aac gcc atg gca gaa gac gga | | | 240 |
| | Val Gln Val Leu Ala Asn Lys Arg Ile Asn Ala Met Ala Glu Asp Gly | | | |
| | 65 | 70 | 75 | 80 |
| | gac ccc ttc gcg aag ctc att gtg gag acc gat act ttt gga agc aga | | | 288 |
| 10 | Asp Pro Phe Ala Lys Leu Ile Val Glu Thr Asp Thr Phe Gly Ser Arg | | | |
| | 85 | 90 | 95 | |
| | gtc cga gtt cgc ggc gca gag aca ggt ctc tac atc tgc atg aac aag | | | 336 |
| 15 | Val Arg Val Arg Gly Ala Glu Thr Gly Leu Tyr Ile Cys Met Asn Lys | | | |
| | 100 | 105 | 110 | |
| | aag ggg aag cta att gcc aag agc aac ggc aaa ggc aag gac tgc gta | | | 384 |
| | Lys Gly Lys Leu Ile Ala Lys Ser Asn Gly Lys Gly Lys Asp Cys Val | | | |
| 20 | 115 | 120 | 125 | |
| | ttc aca gag atc gtg ctg gag aac aac tac acg gcg ctg cag aac gcc | | | 432 |
| | Phe Thr Glu Ile Val Leu Glu Asn Asn Tyr Thr Ala Leu Gln Asn Ala | | | |
| 25 | 130 | 135 | 140 | |
| | aag tac gag ggc tgg tac atg gcc ttt acc cgc aag ggc cgg ccc cgc | | | 480 |
| | Lys Tyr Glu Gly Trp Tyr Met Ala Phe Thr Arg Lys Gly Arg Pro Arg | | | |
| | 145 | 150 | 155 | 160 |
| 30 | aag ggc tcc aag acg cgc cag cat cag cgc gag gtg cac ttc atg aag | | | 528 |
| | Lys Gly Ser Lys Thr Arg Gln His Gln Arg Glu Val His Phe Met Lys | | | |
| | 165 | 170 | 175 | |
| | cgc ctg ccg cgg ggc cac cac acc acc gag cag agc ctg cgc ttc gag | | | 576 |
| 35 | Arg Leu Pro Arg Gly His His Thr Thr Glu Gln Ser Leu Arg Phe Glu | | | |
| | 180 | 185 | 190 | |
| | ttc ctc aac tac ccg ccc ttc acg cgc agc ctg cgc ggc agc cag agg | | | 624 |
| 40 | Phe Leu Asn Tyr Pro Pro Phe Thr Arg Ser Leu Arg Gly Ser Gln Arg | | | |
| | 195 | 200 | 205 | |
| | act tgg gcc ccg gag ccc cga gag ccc aaa tct tgt gac aaa act cac | | | 672 |
| | Thr Trp Ala Pro Glu Pro Arg Glu Pro Lys Ser Cys Asp Lys Thr His | | | |
| 45 | 210 | 215 | 220 | |
| | aca tgc cca ccg tgc cca gca cct gaa ctc ctg ggg gga ccg tca gtc | | | 720 |
| | Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val | | | |
| 50 | 225 | 230 | 235 | 240 |
| | ttc ctc ttc ccc cca aaa ccc aag gac acc ctc atg atc tcc cgg acc | | | 768 |
| | Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr | | | |
| | 245 | 250 | 255 | |
| 55 | cct gag gtc aca tgc gtg gtg gtg gac gtg agc cac gaa gac cct gag | | | 816 |

EP 1 498 491 A1

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
260 265 270
5 gtc aag ttc aac tgg tac gtg gac ggc gtg gag gtg cat aat gcc aag 864
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
275 280 285
10 aca aag ccg cgg gag gag cag tac aac agc acg tac cgt gtg gtc agc 912
Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
290 295 300
15 gtc ctc acc gtc ctg cac cag gac tgg ctg aat ggc aag gag tac aag 960
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
305 310 315 320
20 tgc aag gtc tcc aac aaa gcc ctc cca gcc ccc atc gag aaa acc atc 1008
Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
325 330 335
25 tcc aaa gcc aaa ggg cag ccc cga gaa cca cag gtg tac acc ctg ccc 1056
Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
340 345 350
cca tcc cgg gat gag ctg acc aag aac cag gtc agc ctg acc tgc ctg 1104
Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
355 360 365
30 gtc aaa ggc ttc tat ccc agc gac atc gcc gtg gag tgg gag agc aat 1152
Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
370 375 380
35 ggg cag ccg gag aac aac tac aag acc acg cct ccc gtg ctg gac tcc 1200
Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
385 390 395 400
40 gac ggc tcc ttc ttc ctc tac agc aag ctc acc gtg gac aag agc agg 1248
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
405 410 415
45 tgg cag cag ggg aac gtc ttc tca tgc tcc gtg atg cat gag gct ctg 1296
Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
420 425 430
50 cac aac cac tac acg cag aag agc ctc tcc ctg tct ccg ggt aaa 1341
His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
435 440 445
55 <210> 90

<211> 447

<212> protein

<213> homo sapiens

<400>90

10 Met Gly Ser Pro Arg Ser Ala Leu Ser Cys Leu Leu Leu His Leu Leu
 1 5 10 15
 Val Leu Cys Leu Gln Ala Gln Val Thr Val Gln Ser Ser Pro Asn Phe
 20 25 30
 15 Thr Gln His Val Arg Glu Gln Ser Leu Val Thr Asp Gln Leu Ser Arg
 35 40 45
 Arg Leu Ile Arg Thr Tyr Gln Leu Tyr Ser Arg Thr Ser Gly Lys His
 20 50 55 60
 Val Gln Val Leu Ala Asn Lys Arg Ile Asn Ala Met Ala Glu Asp Gly
 65 70 75 80
 25 Asp Pro Phe Ala Lys Leu Ile Val Glu Thr Asp Thr Phe Gly Ser Arg
 85 90 95
 Val Arg Val Arg Gly Ala Glu Thr Gly Leu Tyr Ile Cys Met Asn Lys
 100 105 110
 30 Lys Gly Lys Leu Ile Ala Lys Ser Asn Gly Lys Gly Lys Asp Cys Val
 115 120 125
 Phe Thr Glu Ile Val Leu Glu Asn Asn Tyr Thr Ala Leu Gln Asn Ala
 130 135 140
 35 Lys Tyr Glu Gly Trp Tyr Met Ala Phe Thr Arg Lys Gly Arg Pro Arg
 145 150 155 160
 Lys Gly Ser Lys Thr Arg Gln His Gln Arg Glu Val His Phe Met Lys
 165 170 175
 40 Arg Leu Pro Arg Gly His His Thr Thr Glu Gln Ser Leu Arg Phe Glu
 180 185 190
 Phe Leu Asn Tyr Pro Pro Phe Thr Arg Ser Leu Arg Gly Ser Gln Arg
 45 195 200 205
 Thr Trp Ala Pro Glu Pro Arg Glu Pro Lys Ser Cys Asp Lys Thr His
 210 215 220
 50 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 225 230 235 240
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 245 250 255
 55 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu

EP 1 498 491 A1

260 265 270
 5 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 275 280 285
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 290 295 300
 10 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 15 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 20 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 385 390 395 400
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415
 30 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 435 440 445
 35
 <210> 91
 40 <211> 1245
 <212> DNA
 <213> *Cricetulus griseus*
 45 <400> 91
 gaacttcacc caagccatgt gacaattgaa ggctgtaccc ccagacccta acatcttgga 60
 gccctgtaga ccaggagtg ctcttgccg tggggtgacc tagctcttct accaccatga 120
 50 acagggcccc tctgaagcgg tccaggatcc tgcgcatggc gctgactgga ggctccactg 180
 cctctgagga ggcagatgaa gacagcagga acaagccgtt tctgctgcgg gcgctgcaga 240
 tcgcgctggt cgtctctctc tactgggtca cctccatctc catggtattc ctcaacaagt 300
 acctgctgga cagcccctcc ctgcagctgg ataccctat ctctgctact ttctaccaat 360
 55 gccctgtgac ctctctgctg tgcaagggcc tcagcactct ggccacctgc tgccctggca 420

EP 1 498 491 A1

ccgttgactt cccacccctg aacctggacc ttaagggtggc ccgcagcgtg ctgccactgt 480
 cggtagtcct cattggcatg ataagtttca ataacctctg cctcaagtac gtaggggtgg 540
 5 ccttctacaa cgtggggcgc tcgtcacca ccgtgttcaa tgtgcttctg tcttacctgc 600
 tgctcaaaca gaccacttcc ttctatgccc tgctcacatg tggcatcatc attggtgggt 660
 tctggctggg tatagaccaa gagggagctg agggcaccct gtccctcata ggcaccatct 720
 10 tcgggggtgct ggccagcctc tgcgtctccc tcaatgccat ctataccaag aaggtgctcc 780
 cagcagtggc caacagcatc tggcgcctaa cttctataa caatgtcaat gcctgtgtgc 840
 tcttcttgcc cctgatggtt ctgctgggtg agctcctgc cctcctigac ttgctcatc 900
 15 tgtacagtgc ccacttctgg ctcatgatga cgctgggtgg cctcttcggc ttgccattg 960
 gctatgtgac aggactgcag atcaaattca ccagtcccct gaccacaaat gtatcaggca 1020
 cagccaaggc ctgtgcgcag acagtgtctg ccgtgctcta ctatgaagag actaagagct 1080
 tcctgtggtg gacaagcaac ctgatgtgct tgggtggctc ctacagctat acctgggtca 1140
 20 ggggctggga gatgcagaag acccaagagg accccagctc caaagagggt gagaagagt 1200
 ctattgggggt gtgagcttct tcaggacact gggactgaac ccaag 1245

25 <210> 92
 <211> 365
 <212> PRT
 30 <213> *Cricetulus griseus*

<400> 92
 Met Asn Arg Ala Pro Leu Lys Arg Ser Arg Ile Leu Arg Met Ala Leu
 35 1 5 10 15
 Thr Gly Gly Ser Thr Ala Ser Glu Glu Ala Asp Glu Asp Ser Arg Asn
 20 25 30
 40 Lys Pro Phe Leu Leu Arg Ala Leu Gln Ile Ala Leu Val Val Ser Leu
 35 40 45
 Tyr Trp Val Thr Ser Ile Ser Met Val Phe Leu Asn Lys Tyr Leu Leu
 50 55 60
 45 Asp Ser Pro Ser Leu Gln-Leu Asp Thr Pro Ile Phe Val Thr Phe Tyr
 65 70 75 80
 Gln Cys Leu Val Thr Ser Leu Leu Cys Lys Gly Leu Ser Thr Leu Ala
 50 85 90 95
 Thr Cys Cys Pro Gly Thr Val Asp Phe Pro Thr Leu Asn Leu Asp Leu
 100 105 110
 55 Lys Val Ala Arg Ser Val Leu Pro Leu Ser Val Val Phe Ile Gly Met
 115 120 125

EP 1 498 491 A1

Ile Ser Phe Asn Asn Leu Cys Leu Lys Tyr Val Gly Val Ala Phe Tyr
 130 135 140
 5 Asn Val Gly Arg Ser Leu Thr Thr Val Phe Asn Val Leu Leu Ser Tyr
 145 150 155 160
 Leu Leu Leu Lys Gln Thr Thr Ser Phe Tyr Ala Leu Leu Thr Cys Gly
 10 165 170 175
 Ile Ile Ile Gly Gly Phe Trp Leu Gly Ile Asp Gln Glu Gly Ala Glu
 180 185 190
 15 Gly Thr Leu Ser Leu Ile Gly Thr Ile Phe Gly Val Leu Ala Ser Leu
 195 200 205
 Cys Val Ser Leu Asn Ala Ile Tyr Thr Lys Lys Val Leu Pro Ala Val
 210 215 220
 20 Asp Asn Ser Ile Trp Arg Leu Thr Phe Tyr Asn Asn Val Asn Ala Cys
 225 230 235 240
 Val Leu Phe Leu Pro Leu Met Val Leu Leu Gly Glu Leu Arg Ala Leu
 245 250 255
 25 Leu Asp Phe Ala His Leu Tyr Ser Ala His Phe Trp Leu Met Met Thr
 260 265 270
 Leu Gly Gly Leu Phe Gly Phe Ala Ile Gly Tyr Val Thr Gly Leu Gln
 275 280 285
 30 Ile Lys Phe Thr Ser Pro Leu Thr His Asn Val Ser Gly Thr Ala Lys
 290 295 300
 Ala Cys Ala Gln Thr Val Leu Ala Val Leu Tyr Tyr Glu Glu Thr Lys
 35 305 310 315 320
 Ser Phe Leu Trp Trp Thr Ser Asn Leu Met Val Leu Gly Gly Ser Ser
 325 330 335
 40 Ala Tyr Thr Trp Val Arg Gly Trp Glu Met Gln Lys Thr Gln Glu Asp
 340 345 350
 Pro Ser Ser Lys Glu Gly Glu Lys Ser Ala Ile Gly Val
 355 360 365
 45

<210> 93

<211> 1095

<212> DNA

<213> Homo sapiens

<400> 93

atgaataggg cccctctgaa gcggtccagg atcctgcaca tggcgctgac cggggcctca 60
 gacccctctg cagaggcaga ggccaacggg gagaagccct ttctgctgag ggcattgcag 120
 5 atcgcgctgg tggctctccct ctacigggtc acctccatct ccatgggtgtt ccttaataag 180
 tacctgctgg acagcccctc cctgcggtg gacaccccca tcttcgtcac cttctaccag 240
 tgcctgggtga ccacgctgct gtgcaaaggc ctacgcgctc tggccgcctg ctgccctggt 300
 10 gccgtggact tccccagctt gcgcctggac ctacgggtgg cccgcagcgt cctgccccctg 360
 tcggtgggtct tcacggcat gatcaccttc aataacctct gectcaagta cgtcgggtgtg 420
 gccttctaca atgtgggccc ctactcacc accgtcttca acgtgctgct ctctacctg 480
 ctgctcaagc agaccacctc cttctatgcc ctgctcacct gcggtatcat catcgggggc 540
 15 ttctggcttg gtgtggacca ggagggggca gaaggcacc tgtcgtggct gggcaccgtc 600
 ttcggcgtgc tggctagcct ctgtgtctcg ctcaacgcca tctacaccac gaaggtgtc 660
 ccggcggtgg acggcagcat ctggcgctg acttcttaca acaacgtcaa cgcctgcac 720
 20 ctcttctctg cctgtctct gctgctcggg gagcttcagg cctgcgtga ctttggccag 780
 ctgggcagtg cccacttctg ggggatgatg acgtgggcg gcctgtttgg ctttgccatc 840
 ggctacgtga caggactgca gatcaagttc accagtcgc tgaccacaa tgtgtcgggc 900
 acggccaagg cctgtgccc gacagtgtg gccgtgtct actacgagga gaccaagagc 960
 25 ttcctctggt ggacgagcaa catgatggtg ctggcggtc cctccgcta cacctgggtc 1020
 aggggctggg agatgaagaa gactccggag gagcccagcc ccaaagacag cgagaagagc 1080
 gccatggggg tgtga 1095

30

<210> 94

<211> 364

35

<212> PRT

<213> Homo sapiens

40

<400> 94

Met Asn Arg Ala Pro Leu Lys Arg Ser Arg Ile Leu His Met Ala Leu

1 5 10 15

Thr Gly Ala Ser Asp Pro Ser Ala Glu Ala Glu Ala Asn Gly Glu Lys

45

20 25 30

Pro Phe Leu Leu Arg Ala Leu Gln Ile Ala Leu Val Val Ser Leu Tyr

35 40 45

Trp Val Thr Ser Ile Ser Met Val Phe Leu Asn Lys Tyr Leu Leu Asp

50

50 55 60

Ser Pro Ser Leu Arg Leu Asp Thr Pro Ile Phe Val Thr Phe Tyr Gln

65 70 75 80

55

Cys Leu Val Thr Thr Leu Leu Cys Lys Gly Leu Ser Ala Leu Ala Ala

EP 1 498 491 A1

| | | | | | | |
|----|---|-----|-----|-----|-----|-----|
| | | 85 | | 90 | | 95 |
| 5 | Cys Cys Pro Gly Ala Val Asp Phe Pro Ser Leu Arg Leu Asp Leu Arg | | | | | |
| | | 100 | | 105 | | 110 |
| | Val Ala Arg Ser Val Leu Pro Leu Ser Val Val Phe Ile Gly Met Ile | | | | | |
| | | 115 | | 120 | | 125 |
| 10 | Thr Phe Asn Asn Leu Cys Leu Lys Tyr Val Gly Val Ala Phe Tyr Asn | | | | | |
| | | 130 | | 135 | | 140 |
| | Val Gly Arg Ser Leu Thr Thr Val Phe Asn Val Leu Leu Ser Tyr Leu | | | | | |
| 15 | 145 | | 150 | | 155 | 160 |
| | Leu Leu Lys Gln Thr Thr Ser Phe Tyr Ala Leu Leu Thr Cys Gly Ile | | | | | |
| | | 165 | | 170 | | 175 |
| | Ile Ile Gly Gly Phe Trp Leu Gly Val Asp Gln Glu Gly Ala Glu Gly | | | | | |
| 20 | | 180 | | 185 | | 190 |
| | Thr Leu Ser Trp Leu Gly Thr Val Phe Gly Val Leu Ala Ser Leu Cys | | | | | |
| | | 195 | | 200 | | 205 |
| 25 | Val Ser Leu Asn Ala Ile Tyr Thr Thr Lys Val Leu Pro Ala Val Asp | | | | | |
| | | 210 | | 215 | | 220 |
| | Gly Ser Ile Trp Arg Leu Thr Phe Tyr Asn Asn Val Asn Ala Cys Ile | | | | | |
| | 225 | | 230 | | 235 | 240 |
| 30 | Leu Phe Leu Pro Leu Leu Leu Leu Leu Gly Glu Leu Gln Ala Leu Arg | | | | | |
| | | 245 | | 250 | | 255 |
| | Asp Phe Ala Gln Leu Gly Ser Ala His Phe Trp Gly Met Met Thr Leu | | | | | |
| | | 260 | | 265 | | 270 |
| 35 | Gly Gly Leu Phe Gly Phe Ala Ile Gly Tyr Val Thr Gly Leu Gln Ile | | | | | |
| | | 275 | | 280 | | 285 |
| | Lys Phe Thr Ser Pro Leu Thr His Asn Val Ser Gly Thr Ala Lys Ala | | | | | |
| 40 | | 290 | | 295 | | 300 |
| | Cys Ala Gln Thr Val Leu Ala Val Leu Tyr Tyr Glu Glu Thr Lys Ser | | | | | |
| | 305 | | 310 | | 315 | 320 |
| | Phe Leu Trp Trp Thr Ser Asn Met Met Val Leu Gly Gly Ser Ser Ala | | | | | |
| 45 | | 325 | | 330 | | 335 |
| | Tyr Thr Trp Val Arg Gly Trp Glu Met Lys Lys Thr Pro Glu Glu Pro | | | | | |
| | | 340 | | 345 | | 350 |
| 50 | Ser Pro Lys Asp Ser Glu Lys Ser Ala Met Gly Val | | | | | |
| | | 355 | | 360 | | 364 |
| 55 | <210> 95 | | | | | |

<211> 2609

<212> DNA

<213> Mus musculus

<400> 95

| | | |
|----|--|------|
| 10 | gagccgaggg tgggtgctgca ggtgcacccg agggcaccgc cgagggtgag caccaggtcc | 60 |
| | ctgcatcagc caggacacca gagcccagtc ggggtggacgg acgggcgcct ctgaagcgg | 120 |
| | ccaggatcct gcgcattggcg ctgactggag tctctgctgt ctccgaggag tcagagagcg | 180 |
| 15 | ggaacaagcc atttctgctc cgggctctgc agatcgcgct ggtggctctct ctctactggg | 240 |
| | tcacctccat ttccatggta ttccctcaaca agtacctgct ggacagcccc tccctgcagc | 300 |
| | tggatacccc catTTTTgtc accttctacc aatgcctggg gacctcactg ctgtgcaagg | 360 |
| | gcctcagcac tctggccacc tgcTgccccg gcattggtaga ctccccacc ctaaacctgg | 420 |
| 20 | acctcaaggt ggcccgaagt gtgctgccgc tgtcagtggg ctttatcggc atgataacct | 480 |
| | tcaataacct ctgcctcaag tacgtagggg tgccttcta caacgtggga cgctcgctca | 540 |
| | ccaccgtgtt caacgttctt ctctctacc tgcTgctcaa acagaccact tccttctatg | 600 |
| 25 | ccctgctcac ctgcggcgct atcatTgtg gtttctggct gggatatagac caagaaggag | 660 |
| | ctgaggggaac cttgtccctg acgggcacca tcttcggggg gctggccagc ctctgcgtct | 720 |
| | ccctcaatgc catctatacc aagaaggTgc tccctgcagt agaccacagt atctggcgcc | 780 |
| | taaccttcta taacaatgTc aatgcctgcg tgcTcttctt gccctgatg atagtgtctg | 840 |
| 30 | gcgagctccg tgccctctg gccttcaTc atctgagcag tgccacttc tggctcatga | 900 |
| | tgacgctggg tggcctgttt ggctttgcca tcggctatgt gacaggactg cagatcaaat | 960 |
| | tcaccagtcc cctgaccat aacgtgtcag gcacggccaa ggctgtgca cagacagtgc | 1020 |
| | tggccgtgct ctactacgaa gagattaaga gcttccTgtg gtggacaagc aacctgatgg | 1080 |
| 35 | tgttgggtgg ctccctccgc tacacctggg tcaggggctg ggagatgcag aagaccagg | 1140 |
| | aggacccag ctccaaagat ggtgagaaga gtgctatcag ggtgtgagct ccttcaggga | 1200 |
| | gccagggtg agctcgggtg ggcctgccc agcacggaag gcttccata gacctactg | 1260 |
| 40 | ggtatggccc tgagcaataa tgtttacatc cttctcagaa gacctctaa gaagagccag | 1320 |
| | gttctttcct gataatgtca gaaagctgcc aaatctcctg cctgccccat cttctagtct | 1380 |
| | tgggaaagcc ctaccaggag tggcacccTt cctgcctcct cctggggcct gtctacctcc | 1440 |
| | atatggTctc tgggttggg gccagctgca ctctttgggc actggactga tgaagtgatg | 1500 |
| 45 | tcttactttc tacacaaggg agatgggttg tgacctact atagctagtt gaaggagct | 1560 |
| | gtgtaacccc acatctctgg ggcctgggc aggtagcata atagctaggt gctattaaca | 1620 |
| | tcaataacac ttcagactac ctttgaggc agttgggagc tgagccgaga gagagagatg | 1680 |
| 50 | gccattctgc cctcttctgt gtggatgggt atgacagacc aactgtccat ggggtgactg | 1740 |
| | acacctccac acttcatatt ttcaacttta gaaaaggggg agccacacgt ttacagatt | 1800 |
| | aagtggagtg atgaatgcct ctacagcccc taacccact ttccctgcct ggcttctctt | 1860 |
| | ggcccagaag ggccaccatc ctgttctcca acacctgacc cagctatctg gctatactct | 1920 |
| 55 | ctttctgtac tcccttcccc ttccccccc cattagcctc ctcccaaca cctccatctt | 1980 |

EP 1 498 491 A1

5 caggcaggaa gtggggtcca ctcagcctct gttcccatct gcttggaccc ctgagcctct 2040
 catgaaggta ggcttatgtt ctctgaggct ggggccggag gagcgactg attctcggag 2100
 ttatcccatc aggcctcctgt cacaaaatag cctaggccgt gtgtctaaga acagtggagg 2160
 ttggcttata actgttctgg gggcagcgaa gccacatca aggtactcat agaccagta 2220
 10 tttctgagga aaccctgtc cacatctca cttggtaaag gcacagataa tctccctcag 2280
 gcctcttgta taggagcact agccctggga gggctccgcc ccatgacctg atcaccccaa 2340
 agccttcaac agaaggattc caacatgaat ttggggacag aagcactcag accacgatgc 2400
 ccagcaccac accctcctat cctcagggtg gctgtcactg tcctagtccc ttctgtttgg 2460
 15 ccttttgtac cctcatttcc ttggcgtcat gtttgatgic tttgtctctt togtgagaag 2520
 atggggaaac catgtcagcc tctgcttccg acttcccatg gggttctaag aagttgggtg 2580
 ggctgatgc cctgagttgt atgtgattt 2609

20
 <210> 96
 <211> 360
 <212> PRT
 25 <213> Mus musculus

<400> 96
 30 Met Ala Leu Thr Gly Val Ser Ala Val Ser Glu Glu Ser Glu Ser Gly
 1 5 10 15
 Asn Lys Pro Phe Leu Leu Arg Ala Leu Gln Ile Ala Leu Val Val Ser
 20 25 30
 35 Leu Tyr Trp Val Thr Ser Ile Ser Met Val Phe Leu Asn Lys Tyr Leu
 35 40 45
 Leu Asp Ser Pro Ser Leu Gln Leu Asp Thr Pro Ile Phe Val Thr Phe
 50 55 60
 40 Tyr Gln Cys Leu Val Thr Ser Leu Leu Cys Lys Gly Leu Ser Thr Leu
 65 70 75 80
 Ala Thr Cys Cys Pro Gly Met Val Asp Phe Pro Thr Leu Asn Leu Asp
 45 85 90 95
 Leu Lys Val Ala Arg Ser Val Leu Pro Leu Ser Val Val Phe Ile Gly
 100 105 110
 50 Met Ile Thr Phe Asn Asn Leu Cys Leu Lys Tyr Val Gly Val Pro Phe
 115 120 125
 Tyr Asn Val Gly Arg Ser Leu Thr Thr Val Phe Asn Val Leu Leu Ser
 130 135 140
 55 Tyr Leu Leu Leu Lys Gln Thr Thr Ser Phe Tyr Ala Leu Leu Thr Cys

EP 1 498 491 A1

145 150 155 160
 Gly Val Ile Ile Gly Gly Phe Trp Leu Gly Ile Asp Gln Glu Gly Ala
 5 165 170 175
 Glu Gly Thr Leu Ser Leu Thr Gly Thr Ile Phe Gly Val Leu Ala Ser
 180 185 190
 10 Leu Cys Val Ser Leu Asn Ala Ile Tyr Thr Lys Lys Val Leu Pro Ala
 195 200 205
 Val Asp His Ser Ile Trp Arg Leu Thr Phe Tyr Asn Asn Val Asn Ala
 210 215 220
 15 Cys Val Leu Phe Leu Pro Leu Met Ile Val Leu Gly Glu Leu Arg Ala
 225 230 235 240
 Leu Leu Ala Phe Thr His Leu Ser Ser Ala His Phe Trp Leu Met Met
 20 245 250 255
 Thr Leu Gly Gly Leu Phe Gly Phe Ala Ile Gly Tyr Val Thr Gly Leu
 260 265 270
 25 Gln Ile Lys Phe Thr Ser Pro Leu Thr His Asn Val Ser Gly Thr Ala
 275 280 285
 Lys Ala Cys Ala Gln Thr Val Leu Ala Val Leu Tyr Tyr Glu Glu Ile
 290 295 300
 30 Lys Ser Phe Leu Trp Trp Thr Ser Asn Leu Met Val Leu Gly Gly Ser
 305 310 315 320
 Ser Ala Tyr Thr Trp Val Arg Gly Trp Glu Met Gln Lys Thr Gln Glu
 325 330 335
 35 Asp Pro Ser Ser Lys Asp Gly Glu Lys Ser Ala Ile Arg Val
 340 345 350

40
 <210> 97
 <211> 1053
 <212> DNA
 45 <213> Rattus norvegiucus

<400> 97
 50 atggcgctga ctggagcctc tgctgtctct gaggaggcag acagcgagaa caagccattt 60
 ctgctacggg ctctgcagat cgcgctgggtg gtttctctct actgggtcac ctccatctcc 120
 atggtattcc tcaacaagta cctgctggac agccctctcc tgcagctgga taccctccatc 180
 ttcgtcacct tctaccaatg cctgggtgacc tcaactgctgt gcaagggcct cagcactctg 240
 55 gccacctgct gccctggcat ggtagacttc cccaccctaa acctggacct caaggtggcc 300

EP 1 498 491 A1

5 cgaagtgtgc tgccgctgtc cgtggctctt atcggaatga taaccttcaa taacctctgc 360
 ctcaagtagc tgggggtggc cttctacaac gtgggacgct cgtcactac cgtgttcaat 420
 gtgcttctct cctacctgct gcttaaacag accacttctt tttatgccct gctcacctgt 480
 gccatcatca ttggtgggtt ctggctggga atagatcaag agggagctga gggcaccctg 540
 tccctgacgg gcaccatctt cgggggtgctg gccagcctct gtgtctcact caatgccatc 600
 10 tacaccaaga aggtgctccc tgccgtagac cacagtatct ggcgccaac cttctataac 660
 aacgtcaacg cctgtgtgct cttcttgccc ctgatgtag tgctgggcga gctccatgct 720
 ctctggcct tcgctcatct gaacagcgcc cacttctggg tcatgatgac gctgggtgga 780
 ctcttcggct ttgccattgg ctatgtgaca ggactgcaga tcaaattcac cagtccccctg 840
 15 acccataatg tgtcgggcac agccaaggcc tgtgcacaga cagtgtggtg tgtgctctac 900
 tatgaagaga ttaagagctt cctgtggtgg acaagcaact tgatggtgct ggtgtgctcc 960
 tctgcctaca cctgggtcag gggctgggag atgcagaaga cccaggagga cccagctcc 1020
 20 aaagagggtg agaagagtgc tatcggggtg tga 1053

25 <210> 98
 <211> 350
 <212> PRT
 <213> Rattus norvegicus

30 <400> 98
 Met Ala Leu Thr Gly Ala Ser Ala Val Ser Glu Glu Ala Asp Ser Glu
 1 5 10 15
 35 Asn Lys Pro Phe Leu Leu Arg Ala Leu Gln Ile Ala Leu Val Val Ser
 20 25 30
 Leu Tyr Trp Val Thr Ser Ile Ser Met Val Phe Leu Asn Lys Tyr Leu
 35 40 45
 40 Leu Asp Ser Pro Ser Leu Gln Leu Asp Thr Pro Ile Phe Val Thr Phe
 50 55 60
 Tyr Gln Cys Leu Val Thr Ser Leu Leu Cys Lys Gly Leu Ser Thr Leu
 45 65 70 75 80
 Ala Thr Cys Cys Pro Gly Met Val Asp Phe Pro Thr Leu Asn Leu Asp
 85 90 95
 50 Leu Lys Val Ala Arg Ser Val Leu Pro Leu Ser Val Val Phe Ile Gly
 100 105 110
 Met Ile Thr Phe Asn Asn Leu Cys Leu Lys Tyr Val Gly Val Ala Phe
 115 120 125
 55 Tyr Asn Val Gly Arg Ser Leu Thr Thr Val Phe Asn Val Leu Leu Ser

EP 1 498 491 A1

130 135 140
 Tyr Leu Leu Leu Lys Gln Thr Thr Ser Phe Tyr Ala Leu Leu Thr Cys
 5 145 150 155 160
 Ala Ile Ile Ile Gly Gly Phe Trp Leu Gly Ile Asp Gln Glu Gly Ala
 165 170 175
 10 Glu Gly Thr Leu Ser Leu Thr Gly Thr Ile Phe Gly Val Leu Ala Ser
 180 185 190
 Leu Cys Val Ser Leu Asn Ala Ile Tyr Thr Lys Lys Val Leu Pro Ala
 195 200 205
 15 Val Asp His Ser Ile Trp Arg Leu Thr Phe Tyr Asn Asn Val Asn Ala
 210 215 220
 Cys Val Leu Phe Leu Pro Leu Met Val Val Leu Gly Glu Leu His Ala
 20 225 230 235 240
 Leu Leu Ala Phe Ala His Leu Asn Ser Ala His Phe Trp Val Met Met
 245 250 255
 25 Thr Leu Gly Gly Leu Phe Gly Phe Ala Ile Gly Tyr Val Thr Gly Leu
 260 265 270
 Gln Ile Lys Phe Thr Ser Pro Leu Thr His Asn Val Ser Gly Thr Ala
 275 280 285
 30 Lys Ala Cys Ala Gln Thr Val Leu Ala Val Leu Tyr Tyr Glu Glu Ile
 290 295 300
 Lys Ser Phe Leu Trp Trp Thr Ser Asn Leu Met Val Leu Gly Gly Ser
 305 310 315 320
 35 Ser Ala Tyr Thr Trp Val Arg Gly Trp Glu Met Gln Lys Thr Gln Glu
 325 330 335
 Asp Pro Ser Ser Lys Glu Gly Glu Lys Ser Ala Ile Gly Val
 340 345 350
 40

<210> 99

45 <211> 7752

<212> DNA

<213> Mus musculus

50 <400> 99

agcgttgcaa gttcagccga gggtggtgct gcaggtgcac ccgagggcac cgccgagggt 60
 gagcaccagg tccctgcac agccaggaca ccagagccca gtcgggtgga cggacggtac 120
 55 gttctggaag ggaaagggcc ccgggaaggg gatacagcat tgagaagctc agaggctttg 180

| | | |
|----|--|------|
| | gtctgggcat ccagaatgga ctttatctgg aggaggtgac atggcttctc gcctctggaa | 240 |
| 5 | gtgctgcttg gatctccggg acttcatgtc ctgactaggt ctggaagcgg tgaatatagg | 300 |
| | ggtaggaaaa aaggagagga ctgcaacaag gtcttcccga gtggcctgag ctcgagggac | 360 |
| | gagggaggtg caacgggtgg gagccgggag caagggtggt gcggagggag ggggggggtc | 420 |
| | tccctaagca gaaagggtgt attccatttt ctgggtagat ggtgaagatg cacctgaccg | 480 |
| 10 | agtctggctg atctgaagat atcaggggaa aagatagtgc ggtgaggagg gagaatgaca | 540 |
| | gaaccttcca gaaaatggga gaggctatag cacttgcaaa ccttccctg atctccgggg | 600 |
| | actcccggaa gaagagggca ggtctgtggg cataggtgca gacttgccgg ggagctcttg | 660 |
| 15 | acggccgcgg gaagtggcaa cggcctgcga gctggccctt taaggcggct cgtaggcgtg | 720 |
| | tcaggaaatg cgcgcagggc ccgcctgct cggttaagtgg ccggggaccc gcgtcgctga | 780 |
| | gccggaactt gaattcggct cgtggcaacc gcagggcctt gctccggtca ggccctgtc | 840 |
| | cgtgtccctc gagacgcctt cctgagcctc ggtgatctcc ctgcagcacg ccttcccttc | 900 |
| 20 | ggctctgcgg gtgcttccgg gggttccgc agcccatgct tcccacgcgg tccgcgtcca | 960 |
| | gttatttctt cctccgctcc gtcttctctt cgtctctcgc ctctcttctt ccttgcgact | 1020 |
| | cacgtgtccc ctgtctcaa actggccatg gctgtcaaag cccacatcct tagttaggcc | 1080 |
| 25 | ccttctccct tccctgggtc ttgtttcgtg acaccacctc cctcccccgc cccgggagcg | 1140 |
| | agcaaatga ggagcgggtg acctcggaac atccgggaagc agaacttcat ccaagaagga | 1200 |
| | ggggaccgat aggtcatccc atgtgacagt tgaaggctgc agccacagac cctagctgct | 1260 |
| | tgaagccctg tagtccaggg actgcttctg gccgtaagggt gaccagctc ttctgccacc | 1320 |
| 30 | atgaacaggg cgcctctgaa gcggtccagg atcctgcgca tggcgctgac tggagtctct | 1380 |
| | gctgtctccg aggagtcaga gagcgggaac aagccatttc tgctccgggc tctgcagatc | 1440 |
| | gcgctggtgg tctctctcta ctgggtcacc tccatttcca tggtaattct caacaagtac | 1500 |
| | ctgctggaca gcccctccct gcagctggat acccccattt ttgtcacctt ctaccaatgc | 1560 |
| 35 | ctgggtgacct cactgctgtg caagggcctc agcaactctg ccacctgtg ccccgcatg | 1620 |
| | gtagacttcc ccaccctaaa cctggacctc aagggtggccc gaagtgtgct gccgctgtca | 1680 |
| | gtggtcttta tcggcatgat aaccttcaat aacctctgcc tcaagtacgt aggggtgccc | 1740 |
| 40 | ttctacaacg tgggacgctc gctcaccacc gtgttcaacg ttcttctctc ctacctgctg | 1800 |
| | ctcaaacaga ccacttccct ctatgccctg ctacactgcg gcgtcatcat tggtagtggtg | 1860 |
| | gactgggggc gtggggagca ggaatcgtaa agatcaatac cacattactc attatctgtc | 1920 |
| | ccagggtcttt tgcaccacca gtcataggga gagacctgta gagaacaaat aacttcttta | 1980 |
| 45 | ctgtgactca gtaagttagg gatccagcca aggtgaacat aataatgtta ggcagacact | 2040 |
| | acagcaaagc cagccagaca ctcagatcta gctaagcatt tgagccatgt taatgtaacg | 2100 |
| | gatccccatt acaaggatata atatagctgc gttttaigga gagaaaccca aggcacagag | 2160 |
| 50 | aagctaagta gctgggatca cacaggtaat cactgacgta gcagaaattt gcacataagc | 2220 |
| | agttacctcc ataggttaca ctcttgacca acacaccact gttctcaaga ggtcaagggt | 2280 |
| | gaactcaggt catcacaatt ggcacaagta cctctacca ctgagccatt tcagtgtgtc | 2340 |
| | agtcaatatg tgtgtgcttt aagaggcttt aactaccttc tcagatgtga ccataagtaa | 2400 |
| 55 | ttaattaccg ataggagcgt tgtgtctgac attacacttg tagcatcctc tttattgtac | 2460 |

| | | |
|----|--|------|
| | ccataagctc tctgagtggc ggcatctctg tgaactgca gctcggagag gctgcgctcc | 2520 |
| 5 | ttgccacagc cccacaacta agaagcagat agtctgggac gcagtcccca gttggtcata | 2580 |
| | ctccctggcc tgtgtttcaa gccagctctg tttgctcctg acccttggga gttagcgcaa | 2640 |
| | tgaaaaccaa cactatcact acagtctaaa tgtgctttta aatgaaagcc caggaaacttt | 2700 |
| 10 | gaagcatccg gccccttaac ggcagccact atgtcttgat tccgccaaca tcttttcagt | 2760 |
| | gccccggcagt cacatggagc aagggcctct tggcttggac agcatgtgtt agggaaactg | 2820 |
| | tttgccactt tgaatgaatt tagtggctgc tgggttacag agaccagggc atctttcccc | 2880 |
| | tcagagtcct gaatgaacga aaagcaacct tcatttgtac ctgctctgga ttttagttcg | 2940 |
| 15 | tcttgttttg cctatttaga tgtccctggg gtctctgagg cccaggctgg gtgctctaga | 3000 |
| | tgtagggacc aggccaacct gtactgtctt ccctagaaac attgccctgg ttgggcagct | 3060 |
| | cctggatcca gggitaaggg gtctgggagg agagaggtca gatagtggca ggatgcctcc | 3120 |
| 20 | cactgcccc acatacatac cctaagagat ctggtactcc tccttcagc ctacaagcta | 3180 |
| | ccgtggggtc ccacttcagt ccaccagccc tgccaacgtt agaggggatg ggcctcctag | 3240 |
| | taggagaact tacatgcagg aaggtacagt ctctggagaa cctgagcccg ggtcccaaa | 3300 |
| | gggacaagta gctgatagtg aggcagctga gccccatggc ggccctgcca agtggcacgg | 3360 |
| 25 | gaaagtggag ctctctgctg cccccactac tggccccatc tcttggctct cccctccctt | 3420 |
| | cctcctgttg agaaggccca tctctggaaa ggcctcctag acatgcggca ctttgcaaag | 3480 |
| | cctgtcgggc tcacagcccc tctagggctt aggcacctga gaatgaagaa tggagtcaact | 3540 |
| | tctagactct agtggttaacc accaggaggt acagggtgct ctgactgtgc agggaaacct | 3600 |
| 30 | accgtgggct ctgctgagcc aagtgcctgt gaggctggag agtctggtgc ccttgttctg | 3660 |
| | agatagcatc ttgctatgtt gccctcaagt ccaggcaac tggggctgca ggagcaccac | 3720 |
| | cttgccctctc tccagcttct tgaagacttg tacctttctc ctagcagctc ctatctgctc | 3780 |
| | tcactccatc cattgagcag ctattagctt gtggccaagt attttccagg ccctgtactg | 3840 |
| 35 | agttttaggg tacaagtttg agaaaggaag ggtggggctc ttgctcctgg tccgtgaatg | 3900 |
| | atgttgatgg cagaaacgat agttacacta gatgctaagg gctgtgggta tctagaggga | 3960 |
| | gcagggagca tgtgggataa cctgagcagg cctagctgaa aagtcattgc tggcatgaga | 4020 |
| 40 | ctgctccagt agtacaggct gggaacacac atttgaatgt ttccctgaaga cagtgggag | 4080 |
| | ccacaggaaa tatccactgt agaaagatta tttagtgtgta agacagagta gtagatttgt | 4140 |
| | taacatagta gcaaaaacgt ggccccagtt tttacagatg aagggaattg gaactcagag | 4200 |
| | aggtttaagta acttctccca agcagctcag ctacaaaaat cacagaacag gcaggggcct | 4260 |
| 45 | gatggctctg atgcctgtgc tgggtccact attccatgtt gctaattcct gcagcagcag | 4320 |
| | taaacctctg ccttgtggaa atgaggagtc taaataaaga gaccatagca ttgccacaag | 4380 |
| | caggtttcta ccaactgggg gtggcaagga atgctgtgtt agcagcagga agctgggaag | 4440 |
| 50 | aggctgagta ctggggggat gaggaaggga tccccaggag aggctgactt tggccttgaa | 4500 |
| | gaatggtgga gtccctggaa agatgcagat acacagagct ctgtggatat acagagaagt | 4560 |
| | ggggagctaa gtaggtggct tggggccatc atgtgacaga ggaagtctggg ctagatgcag | 4620 |
| | gaagcccggg gctgtggcct agggagccat gtaggttctt tgagcagggg gcgggggggg | 4680 |
| 55 | gggggggtga cccaggagtg actgtaaaca acatcaggcc atgagcagct ctgacctaat | 4740 |

| | | |
|----|--|------|
| | gttctcacca agggagccag aaccaaggct tagagccctg tcccttttta ggtccaagg | 4800 |
| | tcactttact ggccctcttc ctttacagct gttggccccc acaggccatc aggcacctat | 4860 |
| 5 | gctattttat tttatagcct tcattacaat gactacaatt gtaattagag agttgacagg | 4920 |
| | gtcacatcig tccttatata ttcccccctc gctaagttct gcctgggaga atgtggaggg | 4980 |
| | tattggtgaa atttggggaa gttataacc cccaccccct gccccaccc cctgctttgc | 5040 |
| 10 | tccctttatc tgcagggcac ttctgtgccc actttagccc atatagctcc caaataaatg | 5100 |
| | acacagaaac ctggtatttt cattaacaaa ctgctggcac tctgctgggc aggttctgag | 5160 |
| | ctgttctaac cctctaagct gctaataccc agatagatgc cccaatgctt gccatccgag | 5220 |
| 15 | tccttctctg gcttgcctc ctccatgtgt gacctcatgg tgaatcctcc tgatttcccc | 5280 |
| | acatggcctc tccacacttt tccttctccc ctctctctac cagggaccct ctactggga | 5340 |
| | cccgatgtcc catctgtact gtcctctccc acccagtcac aggtctgattg agtctttatt | 5400 |
| | aaccaatcag agatgatgga aaaacagttt ttacatagca ctgaggatgg agatgcttga | 5460 |
| 20 | cccttgagat gcttggccgt aacctgtact gtatccagat gtctgggccc ccaaatcagc | 5520 |
| | aggtgaatac acagtacaca ggactgaccc ccaacagagg gggaacacag gttctcactc | 5580 |
| | tgggctccac gccctcggcc ctttcttagt gcaggggtta gactttgtat gtgttgatga | 5640 |
| 25 | tgaggtaagg gccatggaac agtcagaacg gtggtgtcag aatcctgtcc ctctccctcc | 5700 |
| | tgctctcact cctccttacc gtgtcactct tctgtctgtt gcagggtggtt tctggctggg | 5760 |
| | tatagaccaa gaaggagctg agggaaacct gtccctgacg ggccacctct tcggggtgct | 5820 |
| | ggccagcctc tgcgtctccc tcaatgccat ctataccaag aaggtgctcc ctgcagtaga | 5880 |
| 30 | ccacagtatc tggcgcctaa ccttctataa caatgtcaat gcctgcgtgc tcttcttgcc | 5940 |
| | cctgaigata gtgtctggcg agctccgtgc cctcctggcc ttactcactc tgagcagtgc | 6000 |
| | ccacttctgg ctcatgatga cgctgggtgg cctgtttggc tttgccatcg gctatgtgac | 6060 |
| | aggactgcag atcaaatca ccagtcacct gaccataac gtgtcaggca cggccaaggc | 6120 |
| 35 | ctgtgcacag acagtgtctg ccgtgctcta ctacgaagag attaaagact tcctgtgtgtg | 6180 |
| | gacaagcaac ctgatggtgc tgggtggctc ctccgcctac acctgggtca ggggctggga | 6240 |
| | gatgcagaag acccaggagg accccagctc caaagatggt gagaagagt ctatcagggt | 6300 |
| 40 | gtgagctcct tcaggagacc agggctgagc tcgggtgggg cctgcccagc acggaaggct | 6360 |
| | tcccatagag cctactgggt atggccctga gcaataatgt ttacatcctt ctcaagaagac | 6420 |
| | catctaagaa gagccagggt ctttctctgat aatgtcagaa agctgccaaa tctcctgcct | 6480 |
| | gccccatctt ctagtcttgg gaaagcccta ccaggagtgg cacccttctt gcctcctct | 6540 |
| 45 | ggggcctgtc tacctccata tggctctctg ggttggggcc agctgcactc tttgggcact | 6600 |
| | ggactgatga agtgatgtct tactttctac acaagggaga tgggttgtga ccctactata | 6660 |
| | gctagttgaa gggagctgtg taaccccaca tctctggggc cctgggcagg tagcataata | 6720 |
| 50 | gctaggtgct attaacatca ataacacttc agactacctt tggaggcagt tgggagctga | 6780 |
| | gccgagagag agagatggcc attctgcctt cttctgtgtg gatgggtatg acagaccaac | 6840 |
| | tgtccatggg gtgactgaca cctccacact tcatattttc aactttagaa aagggggagc | 6900 |
| | cacacgtttt acagattaag tggagtgtg aatgcctcta cagcccctaa ccccactttc | 6960 |
| 55 | cctgcctggc ttctcttggc ccagaagggc caccatctg ttctccaaca cctgaccag | 7020 |

ctatctggct atactctctt tctgtactcc ctcccccttc cccccccat tagcctcctc 7080
 cccaacacct ccattctcag gcaggaagtg gggccactc agcctctgtt cccatctgct 7140
 5 tggacccttg agcctctcat gaaggtaggc ttatgttctc tgaggctggg gccggaggag 7200
 cgcactgatt ctgggagtta tcccatcagg ctctgtcac aaaatagcct aggccgtgtg 7260
 tctaagaaca gtggagggtg gcttataact gttctggggg cagcgaagcc cacatcaagg 7320
 10 tactcataga cccagtattt ctgaggaaac cctgtccac atcctcactt ggtaaaggca 7380
 cagataatct cctcaggcc tcttgtatag gaggactagc cctgggaggg ctccgcccca 7440
 tgacctgac accccaaagc cttaacaga aggattccaa catgaatttg gggacagaag 7500
 15 cactcagacc acgatgccca gcaccacacc ctctatcct cagggtagct gtcactgtcc 7560
 tagtccccctc tgtttggcct tttgtacct catttccttg gcgtcatgtt tgatgtcttt 7620
 gtctctttcg tgagaagatg gggaaaccat gtcagcctct gcttccgact tccatgggt 7680
 tctaataaag ttgggtgggc ctgatgccct gatttgtatg tgatttaata aaaaaaaaaat 7740
 20 ttttttaaaa ac 7752

<210> 100

<211> 4039

<212> DNA

<213> *Cricetulus griseus*

<400> 100

gaacttcacc caagccatgt gacaattgaa ggctgtaccc ccagacccta acatcttgga 60
 gccctgtaga ccaggagtg cttctggccg tggggtgacc tagctcttct accaccatga 120
 35 acaggggcccc tctgaagcgg tccaggatcc tgcgcatggc gctgactgga ggctccactg 180
 cctctgagga ggcagatgaa gacagcagga acaagccgtt tctgctgcgg gcgctgcaga 240
 tcgcgctggt cgtctctctc tactgggtca cctccatctc catggatttc ctcaacaagt 300
 40 acctgctgga cagccccctc ctgcagctgg ataccctat ctctgtcact ttctaccaat 360
 gcctgggtgac ctctctgctg tgcaagggcc tcagcactct ggccacctgc tgccctggca 420
 ccgttgactt ccccacctg aacctggacc ttaagggtgg ccgcagcgtg ctgccactgt 480
 cggtagtctt cattggcatg ataagtttca ataacctctg cctcaagtac gtaggggtgg 540
 45 ccttctacaa cgtggggcgc tcgctacca ccgtgttcaa tgtgcttctg tctacctgc 600
 tgctcaaaca gaccacttcc ttctatgccc tgctcacatg tggcatcatc attggtgagt 660
 ggggcccggg ggctgtggga gcaggatggg catcgaactg aagccctaaa ggtcaacact 720
 gtaggtacct ttacttactg tcccaggctc ctgcatcag cagttacagg aagagccctg 780
 50 tagaaaacaa ataacttctt tatggtcatt caacaagtta gggaccacag cagggtgaaa 840
 ataatgtag cagcaactac agcaaagatg gctctcgcca ctgcatgat taaaatgtgc 900
 cagggtactca gatcyagca ttggatccac attaaactca ctaatcccta ttacaaggta 960
 55 aaatatatcc gaattttaca gagggaaaac caaggcacag agaggctaag tagcttgacc 1020

| | | |
|----|--|------|
| | aggatcacac agctaataat cactgacata gctgggattt aaacataagc agttacctcc | 1080 |
| | atagatcaca ctatgaccac catgccactg ttccttctca agagttccag gatcctgtct | 1140 |
| 5 | gtccagttct ctttaaagag gacaacacat ctgacattgc taccttgagg taacatttga | 1200 |
| | aatagtgggt agacatatgt ttttaagttt attcttrect tttatgygtg tgtgtttggg | 1260 |
| | gggccaccac agtgtatggg tggagataag gggacaactt aagaattggt cctttctccc | 1320 |
| 10 | accacatggg tgctgaggtc tgaactcagg tcatcaggat tggcacaat ccctttaccc | 1380 |
| | actgagccat ttcactggtc caatatatgt gtgcttttaa gaggctttaa ctattttccc | 1440 |
| | agatgtgaat gtccgtctga tcattatccc cttttacccg gaagccctct gggagggtgcc | 1500 |
| 15 | atccctgtgg tcgtctgcat acaaatgggg aaactgcaac tcagagaaac aaggctactt | 1560 |
| | gccagggccc cacaagtaag ataggctggg atgccatccc agactggcca cactccctgg | 1620 |
| | cctgtgcttc aagccagttt actttgttcc tgcccatgg aagttagcat gttgcagtca | 1680 |
| | aacacaataa ctacaggcca aaagtgcctt taaattaaag tcagatgaac ttttaaacad | 1740 |
| 20 | ccagagctcc tcaactgcag gagttaacaac ctgattctgc aacctcttt gcagtgcccg | 1800 |
| | gtagtcatat gtagctagag gctcttggct aggacagcat gtgttaggaa acatctggcc | 1860 |
| | ctgagatcat tgaattgagt gactgctggg tgacaagac caaggcatcc gttccctgag | 1920 |
| 25 | agtccctggc aagcagcaat gtgacctca tttgtacct ctcaggttct ttatctgtcc | 1980 |
| | tgtttgacct acttagtctc ctctggtgtc tcagaggccc aggctgggta ctctggatgt | 2040 |
| | caggatcagg ccaatgcgca catctgccct agaaatgtcc ccctggttga gcagctctctg | 2100 |
| | aatccatcgg taaagggtct ggaccaggga ggagtcagat aaaaagctga cagcactggg | 2160 |
| 30 | ggactccatg gggaactccc acctgcccc acacatccat cctaagagaa ctggtattcc | 2220 |
| | ttgtttctc ttgtctctac aaggcaccct gggatccac ttcagtctcc cagccttgcc | 2280 |
| | agggttagag ggcatgagcc tccttgtggg gaatttagat gcaagaaggt acagtacta | 2340 |
| | gagaacctga gctcagatcc ccaaagtaac cagtacctga tagtgaggca gctgagaacc | 2400 |
| 35 | gcagcagcct gcctgagtgg ctgaactctg cggcctccgg aactggcccc aactgttggg | 2460 |
| | tctcctcttc ctctctctg tgagggaggg cccatctctg ataagtgtg tggggactct | 2520 |
| | agagttagga ggaggaggag caatctaagc aggccctact gagaagtcct tgctggcatg | 2580 |
| 40 | tggctgcctg aggagtacag actgggaaca cccatttgaa tgagtaaggt ttttctgaa | 2640 |
| | ggccatgggg agccacggag gaaaatcatt ttagttaca gacaaagagt agattggta | 2700 |
| | acatggggagc arggacatgg ccccaatttt cattgatgaa ggaaattgga actcragag | 2760 |
| | gttaagtaac ttctcccaa tagctcagct tcataatcac agaacagtca gagtctagat | 2820 |
| 45 | ctctctgatg cctgtgatgg tcctgccatt ccatgttct gatccctgtg gcatcagtaa | 2880 |
| | gcctctacct tgtgggaatg caggatctaa atgaagagag raagtgtctg ccccatgctg | 2940 |
| | tggcttgga agctatgcag gctctttgag cagagagtga cccacaagt aatagagtc | 3000 |
| 50 | tatgagactc aaagcaacat ccacccttaa gcagctctaa ccaatgctc acactgaggg | 3060 |
| | agccaaagcc aagttagagt cctgtgcttg cccaaggtca ctttgccctg ccctcctct | 3120 |
| | atagcaccgc tgttatctta tagccctcat tacagtgatt acaattataa ttagagaggt | 3180 |
| | aacagggcca cactgtcctt acacattccc ctgctagatt gtagctggga gagggggaga | 3240 |
| 55 | tgtagggtggc tgggggagtg ggagggaaga tgcagatttt cattctgggc tctactccct | 3300 |

EP 1 498 491 A1

| | | |
|----|--|------|
| | cagccatttt ttggtgtggg agttagactt tggatatgtt gatgatgagg taagggccac | 3360 |
| | agaacagtct gaactgtggt atcagaatcc tgtccctctc cctctctcct catccctctt | 3420 |
| 5 | caccttgatc ctccctctgtc tgctacaggt ggtttctggc tgggtataga ccaagaggga | 3480 |
| | gctgagggca ccctgtccct cataggcacc atcttcgggg tgctggccag cctctgcgtc | 3540 |
| | tccctcaatg ccactctatac caagaagggt ctcacagcag tggacaacag catctggcgc | 3600 |
| 10 | ctaaccttct ataacaatgt caatgcctgt gtgctcttct tggccctgat ggttctgctg | 3660 |
| | ggtgagctcc gtgcccctct tgactttgct catctgtaca gtgcccactt ctggctcatg | 3720 |
| | atgacgctgg gtggcctctt cggctttgcc attggctatg tgacaggact gcagatcaaa | 3780 |
| 15 | ttcaccagtc ccttgaccca caatgtatca ggcacagcca aggcctgtgc gcagacagtg | 3840 |
| | ctggccgtgc tctactatga agagactaag agcttcctgt ggtggacaag caacctgatg | 3900 |
| | gtgctgggtg gctcctcagc ctatacctgg gtcaggggct gggagatgca gaagacccaa | 3960 |
| | gaggacccca gctccaaaga ggtgagaag agtgctattg ggtgtgagc ttcttcaggg | 4020 |
| 20 | acctgggact gaaccaag | 4039 |

25

30

35

40

45

50

55

Claims

1. A method for enhancing a binding activity of an antibody composition to Fc γ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule.
2. The method according to claim 1, wherein the modification of a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule is to bind a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain to the Fc region in the antibody molecule.
3. The method according to claim 1 or 2, wherein the sugar chain is synthesized by a cell in which the activity of a protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is decreased or deleted.
4. The method according to claim 3, wherein the protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is selected from the group consisting of the following (a), (b) and (c):
 - (a) a protein relating to synthesis of an intracellular sugar nucleotide, GDP-fucose;
 - (b) a protein relating to modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain;
 - (c) a protein relating to transport of an intracellular sugar nucleotide, GDP-fucose to the Golgi body.
5. The method according to claim 3 or 4, wherein the cell is resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain.
6. The method according to any one of claims 3 to 5, wherein the cell is resistant to at least one lectin selected from the group consisting of the following (a) to (d):
 - (a) a *Lens culinaris* lectin;
 - (b) a *Pisum sativum* lectin;
 - (c) a *Vicia faba* lectin;
 - (d) an *Aleuria aurantia* lectin.
7. The method according to any one of claims 3 to 6, wherein the cell is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.
8. The method according to any one of claims 3 to 7, wherein the cell is selected from the group consisting of the following (a) to (i):
 - (a) a CHO cell derived from a Chinese hamster ovary tissue;
 - (b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 cell;
 - (c) a BHK cell derived from a Syrian hamster kidney tissue;
 - (d) a mouse myeloma cell line NS0 cell;
 - (e) a mouse myeloma cell line SP2/0-Ag14 cell;
 - (f) a hybridoma cell;
 - (g) a human leukemic cell line Namalwa cell;
 - (h) an embryonic stem cell;
 - (i) a fertilized egg cell.
9. The method according to any one of claims 1 to 8, wherein the antibody molecule is selected from the group consisting of the following (a) to (d):
 - (a) a human antibody;
 - (b) a humanized antibody;
 - (c) an antibody fragment comprising the Fc region of (a) or (b);
 - (d) a fusion protein comprising the Fc region of (a) or (b).

10. The method according to any one of claims 1 to 9, wherein the antibody molecule belongs to an IgG class.
11. The method according to any one of claims 1 to 10, wherein, in the complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule, the ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is 20% or more of total complex *N*-glycoside-linked sugar chains.
12. A method for enhancing an antibody-dependent cell-mediated cytotoxic activity of an antibody composition, which comprises using the method according to any one of claims 1 to 11.
13. A process for producing an antibody composition having an enhanced binding activity to Fc γ receptor IIIa, which comprises modifying a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule.
14. The process according to claim 13, wherein the modification of a complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule is to bind a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in the complex *N*-glycoside-linked sugar chain to the Fc region in the antibody molecule.
15. The process according to claim 13 or 14, wherein the sugar chain is synthesized by a cell in which the activity of a protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is decreased or deleted.
16. The process according to claim 15, wherein the protein relating to modification of a sugar chain in which fucose is bound to *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain is selected from the group consisting of the following (a), (b) and (c):
- (a) a protein relating to synthesis of an intracellular sugar nucleotide, GDP-fucose;
 - (b) a protein relating to modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain;
 - (c) a protein relating to transport of an intracellular sugar nucleotide, GDP-fucose to the Golgi body.
17. The process according to claim 15 or 16, wherein the cell is resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain.
18. The process according to any one of claims 15 to 17, wherein the cell is resistant to at least one lectin selected from the group consisting of the following (a) to (d):
- (a) a *Lens culinaris* lectin;
 - (b) a *Pisum sativum* lectin;
 - (c) a *Vicia faba* lectin;
 - (d) an *Aleuria aurantia* lectin.
19. The process according to any one of claims 15 to 18, wherein the cell is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.
20. The process according to any one of claims 15 to 19, wherein the cell is selected from the group consisting of the following (a) to (i):
- (a) a CHO cell derived from a Chinese hamster ovary tissue;
 - (b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 cell;
 - (c) a BHK cell derived from a Syrian hamster kidney tissue;
 - (d) a mouse myeloma cell line NS0 cell;
 - (e) a mouse myeloma cell line SP2/0-Ag14 cell;
 - (f) a hybridoma cell;
 - (g) a human leukemic cell line Namalwa cell;
 - (h) an embryonic stem cell;

(i) a fertilized egg cell.

21. The process according to any one of claims 13 to 20, wherein the antibody molecule is selected from the group consisting of the following (a) to (d):

- (a) a human antibody;
- (b) a humanized antibody;
- (c) an antibody fragment comprising the Fc region of (a) or (b);
- (d) a fusion protein comprising the Fc region of (a) or (b).

22. The process according to any one of claims 13 to 21, wherein the antibody molecule belongs to an IgG class.

23. The method according to any one of claims 13 to 22, wherein, in the complex *N*-glycoside-linked sugar chain which is bound to the Fc region of an antibody molecule, the ratio of a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond is 20% or more of total complex *N*-glycoside-linked sugar chains.

24. A process for producing an antibody composition having an increased antibody-dependent cell-mediated cytotoxic activity, which comprises using the process according to claim 12.

25. An antibody composition produced by the process according to any one of claims 13 to 24.

26. A method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among total complex *N*-glycoside-linked sugar chains bound to the Fc region in an antibody composition, which comprises: reacting an antigen with a tested antibody composition to form a complex of the antigen and the antibody composition; contacting the complex with an Fc γ receptor IIIa to measure the binding activity to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

27. A method for detecting the antibody-dependent cell-mediated cytotoxic activity in an antibody composition, which comprises: reacting an antigen with a tested antibody composition to form a complex of the antigen and the antibody composition; contacting the complex with an Fc γ receptor IIIa to measure the binding activity to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

28. A method for detecting the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain among total complex *N*-glycoside-linked sugar chains bound to the Fc region in an antibody composition, which comprises: contacting a tested antibody composition with a Fc γ receptor IIIa to measure the binding activity of the antibody composition to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

29. A method for detecting the antibody-dependent cell-mediated cytotoxic activity in an antibody composition, which comprises: contacting a tested antibody composition with a Fc γ receptor IIIa to measure the binding activity of the antibody composition to the Fc γ receptor IIIa; and detecting the ratio of a sugar chain in a standard antibody composition with a standard curve showing the binding activity to the Fc γ receptor IIIa.

30. An Fc fusion protein composition produced by using a cell resistant to a lectin which recognizes a sugar chain structure in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain.

31. The Fc fusion protein composition according to claim 30, wherein the cell is selected from the group consisting of the following (a), (b) and (c):

- (a) an enzyme protein relating to synthesis of an intracellular sugar nucleotide, GDP-fucose;
- (b) an enzyme protein relating to modification of a sugar chain in which 1-position of fucose is bound to 6-position of *N*-acetylglucosamine in the reducing end through α -bond in a complex *N*-glycoside-linked sugar chain;
- (c) a protein relating to transport of an intracellular sugar nucleotide, GDP-fucose to the Golgi body,

wherein the activity of the protein is decreased or deleted.

32. The Fc fusion protein composition according to claim 30 or 31, wherein the cell is resistant to at least one lectin selected from the group consisting of the following (a) to (d):

- (a) a *Lens culinaris* lectin;
- (b) a *Pisum sativum* lectin;
- (c) a *Vicia faba* lectin;
- (d) an *Aleuria aurantia* lectin.

33. The Fc fusion protein composition according to any one of claims 30 to 32, wherein the cell is a cell into which a gene encoding an Fc fusion protein is introduced.

34. The Fc fusion protein composition according to claim 33, wherein the Fc is derived from an IgG class of an antibody molecule.

35. The Fc fusion protein composition according to any one of claims 30 to 34, wherein the cell is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.

36. The Fc fusion protein composition according to any one of claims 30 to 35, wherein the cell is a mouse myeloma cell.

37. The Fc fusion protein composition according to claim 36, wherein the mouse myeloma cell is NSO cell or SP2/0-Ag14 cell.

38. The Fc fusion protein composition according to any one of claims 30 to 37, wherein the cell is selected from the group consisting of the following (a) to (g):

- (a) a CHO cell derived from a Chinese hamster ovary tissue;
- (b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 line;
- (c) a BHK cell derived from a Syrian hamster kidney tissue;
- (d) an antibody-producing hybridoma cell;
- (e) a human leukemic cell line Namalwa cell;
- (f) an embryonic stem cell;
- (g) a fertilized egg cell.

39. An Fc fusion protein composition comprising an Fc fusion protein having an complex *N*-glycoside-linked sugar chain at the Fc region of an antibody molecule, wherein the ratio of a sugar chain in which fucose is not bound to *N*-acetylglucosamine in the reducing end in the sugar chain is 20% or more of total complex *N*-glycoside-linked sugar chains which are bound to the Fc region in the composition.

40. The Fc fusion protein composition according to claim 39, wherein the sugar chain in which fucose is not bound is a sugar chain in which 1-position of fucose is not bound to 6-position of *N*-acetylglucosamine in the reducing end in the complex *N*-glycoside-linked sugar chain through α -bond.

41. The Fc fusion protein composition according to claim 39 or 40, wherein the antibody molecule belongs to an IgG class.

42. The Fc fusion protein composition according to any one of claims 30 to 41, wherein the Fc fusion protein composition is Fc-fused fibroblast growth factor-8.

43. A cell which produces the Fc fusion protein composition according to any one of claims 30 to 42.

44. The cell according to claim 43, which is selected from the group consisting of a yeast, an animal cell, an insect cell and a plant cell.

45. The cell according to claim 43 or 44, which is a mouse myeloma cell.

46. The cell according to claim 45, wherein the mouse myeloma cell is NSO cell or SP2/0-Ag14 cell.

47. The cell according to any one of claims 43 to 46, which is selected from the group consisting of the following (a) to (g):

- (a) a CHO cell derived from a Chinese hamster ovary tissue;
- (b) a rat myeloma cell line YB2/3HL.P2.G11.16Ag.20 line;
- (c) a BHK cell derived from a Syrian hamster kidney tissue;
- (d) an antibody-producing hybridoma cell;
- (e) a human leukemic cell line Namalwa cell;
- (f) an embryonic stem cell;
- (g) a fertilized egg cell.

48. A process for producing an Fc fusion protein composition, which comprises culturing the cell according to any one of claims 43 to 47 in a medium to form and accumulate an Fc fusion protein composition in the culture, and recovering the Fc fusion protein composition from the culture.

FIG. 1A




FIG. 1B




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6A





FIG. 6B



- ANTI-GD3 CHIMERIC ANTIBODY (50%)
- ANTI-GD3 CHIMERIC ANTIBODY (45%)
- ANTI-GD3 CHIMERIC ANTIBODY (29%)
- △— ANTI-GD3 CHIMERIC ANTIBODY (24%)
- ▲— ANTI-GD3 CHIMERIC ANTIBODY (13%)
- ×— ANTI-GD3 CHIMERIC ANTIBODY (7%)

FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26





FIG. 27






FIG. 29




FIG. 30




FIG. 31




FIG. 32A




FIG. 32B




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37





FIG. 38



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/04504

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.⁷ C12P21/08, C12N15/09, C12N5/10, C07K16/18, G01N33/50,
C07K16/46//A61K39/395, A61P9/00, A61P29/00, A61P31/04,
A61P31/12, A61P31/14, A61P35/00, A61P37/02, A61P37/04,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.⁷ C12P21/08, C12N15/09, C12N5/10, C07K16/18, G01N33/50,
C07K16/46

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
BIOSIS/MEDLINE/WPIDS (STN)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|--|
| X/Y/A | WO 00/61739 A1 (Kyowa Hakko Kogyo Co., Ltd.), 19 October, 2000 (19.10.00), & EP 117619 A1 & AU 3672800 A | 1-4, 9-16, 21-25, 27, 29, 31, 33, 34, 36, 39-43, 45, 48/ 5-8, 17-20, 30, 32, 35, 37, 38, 44, 46, 47/ 26, 28 |
| Y | Ripka J. et al., Two Chinese hamster ovary glycosylation mutants affected in the conversion of GDP-mannose to GDP-fucose, Arch.Biochem. Biophys., 1986, 249(2), pages 533 to 545 | 5-8, 17-20, 30, 32, 35, 38, 44, 47 |

☒ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not
considered to be of particular relevance

"E" earlier document but published on or after the international filing
date

"L" document which may throw doubts on priority claim(s) or which is
cited to establish the publication date of another citation or other
special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other
means

"P" document published prior to the international filing date but later
than the priority date claimed

"T" later document published after the international filing date or
priority date and not in conflict with the application but cited to
understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be
considered novel or cannot be considered to involve an inventive
step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be
considered to involve an inventive step when the document is
combined with one or more other such documents, such
combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
23 May, 2003 (23.05.03)

Date of mailing of the international search report
22 July, 2003 (22.07.03)

Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/04504

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|------------------------------------|
| X | Shitara K. et al., A new vector for the high level expression of chimeric antibodies in myeloma cells, <i>J.Immunol.Methods</i> , 1994, 167(1-2), pages 271 to 278 | 1-25, 30-45, 47, 48 |
| X | Davies J. et al., Expression of GnTIII in a recombinant anti-CD20 CHO production cell line: Expression of antibodies with altered glycoforms leads to an increase in ADCC through higher affinity for FC gamma RIII, <i>Biotechnol.Bioeng.</i> , 2001, 74(4), pages 288 to 294 | 1, 27, 29 |
| Y | Hackett J. Jr. et al., Recombinant mouse-human chimeric antibodies as calibrators in immunoassays that measure antibodies to <i>Toxoplasma gondii</i> , <i>J. Clin.Microbiol.</i> , 1998, 36(5), pages 1277 to 1284 | 37, 46 |
| Y | Elbashir SM. et al., Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells, <i>Nature</i> , 2001, 411(6836), pages 494 to 498 | 5-8, 17-20, 30, 32, 35, 38, 44, 47 |
| A | WO 97/27303 A (Toyobo Co., Ltd.), 31 July, 1997 (31.07.97), & US 6054304 A & EP 816503 A1 & JP 9-201191 A | 1-48 |
| A | Shields RL. et al., High resolution mapping of the binding site on human IgG1 for Fc gamma RI, Fc gamma RII, Fc gamma RIII, and FcRn and design of IgG1 variants with improved binding to the Fc gamma R, <i>J.Biol.Chem.</i> , 2001, 276(9), pages 6591 to 6604 | 1-48 |
| P,X | Shields RL. et al., Lack of fucose on human IgG1 N-linked oligosaccharide improves binding to human Fc gamma RIII and antibody-dependent cellular toxicity, <i>J.Biol.Chem.</i> , 2002 Jul., 277(30), pages 26733 to 26740 | 1-48 |
| P,X | SHINKAWA, T. et al., The absence of fucose but not the presence of galactose or bisecting N-acetylglucosamine of human IgG1 complex-type oligosaccharides shows the critical role of enhancing antibody-dependent cellular cytotoxicity, <i>J.Biol.Chem.</i> , 2003, 278(5), pages 3466 to 3473 | 1-48 |
| P,X | WO 02/31140 A1 (Kyowa Hakko Kogyo Co., Ltd.), 18 April, 2002 (18.04.02), & AU 9419801 A | 1-48 |

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP03/04504

Continuation of A. CLASSIFICATION OF SUBJECT MATTER
(International Patent Classification (IPC))

Int.Cl⁷ A61P37/08, A61P43/00, G01N33/15, (C12P21/08, C12R1:91),
(C12N5/10, C12R1:91)

(According to International Patent Classification (IPC) or to both national
classification and IPC)

Continuation of B. FIELDS SEARCHEDMinimum Documentation Searched (International Patent Classification (IPC))

Int.Cl⁷

Minimum documentation searched (classification system followed by
classification symbols)